



# ANNUAL DANISH INFORMATIVE INVENTORY REPORT TO UNECE

Emission inventories from the base year of the protocols to year 2013

---

Scientific Report from DCE - Danish Centre for Environment and Energy

No. 145

2015



AARHUS  
UNIVERSITY

DCE - DANISH CENTRE FOR ENVIRONMENT AND ENERGY



*[Blank page]*

# ANNUAL DANISH INFORMATIVE INVENTORY REPORT TO UNECE

Emission inventories from the base year of the protocols to year 2013

---

Scientific Report from DCE – Danish Centre for Environment and Energy

No. 145

2015

Ole-Kenneth Nielsen  
Marlene S. Plejdrup  
Morten Winther  
Mette Hjorth Mikkelsen  
Malene Nielsen  
Steen Gyldenkærne  
Patrik Fauser  
Rikke Albrektsen  
Katja Hjelgaard  
Henrik G. Bruun  
Marianne Thomsen

Aarhus University, Department of Environmental Science



AARHUS  
UNIVERSITY

DCE – DANISH CENTRE FOR ENVIRONMENT AND ENERGY

## Data sheet

Series title and no.:	Scientific Report from DCE – Danish Centre for Environment and Energy No. 145
Title:	Annual Danish Informative Inventory Report to UNECE
Subtitle:	Emission inventories from the base year of the protocols to year 2013
Authors:	Ole-Kenneth Nielsen, Marlene S. Plejdrup, Morten Winther, Mette Hjorth Mikkelsen, Malene Nielsen, Steen Gyldenkærne, Patrik Fauser, Rikke Albrektsen, Katja Hjelgaard, Henrik G. Bruun, Marianne Thomsen
Institution:	Department of Environmental Science
Publisher:	Aarhus University, DCE – Danish Centre for Environment and Energy ©
URL:	<a href="http://dce.au.dk">http://dce.au.dk</a>
Year of publication:	March 2015
Editing completed:	March 2015
Financial support:	No external financial support
Please cite as:	Nielsen, O.-K., Plejdrup, M.S., Winther, M., Mikkelsen, M.H., Nielsen, M., Gyldenkærne, S., Fauser, P., Albrektsen, R., Hjelgaard, K., Bruun, H.G. & Thomsen, M. 2015. Annual Danish Informative Inventory Report to UNECE. Emission inventories from the base year of the protocols to year 2013. Aarhus University, DCE – Danish Centre for Environment and Energy, 484 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 145 <a href="http://dce2.au.dk/pub/SR145.pdf">http://dce2.au.dk/pub/SR145.pdf</a>
	Reproduction permitted provided the source is explicitly acknowledged
Abstract:	This report is a documentation report on the emission inventories for Denmark as reported to the UNECE Secretariat under the Convention on Long Range Transboundary Air Pollution due by 15 February 2015. The report contains information on Denmark's emission inventories regarding emissions of (1) SO <sub>x</sub> for the years 1980-2013, (2) NO <sub>x</sub> , CO, NMVOC and NH <sub>3</sub> for the years 1985-2013, (3) Particulate matter: TSP, PM <sub>10</sub> , PM <sub>2.5</sub> for the years 2000-2013, (4) Heavy Metals: Pb, Cd, Hg, As, Cr, Cu, Ni, Se and Zn for the years 1990-2013, (5) Polyaromatic hydrocarbons (PAH): Benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene and indeno(1,2,3-cd)pyrene, PCDD/F and HCB for the years 1990-2013. Further, the report contains information on background data for emissions inventory.
Keywords:	Emission Inventory; Emissions; Projections; UNECE; EMEP; LRTAP; NO <sub>x</sub> ; CO; NMVOC; SO <sub>x</sub> ; NH <sub>3</sub> ; TSP; PM <sub>10</sub> ; PM <sub>2.5</sub> ; Pb; Cd; Hg; As; Cr; Cu; Ni; Se; Zn; Polyaromatic hydrocarbons; Dioxin; Benzo(a)pyrene, Benzo(b)fluoranthene
Layout:	Ann-Katrine Holme Christoffersen
Front page photo:	Ann-Katrine Holme Christoffersen
ISBN:	978-87-7156-133-3
ISSN (electronic):	2245-0203
Number of pages:	484
Internet version:	The report is available in electronic format (pdf) at <a href="http://dce2.au.dk/pub/SR145.pdf">http://dce2.au.dk/pub/SR145.pdf</a>



# Contents

<b>Acknowledgements</b>	<b>7</b>
<b>Summary</b>	<b>8</b>
I Background information on emission inventories	8
II Trends in emissions	8
III Recalculations and Improvements	13
<b>Sammenfatning</b>	<b>14</b>
I Baggrund for emissionsopgørelser	14
II Udviklingen i emissioner	14
III Genberegninger og forbedringer	19
<b>1 Introduction</b>	<b>20</b>
1.1 Background information on emission inventories	20
1.2 A description of the institutional arrangement for inventory preparation	20
1.3 Brief description of the process of inventory preparation. Data collection and processing, data storage and archiving	21
1.4 Brief description of methodologies and data sources used	24
1.5 Key categories	30
1.6 Information on the Quality Control and Quality Assurance plan including verification and treatment of confidential issues where relevant	30
1.7 General uncertainty evaluation, including data on the overall uncertainty for the inventory totals	30
1.8 General assessment of the completeness	31
1.9 References	32
<b>2 Trends in Emissions</b>	<b>34</b>
2.1 Acidifying gases	34
2.2 Description and interpretation of emission trends by gas	35
2.3 Other air pollutants	37
<b>3 Energy (NFR sector 1)</b>	<b>45</b>
3.1 Overview of the sector	45
3.2 Stationary combustion (NFR sector 1A1, 1A2 and 1A4)	46
3.3 Transport and other mobile sources (NFR sector 1A2, 1A3, 1A4 and 1A5)	128
1.1 Fugitive emissions (NFR sector 1B)	207
<b>4 Industrial processes (NFR sector 2)</b>	<b>235</b>
4.1 Overview of the sector	235
4.2 Mineral products	237
4.3 Chemical industry	246
4.4 Metal production	253
4.5 Non-energy products from fuels and solvent use	264
4.6 Other product use	272
4.7 Other production	273

4.8	Other production, consumption, storage, transportation or handling of bulk products	274
4.9	QA/QC and verification	276
4.10	Uncertainty estimates	276
4.11	References	277
<b>5</b>	<b>Agriculture (NFR sector 3)</b>	<b>281</b>
5.1	Overview of the sector	281
5.2	References – sources of information	284
5.3	Manure management	287
5.4	Soils	299
5.5	Field burning of agricultural residues	309
5.6	Agriculture other	310
5.7	Uncertainties	311
5.8	Quality assurance and quality control (QA/QC)	315
5.9	Recalculations	316
5.10	Planned improvements	317
5.11	References	317
<b>6</b>	<b>Waste</b>	<b>320</b>
6.1	Solid waste disposal	320
6.2	Biological treatment of solid waste	320
6.3	Waste incineration	326
6.4	Wastewater handling	335
6.5	Other waste	335
6.6	Uncertainties and time series consistency	344
6.7	QA/QC and verification	346
6.8	Source-specific recalculations and improvements	347
6.9	Source-specific planned improvements	347
6.10	References	348
<b>7</b>	<b>Other and natural emissions</b>	<b>352</b>
<b>8</b>	<b>Reporting spatially distributed emissions on grid</b>	<b>353</b>
8.1	Background for reporting	353
8.2	Methods and data for disaggregation of emission data	354
8.3	Gridded emission data	359
8.4	References	362
<b>9</b>	<b>Recalculations and Improvements</b>	<b>363</b>
9.1	Energy	363
9.2	Industrial processes	367
9.3	Agriculture	370
9.4	Waste	370
<b>10</b>	<b>Projections</b>	<b>371</b>
10.1	References	371
<b>11</b>	<b>Adjustments</b>	<b>372</b>
11.1	Accepted adjustments	373
11.2	Application for adjustment	374
11.3	References	375
	<b>Annex 1 – Key category analysis</b>	<b>377</b>



<b>Annex 2 – Information on the energy balance</b>	<b>380</b>
<b>Annex 3A - Stationary combustion</b>	<b>381</b>
Annex 3A-1 Correspondence list for SNAP/CRF	382
Annex 3A-2 Fuel rate	385
Annex 3A-3 Default Lower Calorific Value (LCV) of fuels and fuel correspondance list	388
Annex 3A-4 Emission factors	392
Annex 3A-5 Implied emission factors for waste incineration plants and power plants combustion coal	424
Annex 3A-6 Large point sources	425
Annex 3A-7 Uncertainty estimates, 2013	430
Annex 3A-8 Emission inventory 2013 based on SNAP sectors	443
Annex 3A-9 Description of the Danish energy statistics	446
Annex 3A-10 Time-series 1980/1985-2013	453
Annex 3A-11 QA/QC for stationary combustion	455
<b>Annex 3B - Transport and other mobile sources</b>	<b>461</b>
List of content	461
Annex 3B-3: EU directive emission limits for road transportation vehicles	463
Annex 3B-11-9: Basis fuel consumption and emission factors, deterioration factors and transient factors for non road working machinery and equipment and recreational craft. Stock and activity data for certain specific types of machinery	468
<b>Annex 3C - Industrial processes (NFR 2)</b>	<b>479</b>
<b>Non-energy products from fuels and solvent use (NFR 2D)</b>	479
<b>Other product use (NFR 2G)</b>	479
<b>Annex 3D - Agriculture</b>	<b>480</b>
<b>Annex 3E - Waste</b>	<b>482</b>
<b>Annex 4 – Completeness and use of notation keys</b>	<b>483</b>
Not estimated categories	483
Mobile combustion	483
Industrial processes	483
Agriculture	483
Waste	483
Categories reported as IE (Included Elsewhere)	484

*[Blank page]*



## Acknowledgements

The work of compiling the Danish air pollutant emission inventory requires the input of many individuals, companies and institutions. The authors of this report would, in particular, like to thank the following for their valuable input in the work process:

- The Danish Energy Agency, in particular Jane Rusbjerg, Ali Zarnaghi and Kaj Stærkind for valuable discussions concerning energy balance data.
- The Danish Environmental Protection Agency, in particular Hans E. Jensen, Charlotte von Hessberg and Stine S. Justesen for valuable discussions concerning PRTR reporting, environmental accounts from companies and small combustion sources.
- DONG Energy A/S and Vattenfall A/S for providing detailed data on emissions at boiler level.
- Anette Holst, Statoil Refining Denmark A/S, for providing detailed data and information on calorific values and uncertainties related to processes at the refinery.
- Lis R. Rasmussen, A/S Danish Shell, Shell Refinery, for providing detailed data on emissions from the refinery.
- HMN Naturgas for providing detailed data on distribution of natural gas
- NGF Nature Energy Distribution A/S for providing detailed data on distribution of natural gas
- DTU Transport (Technical University of Denmark), in particular Thomas Jensen for valuable input and discussions on road transport fleet and mileage characterisation.
- DCA - Danish Centre for Food and Agriculture, Aarhus University, for valuable input on animals feed consumption and excretion based on the Danish Normative System and Peter Lund for information and discussion on methane conversion rate from livestock and emission particularly from sheep and lambs.
- Annette Vestergaard - Knowledge Centre for Agriculture, for discussions on actual farming practice regarding acidification of manure during application to soils.
- The Danish AgriFish Agency for providing unrestricted access to all agricultural data.

# Summary

## I Background information on emission inventories

### Annual report

This report is Denmark's Annual Informative Inventory Report (IIR) due March 15, 2015 to the UNECE-Convention on Long-Range Transboundary Air Pollution (LRTAP). The report contains information on Denmark's inventories for all years from the base years of the protocols to 2013.

The air pollutants reported under the LRTAP Convention are SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, dioxins/furans, HCB, PCBs and PAHs,.

The annual emission inventory for Denmark is reported in the Nomenclature for Reporting (NFR) 2014 format.

The issues addressed in this report are: trends in emissions, description of each NFR category, uncertainty estimates, recalculations, planned improvements and procedures for quality assurance and control. The structure of the report follows to the extent possible the proposed outline.

Information contained in this report is available to the public on the Danish Centre for Environment and Energy (DCE), Aarhus University's homepage:

<http://envs.au.dk/videnudveksling/luft/emissioner/emissioninventory/>

and this report and the NFR tables are available on the Eionet central data repository:

[http://cdr.eionet.europa.eu/dk/Air\\_Emission\\_Inventories/Submission\\_EMEP\\_UNECE](http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories/Submission_EMEP_UNECE)

### Responsible institute

DCE-Danish Centre for Environment and Energy, Aarhus University, is on behalf of the Danish Ministry of the Environment responsible for the annual preparation and submission to the UNECE-LRTAP Convention of the Annual Danish Emissions Report and the inventories in the NFR format. DCE participates in meetings under the UNECE Task Force on Emission Inventories and Projections and the related expert panels, where parties to the convention prepare the guidelines and methodologies on inventories.

## II Trends in emissions

### Acidifying gases

In 1990, the relative contribution in acid equivalents was almost equal for the three gases SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub>. In 2013, the most important acidification factor in Denmark is ammonia nitrogen and the relative contributions for SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 6 %, 36 % and 58 %, respectively. However, with regard



to long-range transport of air pollution, SO<sub>2</sub> and NO<sub>x</sub> are still the most important pollutants.

### **Sulphur dioxide (SO<sub>2</sub>)**

The main part of the SO<sub>2</sub> emission originates from combustion of fossil fuels, i.e. mainly coal and oil, in public power and district heating plants. From 1990 to 2013, the total emission decreased by 92 %. The large reduction is mainly due to installation of desulphurisation plants in public power and district heating plants and use of fuels with lower content of sulphur. Despite the large reduction of the SO<sub>2</sub> emissions, these plants make up 27 % of the total emission. Also emissions from industrial combustion plants, non-industrial combustion plants, other mobile sources and production of bricks and tiles are important. National sea traffic (navigation and fishing) contributes with about 11 % of the total SO<sub>2</sub> emission in 2013. This is due to the use of residual oil with high sulphur content.

### **Nitrogen oxide (NO<sub>x</sub>)**

The largest sources of emissions of NO<sub>x</sub> are road transport followed by other mobile sources and combustion in energy industries (mainly public power and district heating plants). The transport sector is the sector contributing the most to the emission of NO<sub>x</sub> and, in 2013, 42 % of the Danish emissions of NO<sub>x</sub> stems from road transport, national navigation, railways and civil aviation. Also emissions from national fishing and off-road vehicles contribute significantly to the NO<sub>x</sub> emission. For non-industrial combustion plants, the main sources are combustion of gas oil, natural gas and wood in residential plants. The emissions from energy industries have decreased by 78 % from 1990 to 2013. In the same period, the total emission decreased by 59 %. The reduction is due to the increasing use of catalyst cars and installation of low-NO<sub>x</sub> burners and denitrifying units in power plants and district heating plants.

### **Ammonia (NH<sub>3</sub>)**

The majority of emissions of NH<sub>3</sub> result from agricultural activities. Only a minor part originates from stationary combustion (2 %), road transport (2 %), industrial processes and product use (1 %) and waste (1 %). The share for road transport increased during the 1990'ties and early 2000's due to growing use of catalyst cars. In more recent years the share is again decreasing due to more advanced catalysts being implemented.

The major part of the emission from agriculture stems from livestock manure (51 %) and the largest losses of ammonia occur during the handling of the manure in animal housing systems. The second largest agricultural source is agricultural soils contributing 48 % in 2013; this is mainly emissions from application of mineral fertiliser, application of animal manure and emissions from growing crops.

The total ammonia emission decreased by 42 % from 1985 to 2013. Due to the action plans for the aquatic environment and the Ammonia Action Plan, a series of measures to prevent loss of nitrogen in agricultural production has been initiated. The measures have included demands for improved utilisation of nitrogen in livestock manure, a ban against field application of livestock manure in winter, prohibition of broad spreading of manure, requirements for establishment of catch crops, regulation of the number of livestock per hectare and a ceiling for the supply of nitrogen to crops. As a result, de-

spite an increase in the production of pigs and poultry, the ammonia emission has been reduced considerably.

### **Other air pollutants**

#### **Non-methane volatile organic compounds (NMVOC)**

The emissions of NMVOC originate from many different sources and can be divided into two main groups: incomplete combustion and evaporation. Road vehicles and other mobile sources such as national navigation vessels and off-road machinery are the main sources of NMVOC emissions from incomplete combustion processes. Road transportation vehicles are still the main contributors, even though the emissions have declined since the introduction of catalyst cars in 1990. The evaporative emissions mainly originate from the agricultural sector, use of solvents and the extraction, handling and storage of oil and natural gas. The total anthropogenic emissions have decreased by 44 % from 1990 to 2013, largely due to the increased use of catalyst cars and reduced emissions from use of solvents.

#### **Carbon monoxide (CO)**

Non-industrial combustion plants are the main source to the total CO emission. For the non-industrial sector, emissions from commercial/institutional sources have increased and emissions from agriculture/forestry/fishing sources have decreased from 1990 to 2013, while emissions from the residential sector have been fluctuating, but around the same level in 1990 and 2013.

Transport is the second largest contributor to the total CO emission in 2013, showing a decrease of 81 % from 1990 to 2013. The major transport source is passenger cars, which make up 58 % in 1990, but has decreased to 20 % in 2013. The main driver is the increase of catalyst cars.

In 1990 a law forbidding the burning of agricultural crop residues on fields was implemented, which caused a significant reduction in CO emission.

The total CO emission decreased further by 55 % from 1990 to 2013, largely because of decreasing emissions from road transportation.

#### **Particulate Matter (PM)**

The particulate matter (PM) emission inventory has been reported for the years 2000 onwards. The inventory includes the total emission of particles Total Suspended Particles (TSP), emission of particles smaller than 10 µm (PM<sub>10</sub>) and emission of particles smaller than 2.5 µm (PM<sub>2.5</sub>).

The largest PM<sub>2.5</sub> emission sources are residential plants (63 %), road transport (9 %) and other mobile sources (8 %). Emissions from residential plants have increased by 32 % from 2000 to 2007, followed by a decrease of 32 % from 2007 to 2013. For other mobile sources, the most important sources are off-road vehicles and machinery in the industrial sector and in the agricultural/forestry sector (35 % and 33 %, respectively). For the road transport sector, exhaust emissions account for the major part (57 %) of the emissions. The PM<sub>2.5</sub> emission decreased by 11 % from 2000 to 2012 as the increasing wood consumption in the residential sector has been counterbalanced by decreasing emissions for the remaining sectors, the most important being the transport sector.

The largest TSP emission sources are agriculture and non-industrial combustion (74 % and 18 % of total TSP emission in 2013, respectively). Residential plants is the largest source in the non-industrial combustion sector, making up 16 % of the national total TSP emission in 2013. The TSP emissions from transport are also important and include both exhaust emissions and the non-exhaust emissions from brake and tyre wear and road abrasion. The non-exhaust emissions account for 68 % of the TSP emission from road transport in 2013.

### **Black carbon (BC)**

The black carbon (BC) emission inventory is reported for the years 2000 onwards. The main sources are residential plants and road transport contributing 38 % and 19 % in 2013, respectively. From 2000 to 2013 the total BC emission decreases by 24 %. BC emissions from non-industrial plants have increased by 43 % from 2000 to 2007, followed by a decrease of 30 % from 2007 to 2013. The trend for non-industrial combustion is mainly controlled by the trend for the wood consumption in the residential sector. BC emissions from the transport sector decrease by 49 % from 2000 to 2013, mainly due to implementing of new EURO norms and improved technology. An important factor is the use of particle filters for heavy duty vehicles and personal cars, which reduce the BC emission effectively. BC emissions from fugitive emissions from fuels, which are mainly due to storage of coal, decrease by 23 % from 2000 to 2013, in accordance with the decrease of the coal consumption in electricity and heat production.

### **Heavy metals**

In general, the most important sources of heavy metal emissions are combustion of fossil fuels and waste. The heavy metal emissions have decreased substantially in recent years, except for Cu. The reductions span from 19 % to 91 % for Zn and Pb, respectively. The reason for the reduced emissions is mainly increased use of gas cleaning devices at power and district heating plants (including waste incineration plants). The large reduction in the Pb emission is due to a gradual shift towards unleaded gasoline, the latter being essential for catalyst cars. The major source of Cu is automobile tyre and break wear (93 % in 2013) and the 30 % increase from 1990 to 2013 owe to increasing mileage.

### **Cadmium (Cd)**

The main sources of emissions of cadmium (Cd) to air are mainly combustion of wood, wood waste and municipal waste. Non-industrial combustion contributes 73 % in 2013, of which 97 % comes from residential plants. Emissions from residential plants have increased by 225 % from 1990 to 2013 due to increasing wood consumption. Emissions from energy industries, manufacturing industries and construction, and industrial processes have decreased by 88 % from 1990 to 2013. The decreasing emission from energy industries are related to the decreasing combustion of coal. In the transport sector emissions from passenger cars is the main source contributing with 58 % of the sectoral emission in 2013.

### **Mercury (Hg)**

The largest sources of mercury (Hg) emissions to air are waste incineration and coal combustion in energy industries. Due to improved flue gas cleaning and decreasing coal combustion the emissions from Energy industries decreased by 76 % from 1990-2000. The trend has continued in the following years and the corresponding decrease from 1990-2013 is 92 %. Non-

industrial combustion is dominated by wood combustion in residential plants while the main contributions to emissions from manufacturing industries and construction are food processing, beverages and tobacco, and non-metallic minerals. The variations in emissions from industrial processes owe to shut down in 2002 followed by re-opening and a second shut down in 2005 of the only Danish electro-steelwork.

#### **Lead (Pb)**

The main lead (Pb) emission sources are transport, waste, non-industrial combustion and industrial processes. In earlier years combustion of leaded gasoline was the major contributor to Pb emissions to air but the shift toward use of unleaded gasoline for transport have decreased the Pb emission from transport by 94 % from 1990-2013. The trend in the Pb emission from non-industrial combustion from 1990 to 2013 is almost constant. In the non-industrial combustion sector the dominant source is wood combustion in residential plants, which has been increasing from 1990 to 2013, but counter-balanced by decreasing emissions from stationary combustion in commercial/institutional and in agriculture/forestry/fishing. The decreasing emission from Energy industries (96 % from 1990 to 2013) is caused by the decreasing coal combustion and more efficient particle abatement.

#### **Polycyclic aromatic hydrocarbons (PAHs)**

The present emission inventory for polycyclic aromatic hydrocarbons (PAH) includes four PAHs: benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene and indeno(1,2,3-cd)pyrene. Benzo(b)fluoranthene and Benzo(a)pyrene contribute the major PAH emission by 34 % and 32 %, respectively in 2013. The most important source of PAHs emissions is combustion of wood in the residential sector making up 68 % of the total emission in 2013. The increasing emission trend is due to increasing combustion of wood in the residential sector. The PAH emission from combustion in residential plants has increased by 47 % from 1990 to 2013.

#### **Dioxins and furans**

The major part of the dioxin emission owes to wood combustion in the residential sector, mainly in wood stoves and ovens without flue gas cleaning. Wood combustion in residential plants accounts for 57 % of the national dioxin emission in 2013. The contribution to the total dioxin emission from the waste sector (26 % in 2013) owes to accidental fires, especially building fires. The emissions of dioxins from energy industries mainly owe to the combustion of biomass as wood, wood waste and to a less extend agricultural waste

#### **Hexachlorobenzene (HCB)**

Stationary combustion accounts for 51 % of the estimated national hexachlorobenzene (HCB) emission in 2013. This owes mainly to combustion of municipal solid waste in heating and power plants. Transport is an important source, too, and has increased by 56 % since 1990 due to increasing diesel consumption. The HCB emission from stationary plants has decreased 74 % since 1990 mainly due to improved flue gas cleaning in MSW incineration plants. The emission from agriculture was very high in the early 1990'ties due to the use of pesticides containing impurities of HCB. The HCB emission from agriculture decreased by 94 % from 1990 to 1994 and by 99 % from 1990 to 2013, causing the share of HCB emission from agriculture to drop from 67 % in 1990 to 5 % in 2013.

### **Polychlorinated biphenyls (PCBs)**

Transport accounts for 60 % of the estimated national polychlorinated biphenyls (PCBs) emission in 2013. This owes mainly to combustion of diesel in road transport. The emission from transport has decreased by 73 % since 1990 due to the phase out of leaded gasoline, which has a high PCBs emission factor. This has led to diesel fuel use being the most important source of PCBs emissions from transport in later years. The emission from manufacturing industries and non-industrial combustion is dominated by diesel fuel used in non-road machinery.

### **III Recalculations and Improvements**

In general, considerable work is being carried out to improve the inventories. Investigations and research carried out in Denmark and abroad produce new results and findings which are given consideration and, to the extent which is possible, are included as the basis for emission estimates and as data in the inventory databases. Furthermore, the updates of the EMEP/EEA Guidebook, and the work of the Task Force on Emission Inventories and its expert panels are followed closely in order to be able to incorporate the best scientific information as the basis for the inventories.

The implementation of new results in inventories is made in a way so that improvements, as far as possible, better reflect Danish conditions and circumstances. This is in accordance with good practice. Furthermore, efforts are made to involve as many experts as possible in the reasoning, justification and feasibility of implementation of improvements.

In improving the inventories, care is taken to consider implementation of improvements for the whole time series of inventories to make it consistent. Such efforts lead to recalculation of previously submitted inventories. This submission includes recalculated inventories for the whole time series. A description of the recalculations is provided in Chapter 9 and more detail can be found in the sectoral chapters of this report. For sector specific planned improvements please also refer to the relevant sectoral chapters.



# Sammenfatning

## I Baggrund for emissionsopgørelser

### Årlig rapport

Denne rapport er Danmarks årlige rapport om emissionsopgørelser sendt til UNECE-konventionen om langtransporteret grænseoverskridende luftforurening (LRTAP) 15. marts 2015. Rapporten indeholder oplysninger om Danmarks opgørelser for alle år fra basisårene for protokollerne til 2013.

Luftforureningskomponenterne der rapporteres til LRTAP-konventionen er SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2,5</sub>, As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, dioxiner/furaner, HCB, PCBs og PAH.

Den årlige emissionsopgørelse for Danmark rapporteres i NFR 2013-formatet.

Emnerne behandlet i rapporten er: Udvikling i emissioner, beskrivelse af hver NFR-kategori, usikkerheder, genberegninger, planlagte forbedringer og procedure for kvalitetssikring og -kontrol. Strukturen i rapporten følger, så vidt muligt, den foreslåede disposition.

Informationer fra denne rapport er tilgængelige for offentligheden på Aarhus Universitets hjemmeside:

<http://envs.au.dk/videnudveksling/luft/emissioner/emissioninventory/>

Den fulde rapport samt NFR-skemaer er tilgængelige på Eionets hjemmeside:

[http://cdr.eionet.europa.eu/dk/Air\\_Emission\\_Inventories/Submission\\_EMEP\\_UNECE](http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories/Submission_EMEP_UNECE)

### Ansvarlig institution

DCE – Nationalt Center for Miljø og Energi, Aarhus Universitet, er på vegne af Miljøministeriet ansvarlig for udarbejdelse af den årlige danske emissionsrapport og opgørelserne i NFR. DCE deltager i møder under UNECEs arbejdsgruppe for emissionsopgørelser og –fremskrivninger samt ekspertpaneler, hvor parter i konventionen udarbejder retningslinjer og metoder for emissionsopgørelserne.

## II Udviklingen i emissioner

### Forsurende gasser

I 1990 var det relative bidrag af syreækvivalenter næsten ens for de tre gasarter. I 2013 var ammoniak den vigtigste forsurende faktor i Danmark og de relative bidrag for SO<sub>2</sub>, NO<sub>x</sub> og NH<sub>3</sub> var på henholdsvis 6 %, 36 % og 58 %. Med hensyn til langtransporteret luftforurening er det dog stadig SO<sub>2</sub> og NO<sub>x</sub>, der er de vigtigste forureningskomponenter.

### **Svovldioxid (SO<sub>2</sub>)**

Hovedparten af SO<sub>2</sub>-emissionerne stammer fra forbrænding af fossile brændsler, dvs. primært kul og olie, på kraftværker, kraftvarmeværker og fjernvarmeværker. Fra 1990 til 2013 er den totale udledning reduceret med 92 %. Den store reduktion er primært opnået gennem installation af afsvovlingsanlæg på kraftværker og fjernvarmeværker og brug af brændsler med lavere svovlindhold. Trods den store reduktion er disse værker kilde til 27 % af den samlede udledning. Også emissioner fra industrielle forbrændingsanlæg, ikke-industrielle forbrændingsanlæg, andre mobile kilder samt teglværker og produktion af ekspanderede lerprodukter er væsentlige bidragsydere til emissionen. National søfart (sejlads og fiskeri) bidrager med 13 % af den totale SO<sub>2</sub>-emission. Dette skyldes brug af fuelolie med et højt svovlindhold.

### **Kvælstofite (NO<sub>x</sub>)**

Den største kilde til emissioner af NO<sub>x</sub> er transportsektoren efterfulgt af andre mobile kilder og forbrænding i energisektoren (hovedsageligt kraftværker og fjernvarmeværker). Transportsektoren er den sektor, der bidrager mest til udledningen af NO<sub>x</sub>, og i 2013 stammede 42 % af de danske NO<sub>x</sub>-emissioner fra vejtransport, national søfart, jernbaner og civil luftfart. Også emissioner fra nationalt fiskeri og off-road-køretøjer (entreprenør-, landbrugsmaskiner, m.m.) bidrager betydeligt til NO<sub>x</sub>-emissionen. For ikke-industrielle forbrændingsanlæg er de primære kilder forbrænding af gasolie, naturgas og træ i husholdninger. Emissionerne fra kraftværker og fjernvarmeværker er faldet med 78 % fra 1990 til 2013. I samme periode er den totale emission faldet med 59 %. Reduktionen skyldes øget brug af katalysatorer i biler og installation af lav-NO<sub>x</sub>-brændere og de-NO<sub>x</sub>-anlæg på kraftværker og fjernvarmeværker.

### **Ammoniak (NH<sub>3</sub>)**

Hovedparten af emissioner af NH<sub>3</sub> stammer fra aktiviteter i landbruget. Kun en mindre del skyldes stationær forbrænding (2 %), vejtransport (2 %), industrielle processer og produktanvendelse (1 %) og affald (1 %). Andelen fra transporten var stigende gennem 1990'erne og i starten af 2000'erne pga. den øgede brug af biler med katalysator. I de senere år er andelen igen faldet på grund af mere implementeringen af mere effektive katalysatorer.

Hovedparten af emissionen fra landbruget stammer fra husdyrgødning (51 %), og de største tab af ammoniak optræder under håndtering af gødningen i stalden. Den næststørste kilde indenfor landbrug er landbrugsjorde som bidrager med 48 % i 2013. Emissionen stammer hovedsageligt fra anvendelse af handelsgødning, udbringning af husdyrgødning samt emissioner fra voksende afgrøder.

Den totale ammoniakemission er faldet 42 % fra 1985-2013. Dette skyldes implementeringen af vandmiljøplaner og ammoniakhandlingsplanen som introducerede en række tiltag for at mindske kvælstoftabet i landbruget. Tiltagene har inkluderet krav om forbedret udnyttelse af kvælstof i husdyrgødning, et forbud mod udbringning af husdyrgødning om vinteren, forbud mod bredspredning af gødning, regler for plantning af efterafgrøder, regulering af antallet af tilladte dyr pr. hektar og et loft for gødningsanvendelsen for afgrøder. På trods af en stigning i produktionen af svin og fjerkræ, så er emissionen faldet betydeligt.

## Anden luftforurening

### Flygtige organiske forbindelser (NMVOC)

Emissionen af flygtige organiske forbindelser ekskl. metan (NMVOC) stammer fra mange forskellige kilder og kan opdeles i to hovedgrupper: Ufuldstændig forbrænding og fordampning. Hovedkilderne til NMVOC-emissioner fra ufuldstændige forbrændingsprocesser er brændeovne, vejtrafik og andre mobile kilder, som national sejlads og ikke vejgående maskiner. Køretøjer til vejtransport er fortsat den største bidragsyder, selvom emissionerne er faldet siden introduktionen af biler med katalysator i 1990. Emissionerne fra fordampning stammer hovedsageligt fra landbrug, anvendelse af opløsningsmidler og udvinding, lagring og transport af olie og gas. De totale menneskeskabte emissioner er faldet med 44 % fra 1990 til 2013, primært som følge af øget brug af biler med katalysator og reducerede emissioner fra anvendelse af opløsningsmidler.

### Kulilte (CO)

Ikke-industriel forbrænding er den vigtigste kilde til CO-emissionen. Indenfor ikke-industriel forbrænding er emissionen fra handel & service steget og emissioner fra landbrug/skovbrug/gartneri er faldet fra 1990 til 2013, emissionen fra husholdninger har fluktueret gennem tidserien, men er på samme niveau i 2013 som i 1990.

Transport er den næst vigtigste kilde til CO-emission i 2013, på trods af et fald i emissionen på 81 % mellem 1990 og 2013. Den største bidragsyder til emissionen fra transport er personbiler, som udgjorde 58 % i 1990, mens andelen er faldet til 20 % i 2013. Driveren for faldet i andelen er implementeringen af lovpligtige katalysatorer og stadigt strengere emissionskrav.

I 1990 trådte et forbud mod markafbrænding i kraft, som medførte et markant fald i CO-emissionen fra 1989 til 1990.

Samlet set, så faldt emissionen af CO med 55 % fra 1990 til 2013, hovedsageligt drevet af faldet i emissioner fra transportsektoren.

### Partikler (PM)

Emissionsopgørelsen for partikler (Particulate Matter, forkortet PM) er blevet rapporteret for år 2000 og fremefter. Opgørelsen inkluderer den totale emission af partikler: Total Suspended Particles (TSP), emissionen af partikler mindre end 10 µm (PM<sub>10</sub>) og emissionen af partikler mindre end 2,5 µm (PM<sub>2,5</sub>).

De største kilder til PM<sub>2,5</sub>-emission er husholdninger (63 %), vejtrafik (9 %) og andre mobile kilder (8 %). Emissionen fra husholdninger steg med 32 % fra 2000 til 2007 efterfulgt af et fald på 32 % fra 2007 til 2013. For andre mobile kilder er offroad-køretøjer i industrien samt landbrugs- og skovbrugsmaskiner de vigtigste kilder (hhv. 35 % og 33 %). I transportsektoren tegner udstødningsemissioner sig for størstedelen (57 %). PM<sub>2,5</sub>-emissionen er faldet med 11 % fra 2000 til 2013, da det stigende træforbrug og dermed emissioner fra husholdninger modsvares fald i emissionen fra de øvrige sektorer især transportsektoren.

De største kilder til TSP-emission er landbrugssektoren og husholdningerne. TSP-emissionen fra transport er også vigtig og inkluderer både udstødningsemissioner og ikke-udstødningsrelaterede emissioner fra slid af brem-

ser, dæk og vej. De ikke-udstødningsrelaterede emissioner udgør 68 % af TSP-emissionen fra transport.

### **Sod (BC)**

Emissionsopgørelsen for sod (Black carbon – BC) er rapporteret for året 2000 og fremefter. De vigtigste kilder er husholdninger og vejtransport, der bidrager med henholdsvis 38 % og 19 % i 2013. Fra 2000 til 2013 er den samlede BC-emission faldet med 24 %. BC-emissionen fra ikke-industriel forbrænding steg med 43 % fra 2000 til 2007, men er efterfølgende faldet med 30 %. Udviklingen indenfor ikke-industriel forbrænding er domineret af udviklingen i træforbruget i husholdninger.

BC-emissionen fra transportsektoren er faldet med 49 % fra 2000 til 2013, hvilket skyldes implementeringen af nye EURO-normer og forbedret teknologi. En vigtig faktor er anvendelsen af partikelfiltre for lastbiler og personbiler, som effektivt begrænser udledningen af partikler og også BC.

BC-emissioner fra udvinding/lagring /transport af kul, olie og gas kommer hovedsageligt fra lagring af kul. Emissionen er fladet med 23 % fra 2000 til 2013 på grund af det faldende kulforbrug til produktion af el og varme.

### **Tungmetaller**

Generelt er de vigtigste kilder til emissioner af tungmetaller forbrænding af fossile brændsler og affald. Emissionerne af tungmetaller er med undtagelse af kobber, faldet betydeligt de seneste år. Reduktionerne spænder fra 19 % til 91 % for henholdsvis Zn og Pb. Årsagen til de reducerede emissioner er hovedsageligt den øgede brug af røggasrensning på kraftværker og fjernvarmeverker (inklusive affaldsforbrændingsanlæg). Den store reduktion i emissionen af Pb skyldes et løbende skift til fordel for blyfri benzin, som er nødvendigt for biler med katalysator. Den største kilde til emission af kobber er slid af køretøjers dæk og bremses (93 % i 2013). Emissionen herfra er steget 30 % fra 1990 til 2013 pga. en stigning i antal kørte kilometer.

### **Cadmium (Cd)**

Den største kilde til Cd-emissioner er forbrænding i energisektoren (hovedsageligt forbrænding af træ og husholdningsaffald). Emissionen fra ikke-industriel forbrænding bidrager med 73 % i 2013 hvoraf 97 % kommer fra husholdninger. Emissionen fra husholdninger er steget med 225 % fra 1990 til 2013 på grund af et stigende træforbrug. Emissioner fra kraftværker, fremstillingsvirksomhed og industrielle processer er faldet med 88 % fra 1990 til 2013. Den faldende emission fra kraftværker skyldes faldende forbrug af kul. Emissioner fra personbiler er den dominerende kilde i transportsektoren og udgør 58 % i 2013.

### **Kviksølv (Hg)**

Den største kilde til Hg-emission er forbrænding af affald og kul i energisektoren. Forbedret røggasrensning og faldende kulforbrug har medført et fald i emissionen fra energisektoren på 76 % fra 1990 til 2000. Denne udvikling er fortsat således at emissionen fra 1990 til 2013 samlet set er faldet med 92 %. Emissionen fra ikke-industriel forbrænding kan hovedsageligt tilskrives forbrænding af træ i husholdninger. Emissionen fra fremstillingsvirksomhed stammer hovedsageligt fra forbrænding i undersektorerne fødevarerindustri og mineralsk industri. Emissionerne fra industrielle processer varierer meget pga. lukning af elektrostålvalseværket i 2002 efterfulgt af genåbning og endnu en lukning i 2005.

### **Bly (Pb)**

Den vigtigste kilde til emission af bly er transport, affald, ikke-industriell forbrænding og industrielle processer. I tidligere år var den største kilde forbrænding af blyholdigt benzin, men overgangen til blyfri benzin i transportsektoren har medført et fald i bly-emissionen fra transport på 94 % fra 1990 til 2013. Udviklingen i emissionen fra ikke-industriell forbrænding har været stort set konstant fra 1990-2012. Forbrænding af træ i husholdningsanlæg er den største kilde til emission af bly fra ikke-industriell forbrænding, og denne emission har været stigende fra 1990 til 2013, men er modvirket af faldende emissioner fra handel & service og landbrug/skovbrug/fiskeri. Emissionen fra energifremstilling er faldet med 96 % fra 1990 til 2013 hovedsageligt pga. faldende forbrug af kul og forbedret røggasrensning.

### **Polycykliske aromatiske hydrocarboner (PAH'er)**

Den nuværende emissionsopgørelse for polycykliske aromatiske hydrocarboner (PAH'er) inkluderer de fire PAH'er: Benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene og indeno(1,2,3-cd)pyrene. Hovedparten af den samlede PAH-emission kan tilskrives benzo(b)fluoranthene og Benzo(a)pyrene der står for hhv. 34 % og 32 %. Den vigtigste kilde til emission af PAH er forbrænding af træ i husholdningerne, der udgør 68 % af den samlede PAH-emission i 2013. De stigende emissioner skyldes øget forbrænding af træ i brændeovne og kedler i husholdningerne. Emissionen fra stationær forbrænding i husholdninger er steget med 47 % fra 1990 til 2013.

### **Dioxiner og furaner**

Størstedelen af dioxinemissionen skyldes forbrænding i husholdninger, hovedsageligt forbrænding af træ i brændeovne og -kedler uden røggasrensning. Forbrænding af træ i stationære anlæg i husholdninger udgør 57 % af den nationale dioxin emission i 2013. Emissioner fra affaldssektoren udgør 26 % af den nationale total i 2013, og skyldes hovedsageligt brande i bygninger. Forbrænding af træ og halm er den største kilde til dioxin emission fra energifremstilling.

### **Hexachlorbenzen (HCB)**

Stationær forbrænding udgør 51 % af den samlede beregnede hexachlorbenzen(HCB)-emission i 2013. Emissionen stammer hovedsageligt fra forbrænding af affald i kraftvarmeværker og fjernvarmeværker. Transport er også en vigtig kilde, og emissionen er steget med 56 % siden 1990 pga. det stigende dieselforbrug. HCB-emissionen fra stationære anlæg er faldet med 74 % siden 1990 hovedsageligt pga. forbedret røggasrensning på affaldsforbrændingsanlæg. Emissionen fra landbrug var høj i starten af 1990'erne pga. anvendelse af pesticider, der indeholdt urenheder af HCB. HCB-emissionen fra landbrug faldt med 94 % fra 1990 til 1994 og med 99 % fra 1990 til 2013. Dette har medført, at landbrugets andel af den samlede emission er faldet fra 67 % i 1990 til 5 % i 2013.

### **Polychlorerede bifenyler (PCBs)**

Transport udgør 60 % af den samlede beregnede emission af polychlorerede bifenyler (PCBs) i 2013. Det største bidrag kommer fra forbrænding af diesel i vejtransport. Emissionen er faldet med 73 % siden 1990 pga. udfasningen af blyholdig benzin, som har en høj PCBs emissionsfaktor. Udfasningen af blyholdig benzin har medført, at diesel er blevet den vigtigste kilde til PCBs emission i de senere år. Emissionen fra fremstillingsvirksomhed & bygge/anlæg samt ikke-industriell forbrænding er domineret af diesel forbrænding i ikke-vejgående maskiner.



### **III Genberegninger og forbedringer**

Generelt pågår der et betydeligt arbejde med at forbedre emissionsopgørelserne. Nye undersøgelser og forskning fra Danmark og udlandet inkluderes så vidt muligt som basis for emissionsestimerne. Desuden følges arbejdet med opdateringer af EMEP/EEA Guidebook for emissionsopgørelser nøje, med henblik på at indarbejde de bedste videnskabelige informationer som basis for opgørelserne.

Opgørelserne opdateres løbende med ny viden, således at opgørelserne bedst mulig afspejler danske forhold. Ved forbedringer lægges vægt på at opdateringer omfatter hele tidsserier, for at sikre konsistente data. Disse tiltag medfører genberegning af tidligere indberettede opgørelser. Begrundelserne for genberegningerne er inkluderet i kapitel 9 samt i de enkelte sektorkapitler i denne rapport. For planlagte sektorspecifikke forbedringer henvises der til sektorkapitlerne.

# 1 Introduction

## 1.1 Background information on emission inventories

DCE (Danish Centre for Environment and Energy), Aarhus University is contracted by the Ministry of the Environment and the Ministry of Climate, Energy and Building to complete emission inventories for Denmark. Department of Environmental Science, Aarhus University is responsible for calculation and reporting of the Danish national emission inventory to EU and the UNFCCC (United Nations Framework Convention on Climate Change) and UNECE CLRTAP (Convention on Long Range Transboundary Air Pollution) conventions.

### 1.1.1 Annual report

According to the guidelines for reporting emission data under the Convention on Long-Range Transboundary Air Pollution (ECE/EB.AIR/125) prepared by the Task Force on Emission Inventories and Projections and approved by the Executive Body, countries that are parties to the UNECE-Convention on Long-Range Transboundary Air Pollution should annually submit an informative inventory report to the Secretariat. The current reporting Guidelines (ECE/EB.AIR/125) were accepted at the meeting of the Executive Body in December 2013. Due to a lack of resources, it has not been possible to incorporate the new elements of the reporting guidelines in this submission nor has the previous reporting guidelines (ECE/EB.AIR/75) been fully implemented.

This report is Denmark's Annual Informative Inventory Report due March 15, 2015. The report contains information on Denmark's inventories for all years from the base years of the protocols to 2013.

The annual emission inventory for Denmark is reported in the Nomenclature for Reporting (NFR) 2014 format.

The issues addressed in this report are: trends in emissions, description of each NFR category, uncertainty estimates, recalculations, planned improvements and procedures for quality assurance and control. The outline in annex V of the reporting guidelines is followed as far as possible.

This report and NFR tables are available to the public on the Danish emission inventory webpage:

<http://envs.au.dk/videnudveksling/luft/emissioner/emissioninventory/>

and on the Eionet central data repository:

[http://cdr.eionet.europa.eu/dk/Air\\_Emission\\_Inventories/Submission\\_EMEP\\_UNECE](http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories/Submission_EMEP_UNECE)

## 1.2 A description of the institutional arrangement for inventory preparation

DCE (Danish Centre for Environment and Energy, Aarhus University, is responsible for the annual preparation and submission to the UNECE-LRTAP Convention of the Annual Danish Emissions Report, and the inventories in

the NFR Format in accordance with the guidelines. DCE participates in meetings under the UNECE Task Force on Emission Inventories and Projections and the related expert panels where parties to the convention prepare the guidelines and methodologies on inventories. DCE is also responsible for estimating emissions for reporting to the NEC Directive, but the Danish EPA is responsible for the reporting.

The work concerning the annual emission inventory is carried out in cooperation with other Danish ministries, research institutes, organisations and companies:

Danish Energy Agency (DEA), Ministry of Climate, Energy and Building:  
Annual energy statistics in a format suitable for the emission inventory work and fuel-use data for the large combustion plants.

Danish Environmental Protection Agency (DEPA), Ministry of the Environment:  
Company reporting to e.g. the PRTR. Database on waste.

Statistics Denmark, Ministry of Economic Affairs and the Interior:  
Statistical yearbook, production statistics for manufacturing industries, agricultural statistics and import/export/production figures.

DCA (Danish Centre for Food and Agriculture), Aarhus University:  
Data on use of mineral fertiliser, feeding stuff consumption and nitrogen turnover in animals.

The Road Directorate, Ministry of Transport:  
Number of vehicles grouped in categories corresponding to the EU classification, mileage (urban, rural, highway), trip speed (urban, rural, highway).

Civil Aviation Agency of Denmark, Ministry of Transport:  
City-pair flight data (aircraft type and origin and destination airports) for all flights leaving major Danish airports.

Danish Railways, Ministry of Transport:  
Fuel-related emission factors for diesel locomotives.

Danish companies:  
Audited environmental reports and direct information gathered from producers and agency enterprises.

Formerly, the provision of data was on a voluntary basis, but now formal agreements are in place with the most important data suppliers, e.g. the Danish Energy Agency and DCA.

### **1.3 Brief description of the process of inventory preparation. Data collection and processing, data storage and archiving**

The background data (activity data and emission factors) for estimation of the Danish emission inventories is collected and stored in central databases located at DCE. The databases are in Access format and handled with software developed by the European Environmental Agency (EEA) and DCE. As input to the databases, various sub-models are used to estimate and ag-

gregate the background data in order to fit the format and level in the central databases. The methodologies and data sources used for the different sectors are described in Chapter 1.4 and Chapters 3 to 6. As part of the QA/QC plan (Chapter 1.5), the data structure for data processing support the pathway from collection of raw data to data compilation, modelling and final reporting.

For each submission, databases and additional tools and submodels are frozen together with the resulting NFR-reporting format. This material is placed on central institutional servers, which are subject to routine back-up services. Material, which has been backed up is archived safely. A further documentation and archiving system is the official journal for DCE, for which obligations apply to DCE, as a governmental institute. In this journal system, correspondence, both in-going and out-going, is registered, which in this case involves the registration of submissions and communication on inventories with the UNECE-LRTAP Secretariat, the European Commission, review teams, etc.

Figure 1.1 shows a schematic overview of the process of inventory preparation. The figure illustrates the process of inventory preparation from the first step of collecting external data to the last step, where the reporting schemes are generated for the UNFCCC and EU (in the CRF format (Common Reporting Format)) and to the United Nations Economic Commission for Europe/Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (UNECE/EMEP) (in the NFR format (Nomenclature For Reporting)). For data handling, the software tool is CollectER (Pulles et al., 1999) and for reporting the software tool is developed by DCE. Data files and programme files used in the inventory preparation process are listed in Table 1.1.




Figure 1.1 Schematic diagram of the process of inventory preparation.

Table 1.1 List of current data structure; data files and programme files in use.

QA/QC Level	Name	Application type	Path	Type	Input sources
4 store	CFR Submissions (UNFCCC and EU) NFR Submissions (UNECE and EU)	External report	U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_4a_Storage\	MS Excel, xml	CRF Reporter
3 process	CRF Reporter	Management tool	Working path: local (exe + mdb) machine Archive path: U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_3b_Processes		manual input and Importer2CRF
3 process	Importer2CRF	Help tool	U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_3b_Processes	MS Access	CRF Reporter, Collector2CRF and excel files
3 process	CollectER2CRF	Help tool	U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_3b_Processes	MS Access	NERIRep
2 process 3 store	NERIRep	Help tool	Working path: U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_3a_Storage	MS Access	CollectER databases; dk1972.mdb. .dkxxx.mdb
2 process	CollectER	Management tool	Working path: local (exe +mdb) machine Archive path: U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_2b_Processes		manual input
2 store	dk1980.mdb.dkxxx.mdb	Datastore	U:\ST_ENVS-LUFT-EMI\Inventory\AIY ears\8_AllSectors\Level_2a_Storage	MS Access	CollectER



## **1.4 Brief description of methodologies and data sources used**

Denmark's air emission inventories are based on the EMEP/EEA Guidebook, the CORINAIR methodology as well as the 2006 IPCC Guidelines (IPCC, 2006). CORINAIR (COOrdination of INformation on AIR emissions) is a European air emission inventory programme for national sector-wise emission estimations, harmonised with the IPCC guidelines. In 2013 a new edition of the EMEP/EEA Guidebook (EEA, 2013) was adopted for use by the EMEP Executive Body, the changes in the 2013 edition have only been reflected to a limited extent in this submission and the changes will be implemented depending on the availability of resources. In 2009 the EMEP/CORINAIR Guidebook changed name to the EMEP/EEA Guidebook (EEA, 2009). In this change the Guidebook switched nomenclature from SNAP to NFR.

The Danish inventory is prepared at the more detailed SNAP level rather than at the NFR level that is only suitable for reporting. To ensure estimates are as timely, consistent, transparent, accurate and comparable as possible, the inventory programme has developed calculation methodologies for most subsectors and software for storage and further data processing.

A thorough description of the CORINAIR inventory programme used for Danish emission estimations is given in Illerup et al. (2000). The CORINAIR calculation principle is to calculate the emissions as activities multiplied by emission factors. Activities are numbers referring to a specific process generating emissions, while an emission factor is the mass of emissions per unit activity. Information on activities to carry out the CORINAIR inventory is largely based on official statistics. The most consistent emission factors have been used, either as national values or default factors proposed by international guidelines.

A list of all subsectors at the most detailed level is given in Illerup et al. (2000) together with a translation between CORINAIR and IPCC codes for sector classifications.

### **1.4.1 The specific methodologies regarding stationary combustion**

Stationary combustion plants are part of the CRF emission sources 1A1 Energy Industries, 1A2 Manufacturing Industries and 1A4 Other sectors.

The Danish emission inventory for stationary combustion plants is based on the former CORINAIR system. The emission inventory for stationary combustion is based on activity rates from the Danish energy statistics. General emission factors for various fuels, plants and sectors have been determined. Some large plants, such as power plants, are registered individually as large point sources and plant-specific emission data are used.

The fuel consumption rates are based on the official Danish energy statistics prepared by the Danish Energy Agency (DEA). DCE aggregates fuel consumption rates to SNAP categories. The fuel consumption of the NFR category 1A4 Manufacturing industries and construction is disaggregated to subsectors according to the DEA data prepared and reported to Eurostat.

For each of the fuel and SNAP categories (sector and e.g. type of plant), a set of general emission factors has been determined. Some emission factors refer to the EMEP/EEA Guidebook and some are country specific and refer to

Danish legislation, Danish research reports or calculations based on emission data from a considerable number of plants.

A number of large plants, e.g. power plants, municipal waste incineration plants and large industrial plants are registered individually as large point sources. This enables use of plant-specific emission factors that refer to emission measurements stated in annual environmental reports. Emission factors of SO<sub>2</sub>, NO<sub>x</sub>, HM and PM are often plant specific.

Please refer to Chapter 3.2 and Annex 3A for further information on emission inventories for stationary combustion plants.

#### **1.4.2 Specific methodologies regarding transport**

The emissions from transport referring to SNAP category 07 (Road transport) and the sub-categories in 08 (Other mobile sources) are made up in the NFR categories; 1A3b (Road transport), 1A2f (Industry-other), 1A3a (Civil aviation), 1A3c (Railways), 1A3d (Navigation), 1A4c (Agriculture/forestry/-fisheries), 1A4a (Commercial/institutional), 1A4b (Residential) and 1A5 (Other).

An internal DCE model with a structure similar to the European COPERT IV emission model (EEA, 2013) is used to calculate the Danish annual emissions for road traffic. The emissions are calculated for operationally hot engines, during cold start and fuel evaporation. The model also includes the emission effect of catalyst wear. Input data for vehicle stock and mileage is obtained from DTU Transport and Statistics Denmark, and is grouped according to average fuel consumption and emission behaviour. For each group, the emissions are estimated by combining vehicle type and annual mileage figures with hot emission factors, cold:hot ratios and evaporation factors.

For air traffic, from 2001 onwards estimates are made on a city-pair level, using flight data provided by the Danish Civil Aviation Agency (CAA-DK) for flights between Danish airports and flights between Denmark and Greenland/Faroe Islands, and LTO and distance-related emission factors from the CORINAIR guidelines (Tier 2 approach). For previous years, the background data consists of LTO/aircraft type statistics from Copenhagen Airport and total LTO numbers from CAA-DK. With appropriate assumptions, consistent time series of emissions are produced back to 1990 and include the findings from a Danish city-pair emission inventory in 1998.

Off-road working machines and equipment are grouped in the following sectors: inland waterways (pleasure craft), agriculture, forestry, industry, and household and gardening. The sources for stock and operational data are various branch organisations and key experts. In general, the emissions are calculated by combining information on the number of different machine types and their respective load factors, engine sizes, annual working hours and emission factors.

The inventory for navigation consists of regional ferries, local ferries and other national sea transport (sea transport between Danish ports and between Denmark and Greenland/Faroe Islands). For regional ferries, the fuel consumption and emissions are calculated as a product of number of round trips per ferry route (Statistics Denmark), sailing time per round trip, share of round trips per ferry, engine size, engine load factor and fuel consump-

tion/emission factor. The estimates take into account the changes in emission factors and ferry specific data during the inventory period.

For the remaining navigation categories, the emissions are calculated simply as a product of total fuel consumption and average emission factors. For each inventory year, this emission factor average comprises the emission factors for all present engine production years, according to engine life times.

Please refer to Chapter 3.3 and Annex 3B for further information on emissions from transport.

### **1.4.3 The specific methodologies regarding fugitive emissions**

#### **Fugitive emissions from oil (1.B.2.a)**

Fugitive emissions from oil are estimated according to the methodology described in the Emission Inventory Guidebook (EEA, 2013). The sources include offshore extraction of oil and gas, onshore oil tanks, onshore and offshore loading of ships, and gasoline distribution. Activity data are given in the Danish Energy Statistics by the Danish Energy Agency. The emission factors are based on the figures given in the guidebook except in the case of onshore oil tanks and gasoline distribution where national values are included.

The VOC emissions from petroleum refinery processes cover non-combustion emissions from feed stock handling/storage, petroleum products processing, and product storage/handling. SO<sub>2</sub> is also emitted from non-combustion processes and includes emissions from product processing and sulphur-recovery plants. The emission calculations are based on information from the Danish refineries.

#### **Fugitive emissions from natural gas (1.B.2.b)**

Inventories of NMVOC emission from transmission and distribution of natural gas and town gas are based on annual environmental reports from the Danish gas transmission company and annual reports for the gas distribution companies. The annual gas composition is based on from the national transmission company.

#### **Fugitive emissions from flaring (1.B.2.c)**

Emissions from flaring offshore, in gas treatment and storage plants, and in refineries are included in the inventory. Emissions calculations are based on annual reports from the Danish Energy Agency and environmental reports from gas storage and treatment plants and the refineries. Calorific values are based on the reports for the EU ETS for offshore flaring, on annual gas quality data from Energinet.dk, and on additional data from the refineries. Emission factors are based on national studies and the EMEP/EEA Guidebook (EEA, 2013).

Please refer to Chapter 3.4 for further information on fugitive emissions from fuels.

#### **1.4.4 Specific methodologies regarding industrial processes and product use**

Energy consumption associated with industrial processes and the emissions thereof is included in the inventory for stationary combustion plants. This is due to the overall use of energy balance statistics for the inventory.

##### **Mineral industry**

The sub-sector includes production of cement, lime, container glass/glass wool, mineral wool, other production (consumption of lime), and roofing and road paving with asphalt. The activity data as well as emission data are primarily based on information from Environmental Reports (In Danish: "Grønne regnskaber") prepared by companies according to obligations under Danish law. Also, data on production and import/export from Statistics Denmark are used. The published information is supplemented with information obtained directly from companies or by use of standard emission factors. The distribution of TSP between PM<sub>10</sub> and PM<sub>2.5</sub> is based on European average data.

##### **Chemical industry**

The sub-sector includes production of nitric and sulphuric acid (ceased in 1997 and 2004, respectively), catalysts, fertilisers and pesticides. The activity data as well as emission data are based on information from the companies as accounted for and published in the Environmental Reports combined with information obtained by contact to the companies. The distribution of TSP between PM<sub>10</sub> and PM<sub>2.5</sub> is based on European average data.

##### **Metal industry**

The sub-sector includes electro steelwork, production of steel sheets and bars (electro steelwork until 2005 and thereafter, only rolling mills), cast iron, aluminium (ceased in 2008), lead and lead products and various other metal products. The activity data as well as emission data for the steelworks are based on information from the companies as accounted for and published in the Environmental Reports, combined with information obtained through contact with the companies. The activity data for the other processes are based on information from Statistics Denmark combined with Danish average emission factors and standard emission factors. The particle size distribution of TSP (PM<sub>10</sub> and PM<sub>2.5</sub>) is based on European average data.

##### **Other production**

The sub-sector includes breweries, production of spirits and other activities within the food sector e.g. sugar production, meat curing and production of margarine and solid cooking fats. The activity data are obtained from Statistics Denmark and the emission factors are obtained from the EMEP/EEA Guidebook combined with emission factors (EF) derived from specific emission measurements at the companies.

##### **Solvent and other product use**

The approach for calculating the emissions of Non-Methane Volatile Organic Carbon (NMVOC) from industrial and household use in Denmark focuses on single chemicals rather than activities. This leads to a clearer picture of the influence from each specific chemical, which enables a more detailed differentiation on products and the influence of product use on emissions. The procedure is to quantify the use of the chemicals and estimate the fraction of the chemicals that is emitted as a consequence of use.

The detailed approach in EMEP/EEA Guidebook (2013) is used. Here all relevant consumption data on all relevant solvents must be inventoried or at least those together representing more than 90 % of the total NMVOC emission. Simple mass balances for calculating the use and emissions of chemicals are set up 1) use = production + import – export, 2) emission = use emission factor. Production, import and export figures are extracted from Statistics Denmark, from which a list of more than 400 single chemicals, a few groups and products is generated. For each of these, a “use” amount in tonnes per year (from 1990 to 2013) is calculated. For some chemicals and/or products, e.g. propellants used in aerosol cans and ethanol used in wind-screen washing agents, use amounts are obtained from the industry as the information from Statistics Denmark does not comply with required specificity. It is found that approximately 40 different NMVOCs comprise over 95 % of the total use and it is these 40 chemicals that are investigated further. The “use” amounts are distributed across industrial activities according to the Nordic SPIN (Substances in Preparations in Nordic Countries) database, where information on industrial use categories is available in a NACE coding system. The chemicals are also related to specific products according to the Use Category (UCN) system. Emission factors are obtained from regulators, literature or the industry.

The same method is used for calculating emissions from the use of fireworks, tobacco, candles and charcoal for barbecues (BBQ). These activities lead to emissions of SO<sub>2</sub>, NO<sub>x</sub>, CO, NH<sub>3</sub>, particles, As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, dioxins/furans and PAHs.

Please refer to Chapter 4 and Annex 3C for further information on industrial processes and product use.

#### **1.4.5 Specific methodologies regarding agriculture**

The emission from agricultural activities covers ammonia, NO<sub>x</sub>, NMVOC and particles from animal husbandry/manure management and agricultural soils. Furthermore, the inventory includes emissions from field burning of straw which covers NH<sub>3</sub>, PM, NO<sub>x</sub>, CO, NMVOC, SO<sub>2</sub>, heavy metals, dioxin and PAH.

Emissions from agricultural activities are estimated according to the methodology described in the EMEP/EEA air pollutant emission inventory guidebook (EEA, 2013). Activity data and national data regarding emission factors are collected, evaluated and discussed in cooperation with Statistics Denmark, DCA-Danish Centre for Food and Agriculture (Aarhus University), the Danish Agricultural Advisory Service, Danish Environmental Protection Agency and the Danish AgriFish Agency. It means that data are evaluated continuously according to the latest knowledge and information.

The Danish agricultural emissions are calculated and managed in a comprehensive model complex called IDA (Integrated Database model for Agricultural emissions), which is used to calculate both air pollutants compounds and greenhouse gas related emissions. The livestock production has a great influence on the Danish agricultural emissions. IDA works with approximately 40 different livestock categories, dependent on livestock category, weight class and age. Each of these subcategories is subdivided according to housing type and manure type, which results in about 200 different combinations of subcategories and housing type and the emissions are calculated



from each of these combinations and aggregated to relevant main categories in the reporting format.

Most of the emissions from agricultural activities are directly related to livestock production. The remaining part comes from the use of synthetic fertiliser, growing crops, NH<sub>3</sub> treated straw, field burning of agricultural residues and sewage sludge applied to fields as fertiliser. The number of animals can be considered as the most important activity data in estimation of the agricultural emissions.

The number of animals is mainly based on data from Statistics Denmark. For data covering pigs, bulls and poultry, the number is based on slaughter data also collected from the Agricultural Statistics. The production of sheep, goats and horses typically take place on small farms below 5 hectare, which is not included in the annual statistics and the production of these categories as well as for deer and ostriches are therefore based on the Central House-animal farm Register (CHR) managed by the Ministry of Food, Agriculture and Fisheries.

Data concerning nitrogen excretion, distribution of housing types until 2004 and handling of manure is based on data and information from DCA-Danish Centre for Food and Agriculture at Aarhus University and the Danish Agricultural Advisory Service. From 2005 annual statistics covering housing types are available from the Danish AgriFish Agency.

Data related to use of synthetic fertiliser, both the amount of fertiliser and the nitrogen content is based on statistics published by the Danish AgriFish Agency.

Please refer to Chapter 5 and Appendix 3D for further information on emission inventories for agriculture.

#### **1.4.6 Specific methodologies regarding waste**

The waste sector consists of the four main NFR categories 5A Solid waste disposal, 5B Biological treatment of solid waste, 5C Waste incineration, 5D Wastewater treatment and discharge and 5E Other waste.

Emissions from solid waste disposal and wastewater treatment and discharge are currently not estimated.

Composting includes four types of biological waste; garden and park waste, organic waste from households and other sources, sludge and home composting of garden and vegetable food waste. Individual emission factors are found for each waste category.

Waste incineration covers the cremation of human bodies and animal carcasses. Both are calculated as an activity multiplied by an emission factor.

The Other waste category includes accidental building- and vehicle fires

Emissions from building fires are calculated by multiplying the number of building fires with selected emission factors. Six types of buildings are separated with different emission factors; detached houses, undetached houses,

apartment buildings, industrial buildings, additional buildings and containers.

Activity data for building fires are classified in four categories; full scale, large, medium and small. The emission factors comply for full scale building fires and the activity data are therefore recalculated as a full scale equivalent where it is assumed that a large, medium and small fire leads to 75 %, 30 % and 5 % of a full scale fire respectively.

Emissions from vehicle fires are calculated by multiplying the total burnt vehicle mass with selected emission factors. 14 different vehicle types are included in the total mass of burned vehicle. Emission factors are not available for different vehicle types, why it is assumed that all the different vehicle types lead to similar emissions. As with accidental building fires, four different sizes are known in relation to damage; full scale (100 % burnout), large (75 %), medium (30 %) and small (5 %).

Please refer to Chapter 6 and Annex 3E for further information on emission inventories for agriculture.

## **1.5 Key categories**

The determination of key categories has not been made due to insufficient resources being available at the moment.

## **1.6 Information on the Quality Control and Quality Assurance plan including verification and treatment of confidential issues where relevant**

In the Danish National Inventory Report to UNFCCC (Nielsen et al., 2014) as well as in the QA/QC manual for the Danish Greenhouse gas inventory (Nielsen et al., 2012), the plan for Quality Control (QC) and Quality Assurance (QA) for greenhouse gas emission inventories prepared by the Danish National Environmental Research Institute is outlined. The plan is in accordance with the guidelines provided by the UNFCCC (IPCC, 2006). The ISO 9000 standards are also used as important input for the plan. The plan also, to a limited extent, includes the gases reported to the UNECE-LRTAP Convention. Due to a lack of resources it has not been possible to extend the QA/QC system for the greenhouse gas inventory to also cover the air pollutants.

## **1.7 General uncertainty evaluation, including data on the overall uncertainty for the inventory totals**

The uncertainty estimates are based on the simple Tier 1 approach in the EMEP/CorinAir *Good Practice Guidance for LRTAP Emission Inventories* (Pulles & Aardenne, 2004).

The uncertainty estimates are based on emission data for the base year and year 2013, and on uncertainties for activity rates and emission factors for each of the main SNAP sectors. For particulate matter and black carbon (BC), the year 2000 is considered as the base year, but for all other pollutants 1990 is used as the base year.

Uncertainty estimates include uncertainty of the total emission as well as uncertainty of the trend. The estimated uncertainties are shown in Table 1.2. The uncertainty estimates include all sectors.

Table 1.2 Danish uncertainty estimates, 2013.

Pollutant	Uncertainty Total emis- sion [%]	Trend 1990 <sup>1)</sup> -2013 [%]	Uncertainty Trend [%-age points]
SO <sub>2</sub>	14	-92	1.0
NO <sub>x</sub>	47	-59	8
NMVOC	97	-44	24
CO	47	-55	15
NH <sub>3</sub>	23	-41	11
TSP	191	-11	12
PM <sub>10</sub>	110	-11	22
PM <sub>2.5</sub>	123	-11	33
BC	48	-43	5
Arsenic	130	-73	21
Cadmium	427	-47	192
Chromium	237	-74	54
Copper	936	30	88
Mercury	91	-89	10
Nickel	403	-80	48
Lead	491	-91	30
Selenium	266	-69	49
Zinc	472	-19	208
PCDD/F	315	-68	101
Benzo(b)fluoranthene	671	47	210
Benzo(k)fluoranthene	634	50	141
Benzo(a)pyrene	723	42	178
Indeno(1,2,3-c,d)pyrene	643	15	244
HCB	527	-90	51
PCBs	703	-64	125

<sup>1)</sup>The base year for PM and BC is 2000.

## 1.8 General assessment of the completeness

Annex 4 provides a full and comprehensive explanation on the use of notation keys in the Danish inventory. The Danish emission inventory due 15 February 2015 includes all sources identified by the EMEP/EEA guidebook except the following.

### 1.8.1 Industrial processes

Categories reported as not estimated:

- Emissions from quarrying and mining of minerals other than coal
- Emissions from construction and demolition
- Emissions from storage, handling and transport of mineral products
- Emissions from storage, handling and transport of chemical products
- Emissions from storage, handling and transport of metal products
- Emissions from pulp and paper production
- Emissions from wood processing
- Emissions from production of POPs
- Emissions from consumption of POPs and heavy metals

### 1.8.2 Agriculture

Categories reported as not estimated:

- Emissions from off-farm storage, handling and transport of agricultural products

### 1.8.3 Waste

Categories reported as not estimated:

- Emissions from solid waste disposal on land
- Emissions from wastewater handling
- Emissions from small scale waste burning

### 1.8.4 Categories reported as “included elsewhere”

The following table lists the categories reported as IE (included elsewhere) and provides information on where the associated emissions are reported, more detailed information is provided in Annex 4.

Table 1.3. List of categories reported as included elsewhere.

Category reported as IE	Category where emissions are included
2 A 1 Cement production	1 A 2 f Stationary combustion in manufacturing industries and construction: Non-metallic minerals
2 A 2 Lime production	1 A 2 f Stationary combustion in manufacturing industries and construction: Non-metallic minerals
2 A 3 Glass production	1 A 2 f Stationary combustion in manufacturing industries and construction: Non-metallic minerals

### 1.8.5 General description on the use of notation keys

The NFR as reported by Denmark makes use of five notation keys: NO (Not Occurring), NA (Not Applicable), NE (Not Estimated), IE (Included Elsewhere) and NR (Not Reported).

NO is used in instances where the activity does not occur in Denmark, e.g. adipic acid production, buffaloes, etc.

NA is used in instances where the activity occurs in Denmark but the emission of a certain pollutant is not believed to be relevant, e.g. heavy metals from dairy cattle.

NE is used in instances where the activity occurs in Denmark and emissions of a certain pollutant are thought to occur but the emission has not been estimated; see Chapter 1.8.3 and Annex 4.

IE is used where emissions of a certain pollutant or the whole source category are reported under another source category; see Chapter 1.8.4 and Annex 4.

NR is used for pollutants prior to the base year, e.g. PM emissions prior to the year 2000.

## 1.9 References

EEA, 2009: EMEP/EEA air pollutant emission inventory guidebook — 2009 prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections. Available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (30-12-2014)

EEA, 2013: EMEP/EEA air pollutant emission inventory guidebook — 2013 prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections. Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (30-12-2014)

Illerup, J.B., Lyck, E., Winther, M. & Rasmussen, E. 2000: Denmark's National Inventory Report – Submitted under the United Nations Framework Convention on Climate Change. Samfund og Miljø – Emission Inventories. Research Notes from National Environmental Research Institute, Denmark no. 127, 326 pp. Available at: [http://www.dmu.dk/1\\_viden/2\\_Publikationer/3\\_arbrapporter/rapporter/ar127.pdf](http://www.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/ar127.pdf)

IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. & Tanabe K. (eds). Published: IGES, Japan. Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html> (30-12-2014)

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M. Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K., Hoffmann, L. & Bruun, H.G. 2012. Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 4. Available at: <http://www.dmu.dk/Pub/SR47.pdf>

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L. & Hansen, M.G. 2014. Denmark's National Inventory Report 2014. Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. Aarhus University, DCE – Danish Centre for Environment and Energy, 1214pp. Scientific Report from DCE – Danish Centre for Environment and Energy. <http://dce2.au.dk/pub/SR101.pdf>

Pulles, T., Mareckova, K., Svetlik, J., Linek, M. & Skakala, J. 1999: CollectER - Installation and User Guide, EEA Technical Report No 31. Available at: <http://reports.eea.eu.int/binarytech31pdf/en> (26-01-2014).

Pulles, T. & Aardenne, J.v., 2004: Good Practice Guidance for LRTAP Emission Inventories, 24. June 2004.

## 2 Trends in Emissions

### 2.1 Acidifying gases

Acid deposition of sulphur and nitrogen compounds mainly derives from emissions of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub>. The effects of acidification may appear in a number of ways, including defoliation and reduced vitality of trees, and declining fish stocks in acid-sensitive lakes and rivers.

SO<sub>2</sub> and NO<sub>x</sub> can be oxidised into sulphate (SO<sub>4</sub><sup>-</sup>) and nitrate (NO<sub>3</sub><sup>-</sup>) - either in the atmosphere or after deposition - resulting in the formation of two and one H<sup>+</sup>, respectively. NH<sub>3</sub> may react with H<sup>+</sup> to form ammonium (NH<sub>4</sub><sup>+</sup>) and, by nitrification in soil, NH<sub>4</sub><sup>+</sup> is oxidised to NO<sub>3</sub><sup>-</sup> and H<sup>+</sup> ions are formed.

Weighting the individual substances according to their acidification effect, total emissions in terms of acid equivalents can be calculated as:

$$A = \frac{m_{SO_2}}{M_{SO_2}} \cdot 2 + \frac{m_{NO_x}}{M_{NO_x}} + \frac{m_{NH_3}}{M_{NH_3}} = \frac{m_{SO_2}}{64} \cdot 2 + \frac{m_{NO_x}}{46} + \frac{m_{NH_3}}{17}$$

where  $A$  is the acidification index in Mmole

$m_i$  is the emission of pollutant  $i$  in tonnes

$M_i$  is the mole weight [tonne/Mmole] of pollutant  $i$

The actual effect of the acidifying substances depends on a combination of two factors: the amount of acid deposition and the natural capacity of the terrestrial or aquatic ecosystem to counteract the acidification. In areas where the soil minerals easily weather or have a high lime content, acid deposition will be relatively easily neutralised.

Figure 2.1 shows the emission of Danish acidifying gases in terms of acid equivalents. In 1990, the relative contribution in acid equivalents was almost equal for the three gases. In 2013, the most important acidification factor in Denmark is ammonia nitrogen and the relative contributions for SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 6 %, 36 % and 58 %, respectively. However, with regard to long-range transport of air pollution, SO<sub>2</sub> and NO<sub>x</sub> are still the most important pollutants.




Figure 2.1 Emissions of NH<sub>3</sub>, NO<sub>x</sub> and SO<sub>2</sub> over time in acid equivalents.

## 2.2 Description and interpretation of emission trends by gas

### 2.2.1 Sulphur dioxide (SO<sub>2</sub>)

The main part of the sulphur dioxide (SO<sub>2</sub>) emission originates from combustion of fossil fuels, i.e. mainly coal and oil, in public power and district heating plants. From 1990 to 2013, the total emission decreased by 92 %. The large reduction is mainly due to installation of desulphurisation plant and use of fuels with lower content of sulphur in public power and district heating plants. Despite the large reduction of the SO<sub>2</sub> emissions, these plants make up 27 % of the total emission. Also emissions from industrial combustion plants, non-industrial combustion plants and other mobile sources are important. National sea traffic (navigation and fishing) contributes with about 11 % of the total SO<sub>2</sub> emission in 2013. This is due to the use of residual oil with high sulphur content.




Figure 2.2 SO<sub>2</sub> emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.2.2 Nitrogen oxide (NO<sub>x</sub>)

The largest sources of emissions of nitrogen oxides (NO<sub>x</sub>) are road transport followed by other mobile sources and combustion in energy industries (mainly public power and district heating plants). The transport sector is the sector contributing the most to the emission of NO<sub>x</sub> and, in 2013, 42 % of the Danish emissions of NO<sub>x</sub> stems from road transport, national navigation, railways and civil aviation. Also emissions from national fishing and off-

road vehicles contribute significantly to the NO<sub>x</sub> emission. For non-industrial combustion plants, the main sources are combustion of gas oil, natural gas and wood in residential plants. The emissions from energy industries have decreased by 78 % from 1990 to 2013. In the same period, the total emission decreased by 59 %. The reduction is due to the increasing use of catalyst cars and installation of low-NO<sub>x</sub> burners and denitrifying units in power plants and district heating plants.




Figure 2.3 NO<sub>x</sub> emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.2.3 Ammonia (NH<sub>3</sub>)

Almost all atmospheric emissions of ammonia (NH<sub>3</sub>) result from agricultural activities. Only a minor fraction originates from stationary combustion (2 %), road transport (2 %), industrial processes and product use (1 %) and waste (1 %). The share for road transport was increasing during the 1990's and early 2000's due to increasing use of catalyst cars. In more recent years the share has been decreasing due to more advanced catalysts being implemented. The major part of the emission from agriculture stems from livestock manure (51 %) and the largest losses of ammonia occur during the handling of the manure in animal housing systems. The second largest agricultural source is agricultural soils contributing 48 % in 2013, this is mainly emissions from application of mineral fertiliser, application of animal manure and emissions from growing crops. The total ammonia emission decreased by 42 % from 1985 to 2013. Due to the action plans for the aquatic environment and the Ammonia Action Plan, a series of measures to prevent loss of nitrogen in agricultural production has been initiated. The measures have included demands for improved utilisation of nitrogen in livestock manure, a ban against field application of livestock manure in winter, prohibition of broadspraying of manure, requirements for establishment of catch crops, regulation of the number of livestock pr hectare and a ceiling for the supply of nitrogen to crops. As a result, despite an increase in the production of pigs and poultry, the ammonia emission has been reduced considerably.






Figure 2.4 NH<sub>3</sub> emissions. Distribution on the main sectors (2013) and time series for 1985 to 2013.

## 2.3 Other air pollutants

### 2.3.1 Non-Methane Volatile Organic Compounds (NMVOC)

The emissions of Non-Methane Volatile Organic Compounds (NMVOC) originate from many different sources and can be divided into two main groups: incomplete combustion and evaporation. Road vehicles and other mobile sources such as national navigation vessels and off-road machinery contribute approximately 50 % of the NMVOC emissions from incomplete combustion processes. NMVOC from road transportation vehicles have been decreasing since 1990, due to the introduction of catalyst cars. The evaporative emissions mainly originate from the agricultural sector, use of solvents, and the extraction, handling and storage of oil and natural gas. The total anthropogenic emissions have decreased by 44 % from 1990 to 2013, largely due to the increased use of catalyst cars and reduced emissions from use of solvents.




Figure 2.5 NMVOC emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.3.2 Carbonmonoxid (CO)

Non-industrial combustion plants are the main source to the total CO emission. For the non-industrial sector, emissions from commercial/institutional sources have increased and emissions from agriculture/forestry/fishing sources have decreased from 1990 to 2013, while emissions from the residential sector have been fluctuating, but around the same level in 1990 and 2013. Transport is the second largest contributor to the total CO emission in 2013, showing a decrease of 81 % from 1990 to 2013. The major transport source is passenger cars, which make up 58 % in 1990, but has decreased to 20 % in 2013. The main driver is the increase of catalyst cars. In 1990 a law forbidding the burning of agricultural crop residues on fields was implemented,

which caused a significant reduction in CO emission. The total CO emission decreased further by 55 % from 1990 to 2013, largely because of decreasing emissions from road transportation.




Figure 2.6 CO emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.3.3 Particulate matter (PM)

The particulate matter (PM) emission inventory is reported for the years 2000 onwards. The inventory includes the total emission of particles TSP (Total Suspended Particles), emission of particles smaller than 10  $\mu\text{m}$  ( $\text{PM}_{10}$ ) and emission of particles smaller than 2.5  $\mu\text{m}$  ( $\text{PM}_{2.5}$ ).

The largest  $\text{PM}_{2.5}$  emission source is residential plants (63 %), road transport (9 %) and other mobile sources (8 %). Emissions from residential plants have increased by 32 % from 2000 to 2007, followed by a decrease of 32 % from 2007 to 2013. For road transport, the most important sources are off-road vehicles and machinery in the industrial sector and in the agricultural/forestry sector (35 % and 33 %, respectively). For the road transport sector, exhaust emissions account for the major part (57 %) of the emissions. The  $\text{PM}_{2.5}$  emission decreased by 11 % from 2000 to 2013 as the increasing wood consumption in the residential sector has been counterbalanced by decreasing emissions for the remaining sectors, the most important being the Transport sector.

The largest TSP emission sources are agriculture and non-industrial combustion (74 % and 18 % of total TSP emission in 2013, respectively). Residential plants is the largest source in the non-industrial combustion sector, making up 16 % of the national total TSP emission in 2013. The TSP emissions from transport are also important and include both exhaust emissions and the non-exhaust emissions from brake and tyre wear and road abrasion. The non-exhaust emissions account for 68 % of the TSP emission from road transport in 2013.




Figure 2.7 PM emissions per sector for 2013.




Figure 2.8 PM<sub>2.5</sub> emissions. Distribution according to the main sectors (2013) and time series for 2000 to 2013.

### 2.3.4 Black carbon (BC)

The black carbon (BC) emission inventory is reported for the years 2000 onwards. The main sources are residential plants and road transport contributing 38 % and 19 % in 2013, respectively. From 2000 to 2013 the total BC emission decreases by 24 %. BC emissions from non-industrial plants have increased by 43 % from 2000 to 2007, followed by a decrease of 30 % from 2007 to 2013. The trend for non-industrial combustion is mainly controlled by the trend for the wood consumption in the residential sector. BC emissions from the transport sector decreased by 49 % from 2000 to 2013, mainly due to implementing of new EURO norms and improved technology. An important factor is the use of particle filters for heavy duty vehicles and personal cars, which reduce the BC emission effectively. BC emissions from fugitive emissions from fuels, which is mainly due to storage of coal, decreased by 23 % from 2000 to 2013, in accordance with the decrease of the coal consumption in electricity and heat production.




Figure 2.9 BC emissions. Distribution according to the main sectors (2013) and time series for 2000 to 2013.

### 2.3.5 Heavy metals

In general, the most important sources of heavy metal emissions are combustion of fossil fuels and waste. The heavy metal emissions have decreased substantially in recent years, except for Cu. The reductions span from 19 % to 91 % for Zn and Pb, respectively. The reason for the reduced emissions is mainly increased use of gas cleaning devices at power and district heating plants (including waste incineration plants). The large reduction in the Pb emission is due to a gradual shift towards unleaded gasoline, the latter being essential for catalyst cars. The major source of Cu is automobile tyre and break wear (93 % in 2013) and the 30 % increase from 1990 to 2013 owe to increasing mileage.

Table 2.1 Emissions of heavy metals.

Heavy metals, kilogramme	As	Cd	Cr	Cu	Hg	Ni	Pb	Se	Zn
1990	1.344	1.095	5.811	32.899	3.172	21.062	128.388	4.844	69.122
2013	369	581	1.486	42.808	355	4.220	11.599	1.488	55.797
Reduction, %	73	47	74	-30	89	80	91	69	19

According to the UNECE Heavy Metal Protocol, the priority metals are Pb, Cd and Hg and the objective is to reduce emissions of these heavy metals.

#### Cadmium (Cd)

The main sources of emissions of cadmium (Cd) to air are mainly combustion of wood, wood waste and municipal waste. Non-industrial combustion contributes 73 % in 2013, of which 97 % comes from residential plants. Emissions from residential plants have increased by 225 % from 1990 to 2013 due to increasing wood consumption. Emissions from energy industries, manufacturing industries and construction, and industrial processes have decreased by 88 % from 1990 to 2013. The decreasing emission from energy industries are related to the decreasing combustion of coal. In the transport sector emissions from passenger cars is the main source contributing with 58 % of the sectoral emission in 2013.




Figure 2.10 Cd emissions. Time series for 1990 to 2013 and distribution by main sector for 2013.

### Mercury (Hg)

The largest sources of mercury (Hg) emissions to air are waste incineration and coal combustion in energy industries. Due to improved flue gas cleaning and decreasing coal combustion the emissions from Energy industries decreased by 76 % from 1990-2000. The trend has continued in the following years and the corresponding decrease from 1990-2013 is 92 %. Non-industrial combustion is dominated by wood combustion in residential plants while the main contributions to emissions from manufacturing industries and construction are food processing, beverages and tobacco, and non-metallic minerals. The variations in emissions from industrial processes owe to shut down in 2002 followed by re-opening and a second shut down in 2005 of the only Danish electro-steelwork.




Figure 2.11 Hg emissions. Time series for 1990 to 2013 and distribution by main sector for 2013.

### Lead (Pb)

The main lead (Pb) emission sources are transport, waste, non-industrial combustion and industrial processes. In earlier years combustion of leaded gasoline was the major contributor to Pb emissions to air but the shift toward use of unleaded gasoline for transport have decreased the Pb emission from transport by 94 % from 1990-2013. The trend in the Pb emission from non-industrial combustion from 1990 to 2013 is almost constant. In the non-industrial combustion sector the dominant source is wood combustion in residential plants, which has been increasing from 1990 to 2013, but counterbalanced by decreasing emissions from stationary combustion in commercial/institutional and in agriculture/forestry/fishing. The decreasing emission from Energy industries (96 % from 1990 to 2013) is caused by the decreasing coal combustion and more efficient particle abatement.




Figure 2.12 Pb emissions. Time series for 1990 to 2013 and distribution by main sector for 2013.

### 2.3.6 Polycyclic aromatic hydrocarbons (PAHs)

The present emission inventory for polycyclic aromatic hydrocarbons (PAH) includes four PAHs: benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene and indeno(1,2,3-cd)pyrene. The major part of the total PAH emission is Benzo(b)fluoranthene and benzo(a)pyrene, which contribute by 34 % and 32 %, respectively in 2013.




Figure 2.13 PAH emissions. Distribution according to reported PAHs in 2013.

The most important source of PAHs emissions is combustion of wood in the residential sector making up 68 % of the total emission in 2013. The increasing emission trend is due to increasing combustion of wood in the residential sector. The PAH emission from combustion in residential plants has increased by 47 % from 1990 to 2013.




Figure 2.14 PAH emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.3.7 Dioxins and furans

The major part of the dioxin emission owes to wood combustion in the residential sector, mainly in wood stoves and ovens without flue gas cleaning. Wood combustion in residential plants accounts for 57 % of the national dioxin emission in 2013. The contribution to the total dioxin emission from the waste sector (26 % in 2013) mainly owes to accidental fires, especially building fires. The emissions of dioxins from energy industries are dominated by emissions from combustion of biomass as wood, wood waste and to a less extend agricultural waste.




Figure 2.15 Emissions of dioxins and furans. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.3.8 Hexachlorobenzene (HCB)

Stationary combustion accounts for 51 % of the estimated national hexachlorobenzene (HCB) emission in 2013. This owes mainly to combustion of municipal solid waste in heating and power plants. Transport is an important source, too, making up 25 % of the total emission in 2013. Emissions from transport have increased by 56 % since 1990 due to increasing diesel consumption. The HCB emission from stationary *plants* has decreased 69 % since 1990 mainly due to improved flue gas cleaning in MSW incineration plants. The emission from agriculture was very high in the early 1990'ties due to the use of pesticides containing impurities of HCB. The HCB emission from agriculture decreased by 94 % from 1990 to 1994 and by 99 % from 1990 to 2013, causing the share of HCB emission from agriculture to drop from 67 % in 1990 to 5 % in 2013. The emission from industrial processes has decreased due to the closure of steel production and secondary aluminium production in Denmark.




Figure 2.16 HCB emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.

### 2.3.9 Polychlorinated biphenyls (PCBs)

Transport accounts for 60 % of the estimated national polychlorinated biphenyls (PCBs) emission in 2013. This owes mainly to combustion of diesel in road transport. The emission from transport has decreased by 73 % since 1990 due to the phase out of leaded gasoline, which has a high PCBs emission factor. This has led to diesel fuel use being the most important source of PCBs emissions from transport in later years. The emission from manufacturing industries and non-industrial combustion is dominated by diesel fuel used in non-road machinery.




Figure 2.17 PCBs emissions. Distribution according to the main sectors (2013) and time series for 1990 to 2013.



### 3 Energy (NFR sector 1)

#### 3.1 Overview of the sector

The energy sector is reported in three main chapters:

3.2 Stationary combustion plants (NFR sector 1A1, 1A2 and 1A4)

3.3 Mobile combustion (NFR sector 1A2, 1A3, 1A4 and 1A5)

3.4 Fugitive emissions (NFR sector 1B)

Summary tables for the emissions from the energy sector are shown below.

Table 3.1.1 (3.1?) SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, PM and BC emissions from the energy sector, 2013.

	NO <sub>x</sub>	NMVOC	SO <sub>x</sub>	NH <sub>3</sub>	PM <sub>2.5</sub>	PM <sub>10</sub>	TSP	BC	CO
	kt NO <sub>2</sub>	kt	kt SO <sub>2</sub>	kt	kt	kt	kt	kt	kt
1A1 Energy Industries	20.9	1.4	3.6	0.02	0.68	0.84	1.04	0.40	10.4
1A2 Manufacturing industries and Construction	12.5	1.3	2.8	0.00	0.80	0.88	0.96	0.51	10.6
1A3 Transport	52.5	9.9	1.4	1.27	2.22	2.92	3.82	1.05	88.9
1A4 Other Sectors	21.8	21.6	2.5	1.29	14.80	15.17	15.93	2.47	217.0
1A5 Other	1.5	0.4	0.1	0.00	0.10	0.10	0.10	0.04	3.9
1B1 Fugitive Emissions from fuels, Solid fuels	-	-	-	-	0.03	0.30	0.74	0.50	-
1B2 Fugitive Emissions from fuels, Oil and Natural gas	0.1	9.2	0.7	-	0.00	0.00	0.00	0.00	0.2
<b>Energy, Total</b>	<b>109.3</b>	<b>43.8</b>	<b>11.1</b>	<b>2.58</b>	<b>18.63</b>	<b>20.20</b>	<b>22.61</b>	<b>4.96</b>	<b>330.9</b>

Table 3.1.2 HM emissions from the energy sector, 2013.

	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn
	t	t	t	t	t	t	t	t	t
1A1 Energy Industries	0.46	0.05	0.20	0.18	0.24	0.24	0.64	0.84	0.64
1A2 Manufacturing industries and Construction	0.45	0.03	0.07	0.07	0.09	0.11	1.29	0.08	1.50
1A3 Transport	6.40	0.05	0.03	0.04	0.19	40.08	1.80	0.06	26.05
1A4 Other Sectors	1.28	0.42	0.05	0.03	0.78	0.28	0.20	0.09	17.77
1A5 Other	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.09
1B1 Fugitive Emissions from fuels, Solid fuels	-	-	-	-	-	-	-	-	-
1B2 Fugitive Emissions from fuels, Oil and Natural gas	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
<b>Energy, Total</b>	<b>8.63</b>	<b>0.55</b>	<b>0.34</b>	<b>0.33</b>	<b>1.31</b>	<b>40.72</b>	<b>3.94</b>	<b>1.07</b>	<b>46.05</b>

Table 3.1.3 PAH, dioxin and HCB emissions from the energy sector, 2013.

	PCDD/ PCDF	Ben- zo(a)- pyrene	Ben- zo(b)- fluoran- thene	Ben- zo(k)- fluoran- thene	Indeno- (1,2,3- cd)- pyrene	HCB	PCB
	g I-Teq	t	t	t	t	kg	kg
1A1 Energy Industries	1.16	0.01	0.04	0.02	0.01	1.25	0.28
1A2 Manufacturing industries and Construction	0.06	0.02	0.08	0.02	0.01	0.15	6.82
1A3 Transport	0.18	0.06	0.09	0.08	0.07	0.67	24.09
1A4 Other Sectors	14.25	1.98	2.05	0.72	1.27	0.35	8.02
1A5 Other	0.00	0.00	0.00	0.00	0.00	0.01	0.71
1B1 Fugitive Emissions from fuels, Solid fuels	-	-	-	-	-	-	-
1B2 Fugitive Emissions from fuels, Oil and Natural gas	0.00	0.00	0.00	0.00	0.00	-	-
<b>Energy, Total</b>	<b>15.67</b>	<b>2.07</b>	<b>2.25</b>	<b>0.84</b>	<b>1.35</b>	<b>2.44</b>	<b>39.92</b>

## 3.2 Stationary combustion (NFR sector 1A1, 1A2 and 1A4)

This chapter includes stationary combustion plants in the NFR sectors 1A1, 1A2 and 1A4.

### 3.2.1 Source category description

#### Source category definition

In the Danish emission database, all activity rates and emissions are defined in SNAP sector categories (Selected Nomenclature for Air Pollution) according to the CORINAIR system<sup>1</sup>. The emission inventories are prepared from a complete emission database based on the SNAP sectors. Aggregation to the NFR sector codes is based on a correspondence list between SNAP and NFR enclosed in Annex 3A-1. Stationary combustion is defined as combustion activities in the SNAP sectors 01-03, not including SNAP 0303.

Stationary combustion plants are included in the emission source subcategories:

- 1A1 Energy, Fuel consumption, Energy Industries
  - 1A1a Public electricity and heat production
  - 1A1b Petroleum refining
  - 1A1c Oil and gas extraction
- 1A2 Energy, Fuel consumption, Manufacturing Industries and Construction
  - 1A2a Iron and steel
  - 1A2b Non-ferrous metals
  - 1A2c Chemicals
  - 1A2d Pulp, Paper and Print
  - 1A2e Food processing, beverages and tobacco
  - 1A2f Non-metallic minerals
  - 1A2 g viii Other manufacturing industry
- 1A4 Energy, Fuel consumption, Other Sectors
  - 1A4a i Commercial/Institutional plants.
  - 1A4b I Residential plants.
  - 1A1c I Agriculture/Forestry.

The emission and fuel consumption data included in tables and figures in Chapter 3.2 only include emissions originating from stationary combustion plants of a given NFR sector.

#### Emission share from stationary combustion compared to national total

Table 3.2.1 gives an overview of the emission share from stationary combustion compared to national total. Main emission sources are discussed in chapter 3.2.3. Key category analysis has not been performed.

<sup>1</sup> And some additional SNAP added for industrial combustion.

Table 3.2.1 Emission share from stationary combustion compared to national total.


Pollutant	Emission share, %
SO <sub>2</sub>	63
NO <sub>x</sub>	25
NM VOC	14
CO	34
NH <sub>3</sub>	2
TSP	19
PM <sub>10</sub>	48
PM <sub>2.5</sub>	70
BC	51
As	76
Cd	85
Cr	73
Cu	1
Hg	85
Ni	50
Pb	18
Se	65
Zn	33
HCB	58
PCDD/F	71
Benzo(a)pyrene	88
Benzo(b)fluoranthene	87
Benzo(k)fluoranthene	79
Indeno(123cd)pyrene	85
PCB	1

### 3.2.2 Fuel consumption data

In 2013, the total fuel consumption for stationary combustion plants was 461 PJ of which 331 PJ was fossil fuels and 130 PJ was biomass.

Fuel consumption distributed according to the stationary combustion sub-categories is shown in Figure 3.2.1 and Figure 3.2.2. The majority - 60 % - of all fuels is combusted in the source category, *Public electricity and heat production*. Other source categories with high fuel consumption are *Residential plants* and *Industry*.

### Fuel consumption including biomass



### Fuel consumption, fossil fuels




Figure 3.2.1 Fuel consumption of stationary combustion source categories, 2013. Based on DEA (2014a).

Coal, natural gas and wood are the most utilised fuels for stationary combustion plants. Coal is mainly used in power plants and natural gas is used in power plants and decentralised combined heating and power (CHP) plants, as well as in industry, residential plants and off-shore gas turbines (see Figure 3.2.2). Wood is mainly applied for public electricity and heat production and in residential plants.

Detailed fuel consumption rates are shown in Annex 3A-2.





Figure 3.2.2 Fuel consumption of stationary combustion 2013, disaggregated to fuel type. Based on DEA (2014a).

Fuel consumption time series for stationary combustion plants are presented in Figure 3.2.3<sup>2</sup>. The fuel consumption for stationary combustion was 8 % lower in 2013 than in 1990, while the fossil fuel consumption was 28 % lower and the biomass fuel consumption 3.2 times the level in 1990.

The consumption of natural gas, waste and biomass has increased since 1990.

<sup>2</sup> Time series 1980 whereas the consumption of coal has decreased. onwards are included in Annex 3A-10.




Figure 3.2.3 Fuel consumption time series, stationary combustion. Based on DEA (2014a).

The fluctuations in the time series for fuel consumption are mainly a result of electricity import/export, but also of outdoor temperature variations from year to year. This, in turn, leads to fluctuations in emission levels. The fluctuations in electricity trade, fuel consumption and  $\text{NO}_x$  emission are illustrated and compared in Figure 3.2.4. In 1990, the Danish electricity import was large causing relatively low fuel consumption, whereas the fuel consumption was high in 1996 due to a large electricity export. In 2013, the net electricity import was 4 PJ, whereas there was a 19 PJ electricity import in 2012. The large electricity export that occurs some years is a result of low rainfall in Norway and Sweden causing insufficient hydropower production in both countries.

To be able to follow the national energy consumption as well as for statistical and reporting purposes, the Danish Energy Agency produces a correction of the actual fuel consumption without random variations in electricity imports/exports and in ambient temperature. This fuel consumption

trend is also illustrated in Figure 3.2.4. The corrections are included here to explain the fluctuations in the time series for fuel rate and emissions.




Figure 3.2.4 Comparison of time series fluctuations for electricity trade, fuel consumption and NO<sub>x</sub> emission. Based on DEA (2014a).

Fuel consumption time series for the subcategories to stationary combustion are shown in Figure 3.2.5 – 3.2.7.

Fuel consumption for *Energy Industries* fluctuates due to electricity trade as discussed above. The fuel consumption in 2013 was 1 % higher than in 1990. However, the fossil fuel consumption was 21 % lower. The fluctuation in electricity production is based on fossil fuel consumption in the subcategory *Public electricity and Heat Production*. The energy consumption

in *Oil and gas extraction* is mainly natural gas used in gas turbines in the off-shore industry. The biomass fuel consumption in *Energy Industries* in 2013 added up to 82 PJ, which is 5.1 times the level in 1990 and a 4 % increase since 2012.

The fuel consumption in *Industry* was 19 % lower in 2013 than in 1990 (Figure 3.2.6). The fuel consumption in industrial plants decreased considerably as a result of the financial crisis. The fuel consumption in 2013 was almost equal to the consumption in 2009. The biomass fuel consumption in *Industry* in 2013 added up to 9 PJ which is a 16 % increase since 1990.

The fuel consumption in *Other Sectors* decreased 23 % since 1990 (Figure 3.2.7) and decreased 3 % since 2012. The biomass fuel consumption in *Other sectors* in 2013 added up to 39 PJ which is 2.1 times the consumption in 1990 but a 2 % decrease since 2012. Wood consumption in residential plants in 2013 was 2.1 times the consumption in year 2000.

Time series for subcategories are shown in Chapter 3.2.4.




Figure 3.2.5 Fuel consumption time series for subcategories - 1A1 Energy Industries.






Figure 3.2.6 Fuel consumption time series for subcategories - 1A2 Industry.





Figure 3.2.7 Fuel consumption time series for subcategories - 1A4 Other Sectors.

### 3.2.3 Emissions

#### SO<sub>2</sub>

Stationary combustion is the most important emission source for SO<sub>2</sub> accounting for 63 % of the national emission. Table 3.2.2 presents the SO<sub>2</sub> emission inventory for the stationary combustion subcategories.

*Public electricity and heat production* is the largest emission source accounting for 38 % of the emission. However, the SO<sub>2</sub> emission share is lower than the fuel consumption share for this source category, which is 60 %. This is a result of effective flue gas desulphurisation equipment installed in power plants combusting coal. In the Danish inventory, the source category *Public electricity and heat production* is further disaggregated. Figure 3.2.8 shows the SO<sub>2</sub> emission from *Public electricity and heat production* on a disaggregated level. Power plants >300MW<sub>th</sub> are the main emission source, accounting for 47 % of the emission.

The SO<sub>2</sub> emission from industrial plants adds up to 33 % of the emission from stationary combustion, a remarkably high emission share compared


with fuel consumption. The main emission sources in the industrial category are combustion of coal and emissions from the cement industry, mineral wool industry and sugar production plants. Until year 2000, the SO<sub>2</sub> emission from the industrial category only accounted for a small part of the emission from stationary combustion, but as a result of reduced emissions from power plants, the share has now increased.

The time series for SO<sub>2</sub> emission from stationary combustion is shown in Figure 3.2.9. The SO<sub>2</sub> emission from stationary combustion plants has decreased by 94 % since 1990. The large emission decrease is mainly a result of the reduced emission from *Public electricity and heat production*, made possible due to installation of desulphurisation plants and due to the use of fuels with lower sulphur content. Despite the considerable reduction in emission from public electricity and heat production plants, these still account for 38 % of the emission from stationary combustion, as mentioned above. The emission from other source categories also decreased considerably since 1990. Time series for subcategories are shown in Chapter 3.2.4.

The emission of SO<sub>2</sub> has decreased since 2005, but the emission level has steadied since 2009.

Table 3.2.2 SO<sub>2</sub> emission from stationary combustion plants, 2013<sup>1)</sup>

	SO <sub>2</sub> , Mg
1A1a Public electricity and heat production	3306
1A1b Petroleum refining	302
1A1c Oil and gas extraction	10
1A2 Industry	2822
1A4a Commercial/Institutional	116
1A4b Residential	896
1A4c Agriculture/Forestry/Fisheries	1161
<b>Total</b>	<b>8612</b>



1) Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.8 Disaggregated SO<sub>2</sub> emissions from 1A1a Public electricity and heat production.




Figure 3.2.9 SO<sub>2</sub> emission time series for stationary combustion.

### NO<sub>x</sub>

Stationary combustion accounts for 26% of the national NO<sub>x</sub> emission. Table 3.2.3 shows the NO<sub>x</sub> emission inventory for stationary combustion subcategories.

*Public electricity and heat production* is the largest emission source accounting for 45 % of the emission from stationary combustion plants. The emission from public power boilers > 300 MW<sub>th</sub> accounts for 31 % of the emission in this subcategory.

Industrial combustion plants are also an important emission source accounting for 16 % of the emission. The main industrial emission source is cement production, which accounts for 28 % of the emission.


Residential plants account for 12 % of the NO<sub>x</sub> emission. The fuel origin of this emission is mainly wood accounting for 62 % of the residential plant emission.

*Oil and gas extraction*, which is mainly off-shore gas turbines accounts for 19 % of the NO<sub>x</sub> emission.

Time series for NO<sub>x</sub> emission from stationary combustion are shown in Figure 3.2.10. NO<sub>x</sub> emission from stationary combustion plants has decreased by 73 % since 1990. The reduced emission is largely a result of the reduced emission from public electricity and heat production due to installation of low NO<sub>x</sub> burners, selective catalytic reduction (SCR) units and selective non-catalytic reduction (SNCR) units. The fluctuations in the time series follow the fluctuations in public electricity and heat production, which, in turn, result from electricity trade fluctuations.

The emission has decreased in recent years as shown in Figure 3.2.10.

Table 3.2.3 NO<sub>x</sub> emission from stationary combustion plants, 2013<sup>1)</sup>.



1) Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.10 NO<sub>x</sub> emission time series for stationary combustion.

## NMVOG

Stationary combustion plants account for 14 % of the national NMVOG emission. Table 3.2.4 presents the NMVOG emission inventory for the stationary combustion subcategories.

Residential plants are the largest emission source accounting for 80 % of the emission from stationary combustion plants. For residential plants NMVOG is mainly emitted from wood and straw combustion, see Figure 3.2.11.

Public electricity and heat production is also a considerable emission source, accounting for 8 % of the emission. Lean-burn gas engines have a relatively high NMVOG emission factor and are the most important emission source in this subcategory (see Figure 3.2.11). The gas engines are either natural gas or biogas fuelled.

Agricultural plants accounted for 9 % of the emission in 2013. Combustion of straw was the main emission source in this category.


The time series for NMVOG emission from stationary combustion is shown in Figure 3.2.12. The emission has decreased by 1 % from 1990. The emission increased until 2007 and decreased after 2007. The increased emission is mainly a result of the increasing wood consumption in residential plants and of the increased use of lean-burn gas engines in CHP plants. The decrease in recent years is a result of lower emission from residential wood combustion and the low number of operation hours for the lean burn gas engines.

The emission from residential plants increased 6 % since 1990. The NMVOG emission from wood combustion in 2013 was 46 % higher than the 1990 emission due to increased wood consumption. However, the emission factor has decreased since 1990 due to installation of modern stoves and boilers with improved combustion technology. Further, the emission from straw combustion in farmhouse boilers has decreased (43 %) over this period due to both a decreasing emission factor and decrease in straw consumption in this source category.

The use of wood in residential boilers and stoves was relatively low in 1998-99 resulting in a lower emission level.

The consumption of wood in residential plants increased until 2007. The improved technology that has been implemented in residential wood combustion have led to lower emission factors and thus decreasing NMVOG emission since 2007.

Table 3.2.4 NMVOC emission from stationary combustion plants, 2013<sup>1)</sup>.



1) Only emission from stationary combustion plants in the categories is included.




Figure 3.2.11 NMVOC emission from residential plants and from public electricity and heat production, 2013.




Figure 3.2.12 NMVOC emission time series for stationary combustion.

## CO

Stationary combustion accounts for 34 % of the national CO emission. Table 3.2.5 presents the CO emission inventory for stationary combustion subcategories.

Residential plants are the largest emission source, accounting for 78 % of the emission. Wood combustion accounts for 86 % of the emission from residential plants, see Figure 3.2.13. This is in spite of the fact that the fuel consumption share is only 45 %. Combustion of straw is also a considera-


ble emission source whereas the emission from other fuels used in residential plants is almost negligible.

The time series for CO emission from stationary combustion is shown in Figure 3.2.14. The emission has decreased by 21 % from 1990. The time series for CO from stationary combustion plants follow the time series for CO emission from residential plants.

The increase of wood consumption in residential plants in 1999-2007 is reflected in the time series for CO emission. The consumption of wood in residential plants in 2013 was 3.5 times the 1990 level. The decreased emission in 2007-2013 is a result of implementation of improved residential wood combustion technologies and the fact that the rapid increase of wood consumption until 2007 have stopped.

Both consumption and CO emission factor for have decreased for residential straw combustion plants since 1990.

Table 3.2.5 CO emission from stationary combustion plants, 2013<sup>1)</sup>.



<sup>1)</sup> Only emission from stationary combustion plants in the source categories is included.





Figure 3.2.13 CO emission sources, residential plants, 2013.



### Stationary combustion



### 1A4b Residential plants, fuel origin




Figure 3.2.14 CO emission time series for stationary combustion.

### NH<sub>3</sub>

Stationary combustion plants accounted for only 2 % of the national NH<sub>3</sub> emission in 2013.


Table 3.2.6 shows the NH<sub>3</sub> emission inventory for the stationary combustion subcategories. Residential plants account for 99 % of the emission. Wood combustion accounts for 99 % of the emission from residential plants.

The time series for the NH<sub>3</sub> emission is presented in Figure 3.2.15. The NH<sub>3</sub> emission has increased to 2.0 times the 1990 level.

The NH<sub>3</sub> emission for residential wood combustion have been recalculated according to emission factors from EEA (2013) resulting in higher emission values reported this year than the values reported for stationary combustion last year, see also Chapter 3.2.8.

Table 3.2.6 NH<sub>3</sub> emission from stationary combustion plants, 2013<sup>1)</sup>.

	NH <sub>3</sub> , Mg
1A1a Public electricity and heat production	16
1A1b Petroleum refining	-
1A1c Oil and gas extraction	-
1A2 Industry	-
1A4a Commercial/Institutional	-
1A4b Residential	1287
1A4c Agriculture/Forestry/Fisheries	-
<b>Total</b>	<b>1303</b>



1) Only the emission from stationary combustion plants in the source categories is included.




Figure 3.2.15 NH<sub>3</sub> emission time series, stationary combustion plants.

### Particulate matter (PM)

TSP from stationary combustion accounts for 19 % of the national emission. The emission shares for PM<sub>10</sub> and PM<sub>2.5</sub> are 48 % and 70 %, respectively.

Table 3.2.7 and Figure 3.2.16 show the PM emission inventory for the stationary combustion subcategories. Residential plants are the largest emission source accounting for 90 % of the PM<sub>2.5</sub> emission from stationary combustion plants.

The primary sources of PM emissions are:

- Residential boilers, stoves and fireplaces combusting wood
- Farmhouse boilers combusting straw
- Power plants primarily combusting coal
- Wood combusted in non-residential plants

The PM emission from wood combusted in residential plants is the predominant source. Thus, 86 % of the PM<sub>2.5</sub> emission from stationary combustion is emitted from residential wood combustion. This corresponds to 60 % of the national emission. A literature review (Nielsen et al., 2003) and a Nordic project (Sternhufvud et al., 2004) has demonstrated that the emis-

sion factor uncertainty for residential combustion of wood in stoves and boilers is notably high.

Figure 3.2.17 shows the fuel consumption and the PM<sub>2.5</sub> emission of residential plants. Wood combustion accounts for 95 % of the PM<sub>2.5</sub> emission from residential plants in spite of a wood consumption share of 45 %.

Emission inventories for PM have been reported for the years 2000-2013. The time series for PM emission from stationary combustion is shown in Figure 3.2.18. The emission of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> has increased 11 %, 12 % and 14 %, respectively, since year 2000. The increase is caused by the increased wood combustion in residential plants. However, the PM emission factors have decreased for this emission source category due to installation of modern stoves and boilers. The stabilisation of wood consumption in residential plants in 2007-2013 has resulted in a decrease of PM emission from stationary combustion in recent years.

The time series for PM emission from stationary combustion plants follows the time series for PM emission from residential plants.

Table 3.2.7 PM emission from stationary combustion plants, 2013<sup>1)</sup>.

		TSP, Mg	PM <sub>10</sub> , Mg	PM <sub>2.5</sub> , Mg
1A1a	Public electricity and heat production	930	731	580
1A1b	Petroleum refining	110	104	100
1A1c	Oil and gas extraction	2	1	1
1A2	Industry	351	263	182
1A4a	Commercial/Institutional	147	145	137
1A4b	Residential	14460	13733	13401
1A4c	Agriculture/Forestry/Fisheries	511	478	447
<b>Total</b>		<b>16511</b>	<b>15456</b>	<b>14849</b>

1) Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.16 PM emission sources, stationary combustion plants, 2013.




Figure 3.2.17 Fuel consumption and PM<sub>2.5</sub> emission from residential plants.





Figure 3.2.18 PM emission time series for stationary combustion.

### Black carbon (BC)

BC from stationary combustion accounts for 51 % of the national emission. Residential wood combustion is the main emission source accounting for 97 % of the emission from stationary combustion.

Table 3.2.8 shows the BC emission inventory for the stationary combustion subcategories.

Table 3.2.8 BC emission from stationary combustion plants, 2013<sup>1)</sup>.

	BC, Mg
1A1a Public electricity and heat production	383
1A1b Petroleum refining	15
1A1c Oil and gas extraction	0
1A2 Industry	80
1A4a Commercial/Institutional	42
1A4b Residential	1 899
1A4c Agriculture/Forestry/Fisheries	123
<b>Total</b>	<b>2 543</b>

1) Only emission from stationary combustion plants in the source categories is included.

### Heavy metals

Stationary combustion plants are among the most important emission sources for heavy metals. The emission share for stationary combustion compared to national total is shown for each metal in Table 3.2.9.

Table 3.2.9 and Figure 3.2.19 present the heavy metal emission inventory for the stationary combustion subcategories. The source categories *Public electricity and heat production*, *Residential* and *Industry* are the main emission sources. The emission share for waste incineration plants has decreased considerably since the year 2000 due to installation of new improved flue gas cleaning technology that was initiated based on lower emission limit values in Danish legislation (DEPA, 2011).

Table 3.2.9 Heavy metal emission from stationary combustion plants, 2013<sup>1)</sup>.

	As, kg	Cd, kg	Cr, kg	Cu, kg	Hg, kg	Ni, kg	Pb, kg	Se, kg	Zn, kg
1A1a Public electricity and heat production	149	32	218	196	171	375	390	731	552
1A1b Petroleum refining	31	22	23	45	22	260	66	106	90
1A1c Oil and gas extraction	3	0	0	0	2	0	0	0	0
1A2 Industry	74	24	83	106	70	1 292	437	80	983
1A4a Commercial/Institutional	3	1	4	4	2	25	5	1	15
1A4b Residential	12	410	728	214	23	91	961	17	16 359
1A4c Agriculture/Forestry/Fisheries	9	6	27	44	13	71	278	39	566
Total	281	494	1 083	610	303	2 115	2 138	975	18 564
Emission share from stationary combustion	76%	85%	73%	1%	85%	50%	18%	65%	33%

1) Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.19 Heavy metal emission sources, stationary combustion plants, 2013.

The time series for heavy metal emissions are provided in Figure 3.2.20. Emissions of all heavy metals have decreased considerably (33 % - 89 %) since 1990, see Table 3.2.10. Emissions have decreased despite increased incineration of waste. This has been made possible due to installation and improved performance of gas cleaning devices in waste incineration plants and also in large power plants, the latter being a further important emission source.

Table 3.2.10 Decrease in heavy metal emission 1990-2013.

Pollutant	As	Cd	Cr	Cu	Hg	Ni	Pb	Se	Zn
Decrease since 1990, %	76	49	80	83	89	87	86	75	33




Figure 3.2.20 Heavy metal emission time series, stationary combustion plants.



### Polycyclic aromatic hydrocarbons (PAH)

Stationary combustion plants accounted for more than 79 % of the PAH emission in 2013.

Table 3.2.11 and Figure 3.2.21 present the PAH emission inventories for the stationary combustion subcategories. Residential combustion is the largest emission source accounting for more than 74 % of the emission. Combustion of wood is the predominant source, accounting for more than 96 % of the PAH emission from residential plants, see Figure 3.2.22.

The time series for PAH emissions are presented in Figure 3.2.23. The increasing (15 % - 57 %) emission trend for PAH is a result of the increased combustion of wood in residential plants. The time series for wood combustion in residential plants is also provided in Figure 3.2.23. The stabilisation of the consumption of wood in residential plants since 2007 is reflected in the PAH emission time series.

Table 3.2.11 PAH emission from stationary combustion plants, 2013<sup>1)</sup>.

	Benzo(a)- Pyrene, kg	Benzo(b)- fluoranthene, kg	Benzo(k)- fluoranthene, kg	Indeno(1,2,3- c,d)pyrene, kg
1A1a Public electricity and heat production	9	36	23	6
1A1b Petroleum refining	0	0	0	0
1A1c Oil and gas extraction	0	0	0	0
1A2 Industry	20	72	14	3
1A4a Commercial/Institutional	154	202	67	110
1A4b Residential	1667	1667	616	934
1A4c Agriculture/Forestry/Fisheries	149	165	32	210
<b>Total</b>	<b>1999</b>	<b>2142</b>	<b>752</b>	<b>1264</b>
<b>Emission share from stationary combustion</b>	<b>88%</b>	<b>87%</b>	<b>79%</b>	<b>85%</b>

<sup>1)</sup> Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.21 PAH emission sources, stationary combustion plants, 2013.





Figure 3.2.22 PAH emission from residential combustion plants (stationary), fuel origin.



Combustion of wood in residential plants

Figure 3.2.23 PAH emission time series, stationary combustion plants. Comparison with wood consumption in residential plants.

### Polychlorinated dibenzodioxins and -furans (PCDD/F)


Stationary combustion plants accounted for 71 % of the national emission of polychlorinated dibenzodioxins and –furans (PCDD/F) in 2013.

Table 3.2.12 presents the PCDD/F emission inventories for the stationary combustion subcategories. In 2013, the emission from residential plants accounted for 81 % of the emission. Combustion of wood is the predominant source accounting for 87 % of the emission from residential plants (Figure 3.2.24).

The time series for PCDD/F emission is presented in Figure 3.2.25. The PCDD/F emission has decreased 67 % since 1990 mainly due to installation of dioxin filters in waste incineration plants. The emission from residential plants has increased due to increased wood consumption in this source category. However, the emission factor for residential wood combustion has decreased due to installation of modern stoves and boilers.

Table 3.2.12 PCDD/F emission from stationary combustion plants, 2013<sup>1)</sup>.

	PCDD/F, g I-teq
1A1a Public electricity and heat production	1.16
1A1b Petroleum refining	0.001
1A1c Oil and gas extraction	0.001
1A2 Industry	0.05
1A4a Commercial/Institutional	0.41
1A4b Residential	12.41
1A4c Agriculture/Forestry/Fisheries	1.31
<b>Total</b>	<b>15.36</b>



1) Only emission from stationary combustion plants in the source categories is included.




Figure 3.2.24 PCDD/F emission from residential plants, fuel origin.




Figure 3.2.25 PCDD/F emission time series, stationary combustion plants.

### Hexachlorobenzene (HCB)


Stationary plants accounted for more than 58 % of the estimated national emission of hexachlorobenzene (HCB) in 2013.

Table 3.2.13 shows the HCB emission inventory for the stationary combustion subcategories. *Public electricity and heat production* account for 80 % of the emission. Residential plants account for 12 % of the emission.

The time series for HCB emission is presented in Figure 3.2.26. The HCB emission has decreased 67 % since 1990 mainly due to improved flue gas cleaning in waste incineration plants. The emission from residential plants has increased due to increased wood consumption in this source category.

Table 3.2.13 HCB emission from stationary combustion plants, 2013<sup>1)</sup>.

	HCB, kg
1A1a Public electricity and heat production	1.246
1A1b Petroleum refining	0.0001
1A1c Oil and gas extraction	0.0000000
1A2 Industry	4
1A4a Commercial/Institutional	0.075
1A4b Residential	0.194
1A4c Agriculture/Forestry/Fisheries	0.045
<b>Total</b>	<b>1.567</b>



1) Only the emission from stationary combustion plants in the source categories is included.




Figure 3.2.26 HCB emission time series, stationary combustion plants.

### Polychlorinated biphenyls (PCB)

The previous Danish emission inventories did not include polychlorinated biphenyls (PCB) emissions.

PCB can be emitted in any chemical process involving chloride and organic carbon or emitted due to incomplete combustion of PCB in fuel (waste incineration). In Denmark, waste with high levels of PCB is only incinerated in plants with permission to incinerate this waste fraction as it requires a high combustion temperature.

Different references for PCB emissions are not directly comparable because some PCB emission data are reported for individual PCB congeners, some as a sum of a specified list of PCB congeners and some PCB emission data are reported as toxic equivalence (teq) based on toxicity equivalence factors (TEF) for 12 dioxin-like PCB congeners. The emission measurements reported by Thistlethwaite (2001a and 2001b) show that the emission of non-dioxin-like PCBs is high compared to the emission of dioxin-like PCBs.

Furthermore, teq values based on TEF are reported as WHO<sub>2005</sub>-teq or WHO<sub>1998</sub>-teq. This difference is however typically less than 50 %<sup>3</sup>.

For stationary combustion the emission inventory is a sum of dioxin-like PCB (dl-PCB) emission, no teq values applied.

Stationary plants accounted for 1 % of the estimated national PCB emission in 2013.

Table 3.2.14 shows the dl-PCB emission inventory for the stationary combustion subcategories. *Public electricity and heat production* account for 62 % of the emission. Residential plants account for 21 % of the emission.


The time series for dl-PCB emission is presented in Figure 3.2.26. The dl-PCB emission has decreased 62 % since 1990. The decrease is mainly a re-

<sup>3</sup> Data have been compared for a few datasets in which each dioxin-like PCB congener was specified

sult of the flue gas cleaning devices that have been installed in waste incineration plants for dioxin reduction.

Table 3.2.14 PCB emission from stationary combustion plants, 2013<sup>1)</sup>.

	PCB, kg
1A1a Public electricity and heat production	0.283
1A1b Petroleum refining	0.001
1A1c Oil and gas extraction	0.000
1A2 Industry	0.052
1A4a Commercial/Institutional	0.003
1A4b Residential	0.096
1A4c Agriculture/Forestry/Fisheries	0.018
<b>Total</b>	<b>0.452</b>



1) Only the emission from stationary combustion plants in the source categories is included.




Figure 3.2.26 PCB emission time series, stationary combustion plants.

### 3.2.4 Trend for subsectors



In addition to the data for stationary combustion, this chapter presents and discusses data for each of the subcategories in which stationary combustion is included. Time series are presented for fuel consumption and emissions.

#### 1A1 Energy industries


The emission source category *1A1 Energy Industries* consists of the subcategories:

- 1A1a Public electricity and heat production
- 1A1b Petroleum refining
- 1A1c Oil and gas extraction

Figure 3.2.27 – 3.2.31 present time series for the *Energy Industries*. *Public electricity and heat production* is the largest subcategory accounting for the main part of all emissions. Time series are discussed below for each subcategory.



### Natural gas fuelled engines



### Biogas fuelled engines





Figure 3.2.27 Time series for fuel consumption, 1A1 Energy industries.




Figure 3.2.28 Time series for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO emission, 1A1 Energy industries.




Figure 3.2.29 Time series for PM emission, 1A1 Energy industries.














Figure 3.2.30 Time series for HM emission, 1A1 Energy industries.




Figure 3.2.31 Time series for PAH, PCDD/F, HCB and dl-PCB emission, 1A1 Energy industries.

### **1A1a Public electricity and heat production**

Public electricity and heat production is the largest source category regarding fuel consumption for stationary combustion. Figure 3.2.32 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

The fuel consumption in public electricity and heat production was 4 % lower in 2013 than in 1990. As discussed in Chapter 3.2.2 the fuel consumption fluctuates mainly as a consequence of electricity trade. Coal is the fuel that is affected the most by the fluctuating electricity trade. Coal is the main fuel in the source category even in years with electricity import. The coal consumption in 2013 was 45 % lower than in 1990. Natural gas is also an important fuel and the consumption of natural gas has increased since 1990. A considerable part of the natural gas is combusted in gas engines (Figure 3.2.27). The consumption of waste and biomass has increased.

The SO<sub>2</sub> emission has decreased 97 % from 1990 to 2013. This decrease is a result of both lower sulphur content in fuels and installation and improved performance of desulphurisation plants. The emission was higher in 2013 than in 2012, but the emission has however decreased 57 % since 2005.

The NO<sub>x</sub> emission has decreased 85 % since 1990 due to installation of low NO<sub>x</sub> burners, selective catalytic reduction (SCR) units and selective non-catalytic reduction (SNCR) units. The fluctuations in time series follow the fluctuations in fuel consumption and electricity trade.

The emission of NMVOC in 2013 was 2.9 times the 1990 emission level. This is a result of the large number of gas engines that has been installed in Danish CHP plants. The decreasing emission in 2004-2013 is results of the time series for natural gas consumption in gas engines (Figure 3.2.27). In addition, the emission of NMVOC from engines decreased in 1995-2007 as a result of introduction of an emission limits for unburned hydrocarbon<sup>4</sup> (DEPA, 2005).

The CO emission was 28 % higher in 2013 than in 1990. The fluctuations follow the fluctuations of the fuel consumption. In addition, the emission from gas engines is considerable.

<sup>4</sup> Including methane.








Figure 3.2.32 Time series for 1A1a Public electricity and heat production.

### **1A1b Petroleum refining**

*Petroleum refining* is a small source category regarding both fuel consumption and emissions for stationary combustion. Presently two refineries are operating in Denmark. Figure 3.2.33 shows the time series for fuel consumption and emissions.

The significant decrease in both fuel consumption and emissions in 1996 is a result of the closure of a third refinery.

The fuel consumption has increased 5 % since 1990.

The emission of SO<sub>2</sub> has shown a pronounced decrease (71 %) since 1990, mainly because decreased consumption of residual oil (50 %) also shown in Figure 3.2.33. The increase in SO<sub>2</sub> emission in 1990-1992 also follows the residual oil consumption. The NO<sub>x</sub> emission in 2013 was 11 % lower in 2013 than in 1990. Since 2005, data for both SO<sub>2</sub> and NO<sub>x</sub> are plant specific data stated by the refineries.

The NMVOC emission time series follows the time series for fuel consumption.

A description of the Danish emission inventory for fugitive emissions from fuels is given in Plejdrup et al. (2011) and in Chapter 3.4.









Figure 3.2.33 Time series for 1A1b Petroleum refining.

### 1A1c Oil and gas extraction

The source category *Oil and gas extraction* comprises natural gas consumption in the off-shore industry and in addition a small consumption in the Danish gas treatment plant<sup>5</sup>. Gas turbines are the main plant type. Figure 3.2.34 shows the time series for fuel consumption and emissions.

The fuel consumption in 2013 was 2.5 times the consumption in 1990. The fuel consumption has decreased since 2008.

The emissions follow the increase of fuel consumption.

The decrease of CO emission in 2005 – 2007 is a result of a lower emission factor. This decrease of emission factor is valid for gas turbines in cogeneration plants, but might not be valid for off shore gas turbines. However, the same emission factors have been assumed for CO emission due to the lack of data from off shore gas turbines.




Figure 3.2.34 Time series for 1A1c Oil and gas extraction.

<sup>5</sup> Nybro.

## 1A2 Industry

*Manufacturing industries and construction* (Industry) consists of both stationary and mobile sources. In this chapter, only stationary sources are included.

The emission source category *1A2 Industry* consists of the subcategories:

- 1A2a Iron and steel
- 1A2b Non-ferrous metals
- 1A2c Chemicals
- 1A2d Pulp, Paper and Print
- 1A2e Food processing, beverages and tobacco
- 1A2f Non-metallic minerals
- 1A2 g viii Other manufacturing industry

Figure 3.2.35 - 3.2.39 show the time series for fuel consumption and emissions. The subsectors *Non-metallic minerals*, *Other manufacturing industry* and *Food processing, beverages and tobacco* are the main subsectors for fuel consumption and emissions.

The total fuel consumption in industrial combustion was 19 % lower in 2013 than in 1990. The consumption of natural gas has increased since 1990 whereas the consumption of coal has decreased. The consumption of residual oil has decreased, but the consumption of petroleum coke increased. The biomass consumption has increased 46 % since 1990.

The SO<sub>2</sub> emission has decreased 82 % since 1990. This is mainly a result of lower consumption of residual oil in the industrial sector (Figure 3.2.35). Further, the sulphur content of residual oil and several other fuels has decreased since 1990 due to legislation and tax laws.

The NO<sub>x</sub> emission has decreased 58 % since 1990 due to the reduced emission from industrial boilers in general. Cement production is the main emission source accounting for more than 50 % of the industrial emission in 1990-2009<sup>6</sup>. After 2009 the NO<sub>x</sub> emission from cement production was reduced considerably and in 2013, the NO<sub>x</sub> emission from cement industry was 28 % of the total emission from manufacturing industries and construction (stationary combustion). The NO<sub>x</sub> emission from cement production was reduced 78 % since 1990. The reduced emission is a result of installation of SCR on all production units at the cement production plant in 2004-2007<sup>7</sup> and improved performance of the SCR units in recent years. A NO<sub>x</sub> tax was introduced in 2010 (DMT, 2008).

The NMVOC emission has decreased 76 % since 1990. The decrease is mainly a result of decreased emission factor for combustion of wood in industrial boilers. The emission from gas engines has however increased considerably after 1995 due to the increased fuel consumption that is a result of the installation of a large number of industrial CHP plants (Figure

<sup>6</sup> More than 90 % of sector 1A2f i.

<sup>7</sup> To meet emission limit.



3.2.35). The NMVOC emission factor for gas engines is much higher than for boilers regardless of the fuel.

The CO emission in 2013 was 6 % lower than in 1990. The main sources of emission are combustion of wood and cement production. The CO emission from mineral wool production is included in the industry sector (2A6).

The large decrease of Hg emission since 2009 is related to a large decrease of particulate matter emission and to a large decrease of coal consumption since 2009.




Figure 3.2.35 Time series for fuel consumption, 1A2 Industry.




Figure 3.2.36 Time series for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO emission, 1A2 Industry.




Figure 3.2.37 Time series for PM emission, 1A2 Industry.




Figure 3.2.38 Time series for HM emission, 1A2 Industry.




Figure 3.2.39 Time series for PAH, PCDD/F, HCB and dioxin-like PCB emission, 1A2 Industry.

### 1A2a Iron and steel

*Iron and steel* is a very small emission source category. Figure 3.2.40 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

Natural gas is the main fuel in the subsector.




Figure 3.2.40 Time series for 1A2a Iron and steel.

### 1A2b Non-ferrous metals

*Non-ferrous metals* is a very small emission source category. Figure 3.2.41 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

Natural gas is the main fuel in the subsector. The consumption of residual oil has decreased and the SO<sub>2</sub> emission follows this decrease. The emissions of NO<sub>x</sub>, NMVOC and CO follow the fuel consumption.

The fuel consumption is very low after 2009. This is in agreement with the data reported by DEA to Eurostat (DEA, 2014c).




Figure 3.2.41 Time series for 1A2b Non-ferrous metals.

### 1A2c Chemicals

Chemicals is a minor emission source category. Figure 3.2.42 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

Natural gas is the main fuel in this subsector. The consumption of residual oil has decreased and the SO<sub>2</sub> emission follows this fuel consumption.




Figure 3.2.42 Time series for 1A2c Chemicals.

### 1A2d Pulp, paper and print

*Pulp, paper and print* is a minor emission source category. Figure 3.2.43 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

Natural gas - and since 2007 also wood - are the main fuels in the subsector. The consumption of coal and residual oil has decreased and this is reflected in the SO<sub>2</sub> emission time series. The increased consumption of wood since 2007 has resulted in a considerable increase in NMVOC and CO emission in 2007-2013.




Figure 3.2.43 Time series for 1A2d Pulp, paper and print.



### 1A2e Food processing, beverages and tobacco

Food processing, beverages and tobacco is a considerable industrial subsector. Figure 3.2.44 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

Natural gas and residual oil are the main fuels in the subsector. The consumption of residual oil has decreased and this is reflected in the SO<sub>2</sub> emission time series. The consumption of natural gas has increased.




Figure 3.2.44 Time series for 1A2e Food processing, beverages and tobacco.

### 1A2f Non-metallic minerals

*Non-metallic minerals* is a considerable industrial subsector. Figure 3.2.45 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO. The subsector includes cement production that is a major industrial emission source in Denmark.

Petroleum coke, natural gas, industrial waste and coal are the main fuels in the subsector in recent years. The consumption of coal and residual oil has decreased.

The NO<sub>x</sub> time series is discussed above on page 84.




Figure 3.2.45 Time series for 1A2f Non-metallic minerals

### 1A2g Other manufacturing industry

*Other manufacturing industry* is a considerable industrial subsector. Figure 3.2.45 shows the time series for fuel consumption and emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO.

Natural gas and wood are the main fuels in the subsector in recent years. The consumption of coal and residual oil has decreased.




Figure 3.2.45 Time series for 1A2g Other manufacturing industry.

### 1A4 Other Sectors



The emission source category *1A4 Other Sectors* consists of the subcategories:

- 1A4a Commercial/Institutional plants.
- 1A4b Residential plants.
- 1A4c Agriculture/Forestry.


Figure 3.2.46 – 3.2.50 present time series for this emission source category. *Residential plants* is the largest subcategory accounting for the largest part of all emissions. Time series are discussed below for each subcategory.

The HCB emission time series follows the fuel consumption of coal in residential plants. The HCB emission factor for coal used in residential plants is high compared to other fuels.


#### 1A4 Other Sectors




#### Gas engines, biogas (subsectors to Other Sectors)



#### Gas engines, natural gas (subsectors to Other Sectors)



#### Combustion of wood in Other Sectors



#### Combustion of straw in Other Sectors




Figure 3.2.46 Time series for fuel consumption, 1A4 Other Sectors.




Figure 3.2.47 Time series for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO emission, 1A4 Other Sectors.




Figure 3.2.48 Time series for PM emission, 1A4 Other Sectors.




Figure 3.2.49 Time series for HM emission, 1A4 Other Sectors.




Figure 3.2.50 Time series for PAH, PCDD/F, HCB and dioxin-like PCB emission, 1A4 Other Sectors.

#### **1A4a i Commercial and institutional plants**

The emission source category *Commercial and institutional plants* consists of both stationary and mobile sources. In this chapter, only stationary sources are included (1A4a i).

The fuel consumption and the emissions from commercial and institutional plants are low compared to the other stationary combustion emission source categories. Figure 3.2.51 shows the time series for fuel consumption and emissions.

The fuel consumption in commercial/institutional plants has decreased 32 % since 1990 and there has been a change of fuel type. The fuel consumption consists mainly of gas oil and natural gas. The consumption of gas oil has decreased and the consumption of natural gas has increased since 1990. The consumption of wood and biogas has also increased. The wood consumption in 2013 was 4.5 times the consumption in 1990 (see Figure 3.2.46).

The SO<sub>2</sub> emission has decreased 94 % since 1990. The decrease is a result of both the change of fuel from gas oil to natural gas and of the lower sulphur content in gas oil and in residual oil. The lower sulphur content (0.05 % for gas oil since 1995 and 0.7 % for residual oil since 1997) is a result of Danish tax laws (DEPA, 1998).

The NO<sub>x</sub> emission was 46 % lower in 2013 than in 1990. The decrease is mainly a result of the lower fuel consumption but also the change from gas oil to natural gas has contributed to the decrease. The emission from gas engines and wood combustion has increased.

The NMVOC emission in 2013 was 52 % higher than the 1990 emission level. The large increase is a result of the increased combustion of wood that is the main source of emission. The increased consumption of natural gas in gas engines (Figure 3.2.46) also contributes to the increased NMVOC emission.

The CO emission has decreased 19 % since 1990. The emission from wood and from natural gas fuelled engines and boilers have increased whereas the emission from gas oil has decreased. This is a result of the change of fuels used in the sector.










Figure 3.2.51 Time series for 1A4a Commercial /institutional.

### **1A4b i Residential plants**

The emission source category *Residential plants* consists of both stationary and mobile sources. In this chapter, only stationary sources are included (1A4b i). Figure 3.2.52 shows the time series for fuel consumption and emissions.

For residential plants, the total fuel consumption was 19 % lower in 2013 than in 1990. The large decrease from 2010 to 2011 was caused by higher temperature in the winter season of 2011. The consumption of gas oil has decreased since 1990 whereas the consumption of wood has increased considerably (3.5 times the 1990 level). The consumption of natural gas has also increased since 1990.

The large decrease (86 %) of SO<sub>2</sub> emission from residential plants is mainly a result of a change of sulphur content in gas oil since 1995. The lower sulphur content (0.05 %) is a result of Danish tax laws (DEPA, 1998). In addition, the consumption of gas oil has decreased and the consumption of natural gas that results in very low SO<sub>2</sub> emissions has increased.

The NO<sub>x</sub> emission has decreased by 16 % since 1990. The emission factor for wood is higher than for natural gas and gas oil and both consumption and the emission factor for wood have increased. However, the NO<sub>x</sub> emission factor for natural gas has decreased.

The emission of NMVOC has increased 6 % since 1990 as a result of the increased combustion of wood. The emission factor for wood has decreased since 1990, but not as much as the increase in consumption of wood. The emission factors for wood and straw are higher than for liquid or gaseous fuels.

The CO emission has decreased 10 % since 1990. The use of wood that is the main source of emission has increased whereas the emission factor has decreased. The emission from combustion of straw has decreased since 1990.








Figure 3.2.52 Time series for 1A4b Residential plants.

#### **1A4c i Agriculture/forestry**

The emission source category Agriculture/forestry/fishing consists of both stationary and mobile sources. In this chapter, only stationary sources are included (1A4c i). Figure 3.2.53 shows the time series for fuel consumption and emissions.

For plants in agriculture/forestry, the fuel consumption has decreased 41 % since 1990. A remarkable decrease of fuel consumption has taken place since year 2000.

The type of fuel that has been applied has changed since 1990. In the years 1994-2004, the consumption of natural gas was high, but in recent years, the consumption decreased again. A large part of the natural gas consumption has been applied in gas engines (Figure 3.2.46). Most CHP plants in agriculture/forestry based on gas engines came in operation in 1995-1999. The decrease in later years is a result of the liberalisation of the electricity market.

The consumption of straw has decreased since 1990. The consumption of both residual oil and gas oil has increased after 1990 but has decreased again in recent years.

The SO<sub>2</sub> emission was 64 % lower in 2013 than in 1990. The emission decreased mainly in the years 1996-2002. The main emission sources are coal, residual oil and straw.

The emission of NO<sub>x</sub> was 50 % lower in 2013 than in 1990.

The emission of NMVOC has decreased 42 % since 1990. The major emission source is combustion of straw. The consumption of straw has decreased since 1990. The emission from gas engines has increased mainly due to increased fuel consumption.

The CO emission has decreased 71 % since 1990. The major emission source is combustion of straw. In addition to the decrease of straw consumption, the emission factor for straw has also decreased since 1990.




Figure 3.2.53 Time series for 1A4c Agriculture/Forestry.

### 3.2.5 Methodological issues

The Danish emission inventory is based on the CORINAIR (CORe INventory on AIR emissions) system, which is a European program for air emission inventories. CORINAIR includes methodology structure and software for inventories. The methodology is described in the EMEP/EEA Guidebook (EEA 2013). Emission data are stored in an Access database, from which data are transferred to the reporting formats.

The emission inventory for stationary combustion is based on activity rates from the Danish energy statistics. General emission factors for various fuels, plants and sectors have been determined. Some large plants, such as power plants, are registered individually as large point sources and plant-specific emission data are used.

#### Large point sources

Large emission sources such as power plants, industrial plants and refineries are included as large point sources in the Danish emission database.

Each point source may consist of more than one part, e.g. a power plant with several units. By registering the plants as point sources in the database, it is possible to use plant-specific emission factors.

In the inventory for the year 2013, 78 stationary combustion plants are specified as large point sources. Plant specific emission data are available from 73 of the plants. The point sources include:

- Power plants and decentralised CHP plants.
- Waste incineration plants.
- Large industrial combustion plants.
- Petroleum refining plants.

The criteria for selection of point sources consist of the following:

- All centralized power plants, including smaller units.
- All units with a capacity of above 25 MW<sub>e</sub>.
- All district heating plants with an installed effect of 50 MW<sub>th</sub> or above and significant fuel consumption.
- All waste incineration plants obligated to report environmental data annually according to Danish law (DEPA, 2010b).
- Industrial plants,
  - With an installed effect of 50 MW<sub>th</sub> or above and significant fuel consumption.
  - With a significant process related emission.

The fuel consumption of stationary combustion plants registered as large point sources in the 2013 inventory was 265 PJ. This corresponds to 57 % of the overall fuel consumption for stationary combustion.

A list of the large point sources for 2013 is provided in Annex 3A-6. The number of large point sources registered in the databases increased from 1990 to 2013. Aggregated fuel consumption rates for the large point sources are also shown in Annex 3A-6.

The emissions from a point source are based either on plant specific emission data or, if plant specific data are not available, on fuel consumption data and the general Danish emission factors. Annex 3A-6 shows which of the emission data for large point sources are plant-specific and the corresponding share of the emission from stationary combustion.

The emission shares from point sources with plant specific data are shown in Table 3.2.15.

Table 3.2.15 Emission share, plant specific data.

Pollutant	Share from plant specific data, %
SO <sub>2</sub>	50%
NO <sub>x</sub>	42%
NMVOC	0.05%
CO	2.5%
NH <sub>3</sub>	0.6%
TSP	4.2%
PM <sub>10</sub>	3.8%
PM <sub>2.5</sub>	3.1%
As	33%
Cd	3%
Cr	11%
Cu	17%
Hg	59%
Ni	7%
Pb	7%
Se	68%
Zn	2.0%
PCDD/F	0.9%

SO<sub>2</sub> and NO<sub>x</sub> emissions from large point sources are often plant-specific based on continuous emission measurements. Emissions of CO, NMVOC, PM, heavy metals and PCDD/F are also plant-specific for some plants. Plant-specific emission data are obtained from:

- Annual environmental reports / environmental reporting available on the Danish EPA home page<sup>8</sup> (PRTR data), DEPA (2014)
- Annual plant-specific reporting of SO<sub>2</sub> and NO<sub>x</sub> from power plants >25MW<sub>e</sub> prepared for the Danish Energy Agency (DEA) and Energinet.dk
- Emission data reported by DONG Energy and Vattenfall, the two major power plant operators
- Emission data reported from industrial plants

Annual environmental reports for the plants include a considerable number of emission data sets. Emission data from annual environmental reports are, in general, based on emission measurements, but some emissions have potentially been calculated from general emission factors.

If plant-specific emission factors are not available, general area source emission factors are used.

#### Area sources

Fuels not combusted in large point sources are included as source category specific area sources in the emission database. Plants such as residential boilers, small district heating plants, small CHP plants and some industrial boilers are defined as area sources. Emissions from area sources are based on fuel consumption data and emission factors. Further information on emission factors is provided below.

<sup>8</sup> <http://www3.mst.dk/Miljoeoplysninger/PrtrPublicering/Index>

### **Activity rates, fuel consumption**

The fuel consumption rates are based on the official Danish energy statistics prepared by DEA. DCE aggregates fuel consumption rates to SNAP categories. Some fuel types in the official Danish energy statistics are added to obtain a less detailed fuel aggregation level cf. Annex 3A-3. The calorific values on which the energy statistics are based are also enclosed in Annex 3A-3. The calorific values shown in the annex are default values but plant specific reporting to the energy statistics is based on plant specific calorific values if data are available. The correspondence list between the energy statistics and SNAP categories is enclosed in Annex 3A-9.

The fuel consumption of the NFR category *Manufacturing industries and construction* (corresponding to SNAP category 03) is disaggregated into industrial subsectors based on the DEA data set aggregated for the Eurostat reporting (DEA, 2014c).

Both traded and non-traded fuels are included in the Danish energy statistics. Thus, for example, estimation of the annual consumption of non-traded wood is included.

Petroleum coke purchased abroad and combusted in Danish residential plants (border trade of 628 TJ in 2013) is not included in the Danish inventory. This is in agreement with the IPCC Guidelines (1996).

The fuel consumption data for large point sources refer to the EU Emission Trading Scheme (EU ETS) data for plants for which the Danish CO<sub>2</sub> emission inventory also refer to EU ETS.

For all other large point sources, the fuel consumption refers to a DEA database (DEA, 2014b). The DEA compiles a database for the fuel consumption of each district heating and power-producing plant, based on data reported by plant operators.

The fuel consumption of area sources is calculated as total fuel consumption minus fuel consumption of large point sources.

The Danish national energy statistics includes three fuels used for non-energy purposes, bitumen, white spirit and lubricants. The total consumption for non-energy purposes is relatively low, e.g. 11.6 PJ in 2013. The use of fuels for non-energy purposes is included in the inventory in sector 2D Non-energy products from fuels and solvent use, see Chapter 4.5.

In Denmark, all waste incineration is utilised for heat and power production. Thus, incineration of waste is included as stationary combustion in the source category *Energy*.

Fuel consumption data are presented in Chapter 3.2.2.

### **Town gas**

Town gas has been included in the fuel category natural gas. The consumption of town gas in Denmark is very low, e.g. 0.6 PJ in 2013. In 1990,



the town gas consumption was 1.6 PJ and the consumption has been steadily decreasing throughout the time series.

In Denmark, town gas is produced based on natural gas. The use of coal for town gas production ceased in the early 1980s.

An indicative composition of town gas according to the largest supplier of town gas in Denmark is shown in Table 3.2.16 (KE, 2015).

Table 3.2.16 Composition of town gas currently used (KE, 2015).

Component	Town gas, % (mol.)
Methane	43.9
Ethane	2.9
Propane	1.1
Butane	0.9
Carbon dioxide	0.4
Nitrogen	40.9
Oxygen	10.7

In earlier years, the composition of town gas was somewhat different. Table 3.2.17 is constructed with the input from Københavns Energi (KE) (Copenhagen Energy) and Danish Gas Technology Centre (DGC), (Jeppesen, 2007; Kristensen, 2007). The data refer to three measurements performed several years apart; the first in 2000 and the latest in 2005.

Table 3.2.17 Composition of town gas, information from the period 2000-2005.

Component	Town gas % (mol.)
Methane	22.3-27.9
Ethane	1.2-1.9
Propane	0.5-0.9
Butane	0.13-0.2
Higher hydrocarbons	0-0.9
Carbon dioxide	8-11.9
Nitrogen	15.6-20.9
Oxygen	2.3-3.2
Hydrogen	35.4-40.9
Carbon monoxide	2.6-2.9

Due to the scarce data available and the very low consumption of town gas compared to consumption of natural gas, the methodology will be applied unchanged in future inventories.

#### Emission factors

For each fuel and SNAP category (sector and e.g. type of plant), a set of general area source emission factors has been determined. The emission factors are either nationally referenced or based on the international guidebooks: EMEP/EEA Guidebook (EEA, 2013)<sup>9</sup> and IPCC Guidelines (IPCC, 2006).

A complete list of emission factors including time series and references is provided in Annex 3A-4.

<sup>9</sup> And former editions of the EMEP/EEA Guidebook.

### **SO<sub>2</sub> and NO<sub>x</sub> emission factors**

Emission factors for SO<sub>2</sub> and NO<sub>x</sub> are listed in Annex 3A-4. The appendix includes references and time series. Further details about the references, additional references, assumptions and discussions are included in Nielsen et al. (2014). An extract from this report have been enclosed in Annex 3A-12.

The emission factors refer to:

- The EMEP/EEA Guidebook: EEA (2013) and former editions.
- The IPCC Guidelines (IPCC, 2006).
- Danish legislation:
- Danish research reports including:
  - Two emission measurement programs for decentralised CHP plants (Nielsen et al. 2010; Nielsen & Illerup, 2003).
  - Research and emission measurements programs for biomass fuels
  - Research and environmental data from the gas sector
- Aggregated emission factors for residential wood combustion based on technology distribution (DEPA, 2013) and technology specific emission factors (EEA, 2013; DEPA, 2010a).
- Calculations based on plant-specific emissions from a considerable number of power plants.
- Calculations based on plant-specific emission data from a considerable number of waste incineration plants. These data refer to annual environmental reports published by plant operators.
- Sulphur content data from oil companies and the Danish gas transmission company, Energinet.dk.
- Additional personal communication.

Emission factor time series have been estimated for a considerable number of the emission factors. These are provided in Annex 3A-4.

### **NMVOC emission factors**

Emission factors for NMVOC are listed in Annex 3A-4. The annex includes references and time series. The emission factors for NMVOC refer to:

- An emission measurement program for decentralised CHP plants (Nielsen et al., 2010a).
- The EMEP/EEA Guidebook (EEA, 2013) and former editions.
- Aggregated emission factor based on the technology distribution for residential wood combustion and guidebook (EEA, 2013) emission factors. Technology distribution based on DEPA (2013).
- DGC Danish Gas Technology Centre 2001, Naturgas – Energi og miljø (DGC, 2001).
- Gruijthuijsen & Jensen (2000). Energi- og miljøoversigt, Danish Gas Technology Centre, 2000 (In Danish).

### **CO emission factors**

Emission factors for CO are listed in Annex 3A-4. The annex includes references and time series. The emission factors for CO refer to:

- The EMEP/EEA Guidebook (EEA, 2013) and the former editions.
- IPCC Guidelines (IPCC, 2006)

- An emission measurement program for decentralised CHP plants (Nielsen et al., 2010a).
- Danish legislation (DEPA, 2001)
- Aggregated emission factor based on the technology distribution for residential wood combustion and guidebook (EEA, 2013) emission factors. Technology distribution based on DEPA (2013).
- DCE estimate based on annual environmental reports for Danish waste incineration plants without power production, year 2000.
- Nikolaisen et al. (1998)
- Jensen & Nielsen (1990)
- Bjerrum (2002)
- Sander (2002)
- Gruijthuijsen & Jensen (2000)

### **NH<sub>3</sub> emission factors**

Emission factors have been included for residential wood combustion, residential straw combustion, waste incineration in public power production and residential combustion of coal, BKB and coke oven coke. The emission factor for waste incineration plants refers to a Danish emission measurement programme (Nielsen et al., 2010a) and all other emission factors refer to the EMEP/EEA Guidebook (EEA, 2009). Time series have been estimated for residential wood combustion.

### **Particulate matter (PM) emission factors**

Emission factors for PM and references for the emission factors are listed in Annex 3A-4. The emission factors are based on:

- The TNO/CEPMEIP emission factor database (CEPMEIP, 2001).

In addition, a considerable number of country-specific factors referring to:

- Danish legislation:
  - DEPA (2001), The Danish Environmental Protection Agency, Luftvejledningen (legislation from Danish Environmental Protection Agency).
  - DEPA (1990), The Danish Environmental Protection Agency, Bekendtgørelse 698 (legislation from Danish Environmental Protection Agency).
- Calculations based on plant-specific emission data from a considerable number of waste incineration plants.
- Aggregated emission factors for residential wood combustion based on technology distribution (DEPA, 2013) and technology specific emission factors (EEA, 2013; DEPA, 2010a).
- Two emission measurement programs for decentralised CHP plants (Nielsen et al., 2010a; Nielsen & Illerup, 2003).
- An emission measurement program for large power plants (Livbjerg et al., 2001).
- Research leading to the first Danish PM emission inventory for stationary combustion (Nielsen et al., 2003)
- Additional personal communication concerning straw combustion in residential plants.

Emission factor time series have been estimated for residential wood combustion and waste incineration. All other emission factors have been considered constant in 2000-2013.

#### **Black carbon (BC) emission factors**

Emission factors for BC all refer to EEA (2013). All emission factors are expressed as percentage of PM<sub>2.5</sub>. The applied emission factors and references are shown in Annex 3A-4.

Time series have been estimated for residential wood combustion and for waste incineration. The BC fraction of PM<sub>2.5</sub> is considered constant for each fuel/technology.

#### **Heavy metals emission factors**

Emission factors for 2013 for heavy metals (HM) are shown in Annex 3A-4. The annex includes references and time series. The emission factors refer to:

- Two emission measurement programs carried out on Danish decentralised CHP plants (Nielsen et al., 2010a; Nielsen & Illerup, 2003).
- Implied Emission Factors for power plants based on plant specific data reported by the power plant owners.
- Research concerning heavy metal emission factors representative for Denmark (Illerup et al., 1999).
- A CONCAWE study (Denier van der Gon & Kuenen, 2010)
- Data for Danish natural gas (Gruijthuijsen, 2001; Energinet.dk, 2012)
- Emission factors without national reference all refer to EEA (2009).

Time series have been estimated for coal and for waste incineration. For all other sources, the same emission factors have been applied for 1990-2013.

The heavy metal emission inventory has been documented in detail in Nielsen et al. (2013c).

#### **PAH emission factors**

Emission factors for PAH are shown in Annex 3A-4. The appendix includes references. The PAH emission factors refer to:

- Research carried out by TNO (Berdowski et al., 1995).
- Research carried out by Statistics Norway (Finstad et al., 2001).
- An emission measurement program performed on biomass fuelled plants. The project was carried out for the Danish Environmental Protection Agency (Jensen & Nielsen, 1996).
- Two emission measurement programs carried out on Danish decentralised CHP plants (Nielsen et al., 2010a; Nielsen & Illerup, 2003).
- Additional information from the gas sector (Jensen, 2001).

For residential wood combustion, country specific emission factors have been aggregated based on technology distribution in the sector (DEPA, 2013) and technology specific emission factors (EEA, 2013; DEPA 2010a).

Emission factor time series have been estimated for residential wood combustion, natural gas fuelled engines, biogas fuelled engines and waste in-

incineration plants. All other emission factors have been considered constant from 1990 to 2013. In general, emission factors for PAH are uncertain.

#### PCDD/F emission factors

Emission factors 2013 for PCDD/F are shown in Annex 3A-4.

The emission factor for residential wood combustion refers to technology specific emission factors (EEA 2013; DEPA 2010a) and to updated technology distribution data (DEPA, 2013).

The emission factors for decentralised CHP plants<sup>10</sup> refer to an emission measurement program for these plants (Nielsen et al. 2010a).

All other emission factors refer to research regarding PCDD/F emission carried out by NERI (now DCE) to prepare a new PCDD/F emission inventory (Henriksen et al., 2006).

Time series have been estimated for residential wood combustion and for incineration of waste. For all other sources, the same emission factors have been applied for 1990-2013.

#### HCB emission factors

The HCB emission inventory has been documented in Nielsen et al. (2014).

Table 3.2.18 shows the emission factors and references for the Danish emission factors.

Table 3.2.18 Emission factors for HCB, stationary combustion

Fuel	NFR (SNAP)	Emission factor, Reference ng/GJ
Coal	1A1, 1A2	6,700 Grochowalski & Koniecznyński (2008); EEA (2013)
Coal	1A4b	1,200,000 Syc et al. (2011)
Coal	1A4a and 1A4c	23,000 Syc et al. (2011)
Other solid fuels	1A1, 1A2	6,700 Assumed equal to coal.
Other solid fuels	1A4	1,200,000 Assumed equal to coal.
Liquid fuels <sup>1)</sup>	1A1, 1A2, 1A4	220 Nielsen et al. (2010)
Gaseous fuels	1A1, 1A2, 1A4	- Negligible
Waste	1A1, 1A2, 1A4	4300 Nielsen et al. (2010). A time series have been estimated. The emission factor for 1990 (190,000 ng/GJ) refer to Pacyna et al. (2003).
Wood	1A1, 1A2	5,000 EEA (2013)
Wood	1A4	5,000 EEA (2013)
Straw	1A1, 1A2	113 Nielsen et al. (2010)
Straw	1A4	5,000 EEA (2013)
Biogas	1A1, 1A2, 1A4	190 Nielsen et al. (2010)
Producer gas	1A1, 1A2, 1A4	800 Nielsen et al. (2010)

1) The emission factor for LPG and refinery gas is negligible.

<sup>10</sup> Natural gas fuelled engines, biogas fuelled engines, gasoil fuelled engines, engines fuelled by biomass producer gas, CHP plants combusting straw or wood and waste incineration plants.

For coal, the emission factor from Grochowalski & Koniecznyński (2008) is applied for energy industries and for industrial plants. This emission factor is also applied in the EEA Guidebook (EEA, 2013).

For residential plants, the emission factor 1,200,000 ng/GJ is applied referring to Syc et al. (2011). For commercial/institutional plants and for plants in agriculture / forestry the lower end of the value in Syc et al. (2011) (23,000 ng/GJ) is applied.

The emission factor for gas oil fuelled CHP engines (220 ng/GJ) referring to Nielsen et al. (2010) is applied for all liquid fuels except for LPG and refinery gas.

For gaseous fuels, LPG and refinery gas no data are available and the emission is negligible.

For waste combustion, emission data from Danish plants are available and these data are applied (Nielsen et al., 2010). The emission factor 4,300 ng/GJ is applied for 2005 onwards. The HCB emission factor for 1990 refers to Pacyna et al. (2003). The emission of HCB is related to emission of PCDD/F and the decline rate between 1990 and 2005 is based on the decline rate for PCDD/F.

Recent emission measurements from Polish industrial waste incineration plants confirms the emission factor level for waste incineration considering that the PCDD/F emission level is 15 times the PCDD/F emission level for Danish plants.

For wood combustion, the emission factors from EEA (2013) are applied for both energy industries, industrial plants and for non-industrial plants. For residential wood combustion, it would be relevant to estimate a time series. However, the currently available data are considered insufficient for this estimate.

The Cl content in straw is higher than in wood (Villeneuve et al., 2013) and thus the emission from straw combustion might potentially be higher. However, the emission factor for CHP plants combusting straw reported in Nielsen et al. (2010) is lower than the emission factor applied for wood.

The emission factor for energy industries and industrial combustion refer to Nielsen et al. (2010). For non-industrial plants, the EEA (2013) emission factor is applied.

The emission factors for biogas and producer gas both refer to Nielsen et al (2010).

#### **PCB emission factors**

The PCB emission inventory has been documented in Nielsen et al. (2014).

PCB emission is strongly related to the Cl content of the fuel (Syc et al., 2011) and to the emission level for PCDD/F (Hedman et al., 2006; Syc et al., 2011; Pandelova et al., 2009).

The Cl content of straw, bark and manure is higher than for wood (Ville-neuve et al., 2012). Villeneuve et al. (2012) states the Cl contents 50-60 mg/kg wood, 100-370 mg/kg bark, 1000-7000 mg/kg straw.

Different references for PCB emissions are not directly comparable because some PCB emission data are reported for individual PCB congeners, some as a sum of a specified list of PCB congeners and some PCB emission data are reported as toxic equivalence (teq) based on toxicity equivalence factors (TEF) for 12 dioxin-like PCB congeners. The emission measurements reported by Thistlethwaite (2001a and 2001b) show that the emission of non-dioxin-like PCBs is high compared to the emission of dioxin-like PCBs.

Furthermore, teq values based on TEF are reported as WHO<sub>2005</sub>-teq or WHO<sub>1998</sub>-teq. This difference is however typically less than 50 %<sup>11</sup>.

Table 3.2.19 shows the emission factors that have been selected for the Danish PCB emission inventory and reference for each emission factor. All emission factors are dioxin-like PCBs (but not teq values). PCB emission factors have been added for all fuels except LPG, refinery gas and natural gas. The emission from these three fuels is considered negligible.

<sup>11</sup> Data have been compared for a few datasets in which each dioxin-like PCB congener was specified.

Table 3.2.19 Emission factors for  $\Sigma$ dl-PCB, stationary combustion, 2013.

Fuel	NFR (SNAP)	Emission factor, $\Sigma$ dl-PCB, ng/GJ	Emission factor, PCB, ng WHO <sub>1998</sub> - teq/GJ	Reference
Coal	1A1	839	3.16	Grochowalski & Koniecznyński (2008)
Coal	1A2	5,700	53	Thistlethwaite (2001a)
Coal	1A4	7,403	66	Syc et al. (2011)
Other solid fuels	1A1	839	3.16	Assumed equal to coal.
Other solid fuels	1A2	5,700	53	Assumed equal to coal.
Other solid fuels	1A4	7,403	66	Assumed equal to coal.
Residual oil	1A1, 1A2, 1A4	839	3.2	The teq value refers to Dyke et al. (2003). The TEQ value is equal to the emission factor for coal combustion in power plants and the sum of dioxin-like PCB congeners has been assumed equal to the corresponding factor for coal.
Gas oil	1A1, 1A2, 1A4	93	0.11	Nielsen et al. (2010)
Other liquid fuels <sup>1)</sup>	1A1, 1A2, 1A4	93	0.11	Assumed equal to gas oil.
Gaseous fuels	1A1, 1A2, 1A4	-	-	Negligible
Waste	1A1, 1A2, 1A4	109 (time series)	0.28 (time series)	Nielsen et al. (2010). A time series have been estimated. The emission factor for 1990 (46,000 ng/GJ or 117 ng WHO1998teq/GJ) have been estimated based on the assumption that the PCB emission factor time series follow the PCDD/F time series.
Wood	1A1, 1A2, 1A4a/c	2,800	21	Thistlethwaite (2001a)
Wood	1A4b	2,752 (time series)	20.6)	Hedman et al. (2006). A time series have been estimated based on time series for technologies applied in Denmark.
Straw	1A1, 1A2	3,110	31.2	Assumed equal to residential plants.
Straw	1A4	3,110	31.2	Syc et al. (2011)
Biogas	1A1, 1A2, 1A4	90	0.13	Nielsen et al. (2010)
Producer gas	1A1, 1A2, 1A4	144	0.17	Nielsen et al. (2010)

1) Except LPG and refinery gas.

The emission factor for waste incineration refers to recent Danish field measurements. Historical data are not available, but a time series have been estimated based on the assumption that the dl-PCB emission factor follows the PCDD/-F emission factor. The estimated emission factor for 1990 is 45,671 ng/GJ or 117 ng WHO-teq/GJ. This emission level is confirmed by other references (Kakareka & Kukharchyk, 2005; Andrijewski et al., 2004). The emission factor time series is shown in Table 3.2.20.

For residential wood combustion, technology specific emission factors in toxicological equivalence are available from Hedman et al. (2006). However, sums of dioxin-like PCBs are not included in the reference. The emission factors for dioxin-like PCBs have been estimated based on the data for toxicological equivalence and the sum of dioxin-like PCBs in Thistlethwaite (2001a). Thus, the teq factors referring to Hedman (2006) have been multiplied by 2800/21. This assumption is highly uncertain, but the resulting emission factors seem to be in agreement with other references for residential wood combustion. A technology distribution time series for residential wood combustion in Denmark is available and have been ap-



plied for estimating the time series for the aggregated emission factor shown in Table 3.2.20.

Emission factor time series for waste incineration and for residential wood combustion are shown in Table 3.2.20.

Table 3.2.20 Emission factor time series for waste incineration and for residential wood combustion.

Year	Waste incineration		Residential wood combustion	
	$\Sigma$ dl-PCB, ng/GJ	dl-PCB, ng WHO-teq/ GJ	$\Sigma$ dl-PCB, ng/GJ	dl-PCB, ng WHO-teq/ GJ
1990	45671	117	6649	50
1991	38063	98	6623	50
1992	30433	78	6589	49
1993	22825	59	6560	49
1994	19773	51	6511	49
1995	16721	43	6446	48
1996	13690	35	6329	47
1997	10638	27	6232	47
1998	7586	19	6103	46
1999	5515	14	5696	43
2000	3423	9	5390	40
2001	3423	9	4834	36
2002	3423	9	4609	35
2003	3423	9	4548	34
2004	1766	4.5	4477	34
2005	109	0.28	4267	32
2006	109	0.28	4028	30
2007	109	0.28	4050	30
2008	109	0.28	3801	29
2009	109	0.28	3496	26
2010	109	0.28	3309	25
2011	109	0.28	3138	24
2012	109	0.28	2974	22
2013	109	0.28	2752	21

#### Emission factors for residential wood combustion

For the pollutants NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, PCDD/F, PCB and PAH emission factors have been based on fuel consumption data and emission factors for 10 different technologies. Technology categories, emission factors and implied emission factors for 2013 are shown in Table 3.2.21. For other pollutants, time series have not been estimated and the emission factors are shown in Annex 3A-4.

References and assumptions for each of the emission factors shown in Table 3.2.21 are included in Annex 3A-4.

Table 3.2.21 Technology specific emission factors for residential wood combustion.

Technology	NO <sub>x</sub> , g/GJ	NMVOC, g/GJ	CO, g/GJ	NH <sub>3</sub> , g/GJ	TSP, g/GJ	PM <sub>10</sub> , g/GJ	PM <sub>2.5</sub> , g/GJ	BC, g/GJ	PCDD/F, ng/GJ	dl-PCB, ng/GJ	Benzo (a) pyrene, mg/GJ	Benzo (b) fluoran- thene, mg/GJ	Benzo (k) fluoran- thene, mg/GJ	Indeno (1.2.3- c,d) pyrene, mg/GJ
Old stove	50	1200	8000	70	1000	950	930	93	800	7049	121	111	42	71
New stove	50	600	4000	70	800	760	740	74	800	7049	121	111	42	71
Stove according to resent Danish legislation (2008)	80	350	4000	37	640	608	608	97	250	931	61	56	21	36
Eco labelled stove / new advanced stove	95	175	1117	37	222	211	206	58	100	466	10	16	5	4
Other stoves	50	600	4000	70	800	760	740	74	800	7049	121	111	42	71
Old boilers with hot water storage	80	350	4000	74	1000	950	900	144	550	7049	121	111	42	71
Old boilers without hot water storage	80	350	4000	74	2000	1900	1800	288	550	7049	121	111	42	71
New boilers with hot water storage	95	175	1117	37	222	211	206	58	100	466	10	16	5	4
New boilers without hot water storage	95	350	2234	37	444	422	413	116	200	931	20	32	10	8
Pellet boilers	80	10	300	12	31	29	29	4	100	466	10	16	5	4
IEF residential wood combustion, 2013	75	338	2491	41	441	419	409	59	347	2752	52204	52192	19084	29455

### Implied emission factors

A considerable part of the emission data for waste incineration plants and large power plants are plant-specific. Thus, the area source emission factors do not necessarily represent average values for these plant categories. To attain a set of emission factors that expresses the average emission for power plants combusting coal and for waste incineration plants, implied emission factors have been calculated for these two plant categories. The implied emission factors are presented in Annex 3A-5. The implied emission factors are calculated as total emission divided by total fuel consumption.

#### 3.2.1 Uncertainty

According to the EEA Guidelines (EEA, 2013) uncertainty estimates should be estimated.

Uncertainty estimates include uncertainty with regard to the total emission inventory as well as uncertainty with regard to trends.

#### Methodology

The Danish uncertainty estimates are based on the simple Tier 1 approach.

The uncertainty estimates are based on emission data for the base year and year 2013 as well as on uncertainties for fuel consumption and emission factors for each of the NFR source categories. Residential plants have however been split in two parts: Residential wood combustion and other residential plants.

For particulate matter and BC, year 2000 is considered to be the base year, but for all other pollutants, the base year is 1990.

The uncertainty for fuel consumption in stationary combustion plants is based on EEA (2013). The uncertainties are shown in Table 3.2.22.

The applied uncertainties for activity rates and emission factors are based on EEA (2013). The uncertainty for emission factors that are based on recent Danish emission measurements are however estimated lower than suggested in the Guidebook. The applied uncertainties for emission factors are listed in Table 3.2.23.

Table 3.2.22 Uncertainty rates for fuel consumption, %.

Sector	%
1A1a Public electricity and heat production	1
1A1b Petroleum refining	1
1A1c_ii Oil and gas extraction	1
1A2 Manufacturing industries and construction	2
1A4a_i Commercial / institutional	3
1A4b_i Residential (excluding wood)	3
1A4b_i Residential wood	20
1A4c_i Agriculture / forestry / fishing	3

Table 3.2.23 Uncertainty rates for emission factors, %.

Sector	SO <sub>2</sub>	NO <sub>x</sub>	NMVOC	CO	PM	HM	PAH	HCB	Dioxin	NH <sub>3</sub>	PCB	BC
1A1a Public electricity and heat production	10	15	50	20	20	50	100	1000	200	1000	1000	1000
1A1b Petroleum refining	10	20	50	20	50	100	100	1000	1000	1000	1000	1000
1A1c_ii Oil and gas extraction	10	20	50	20	50	100	100	1000	1000	1000	1000	1000
1A2 Manufacturing industries and construction	10	20	50	20	30	100	100	1000	1000	1000	1000	1000
1A4a_i Commercial / institutional	20	50	50	50	50	300	1000	1000	1000	1000	1000	1000
1A4b_i Residential (excluding wood)	20	30	50	50	50	300	1000	1000	1000	1000	1000	1000
1A4b_i Residential wood	20	50	100	100	200	1000	1000	500	600	100	1000	1000
1A4c_i Agriculture/forestry /fishing	20	50	50	50	50	300	1000	1000	1000	1000	1000	1000

### Results

The uncertainty estimates for stationary combustion emission inventories are shown in Table 3.2.24. Detailed calculation sheets are provided in Annex 3A-7.

The total emission uncertainty is 6.1 % for SO<sub>2</sub> and 10 % for NO<sub>x</sub>.

Table 3.2.24 Uncertainty estimates, tier 1 approach, 2013.

Pollutant	Uncertainty	Trend	Uncertainty Trend,
	Total emission, %	1990-2013, %	%-age points
SO <sub>2</sub>	±6.1	-94	±0.3
NO <sub>x</sub>	±10	-73	±2
NMVOC	±69	-1	±30
CO	±69	-21	±31
NH <sub>3</sub>	±101	105	±82
TSP <sup>1)</sup>	±167	11	±30
PM <sub>10</sub> <sup>1)</sup>	±170	12	±30
PM <sub>2.5</sub> <sup>1)</sup>	±173	14	±30
BC <sup>1)</sup>	±741	116	±205
As	±46	-76	±7
Cd	±492	-49	±216
Cr	±266	-80	±51
Cu	±309	-83	±50
Hg	±41	-89	±3
Ni	±67	-87	±5
Pb	±165	-86	±22
Se	±42	-75	±3
Zn	±173	-33	±97
HCB	±799	-73	±42
PCDD/F	±445	-67	±119
Benzo(b)fluoranthene	±770	51	±237
Benzo(k)fluoranthene	±799	57	±157
Benzo(a)pyrene	±822	42	±202
Indeno(1,2,3-c,d)pyrene	±751	15	±285
PCB	±666	-62	±75

<sup>1)</sup> The base year for PM is year 2000.

### **3.2.7 Source specific QA/QC and verification**

An updated quality manual for the Danish emission inventories was published in 2013 (Nielsen et al., 2013). The quality manual describes the concepts of quality work and definitions of sufficient quality, critical control points and a list of Point for Measuring (PM). Details about the source specific QA/QC is included in Annex 3A-11.

Documentation concerning verification of the Danish emission inventories was published by Fauser et al. (2013).

A reviewed sector report for stationary combustion was published in 2014 (Nielsen et al., 2014b). Former editions of the sector report for stationary combustion have been reviewed by external experts in 2004, 2006 and 2009.

### **3.2.8 Source specific improvements and recalculations**

For stationary combustion plants, the emission estimates for the years 1990-2012 have been updated according to the latest energy statistics published by the Danish Energy Agency. The update included both end use and transformation sectors as well as a source category update. The changes in the energy statistics are largest for the years 2010, 2011 and 2012.

The consumption of wood in residential plants in 2012 is 4% lower in the revised energy statistics than in the energy statistics applied last year. This causes a lower emission of PM, PCDD/F, PAHs and PCB reported for 2012 this year.

A NH<sub>3</sub> emission factor for residential wood combustion has been added and this cause the higher NH<sub>3</sub> emission reported from stationary combustion this year.

The NMVOC emission factor for straw combusted in agricultural plants has been revised according to EEA (2013). This has caused a higher reported emission of NMVOC in this source category.

Double counting for SO<sub>2</sub> and NO<sub>x</sub> in mineral wool production plants in 1985-1994 have been corrected and cause a decrease for the industrial plants.

HM emission factors for residential wood combustion have been revised according to EEA (2013). This has caused higher reported emission values for Cd, Cr and Zn and lower reported emission values for As, Cu and Cu.

Emission factor time series for residential wood combustion have been estimated for 1990-2000. In previous inventories the emission factors were assumed constant in 1990-2000. This cause higher emission reported for residential plants for PAHs, PCDD/F, PCB, NMVOC and CO in 1990-2000.

An emission inventory for BC has been added.

And finally, the uncertainty estimates have been improved. The estimates are now based on NFR categories.

Recalculations for stationary combustion as a whole are shown in Table 3.2.25.

Table 3.2.25 Recalculations for stationary combustion, emissions reported in 2015 compared to emissions reported in 2014.

	1990	1995	2000	2005	2006	2007	2008	2009	2010	2011	2012
	Per cent										
SO <sub>2</sub>	99.6	103.6	100.0	100.3	100.2	101.1	100.1	99.8	100.5	101.1	99.6
NO <sub>x</sub>	99.6	100.0	100.0	100.0	100.0	100.0	100.1	100.3	100.2	100.4	100.1
NMVOG	113.3	117.7	109.6	108.0	107.9	107.0	107.5	108.1	107.7	108.8	106.6
CO	110.6	109.6	99.5	99.9	99.9	100.0	100.1	100.0	100.1	100.1	97.6
TSP			98.8	99.8	100.0	100.1	100.2	100.2	100.2	100.2	96.9
PM <sub>10</sub>			98.7	99.8	100.0	100.0	100.1	100.1	100.3	100.2	96.9
PM <sub>2.5</sub>			98.7	99.9	100.0	100.0	100.1	100.2	100.2	100.1	96.9
BC	New	New	New	New	New	New	New	New	New	New	New
NH <sub>3</sub>	947.4	1025.2	943.0	881.0	856.4	892.3	860.9	823.2	806.4	768.0	728.1
As	99.8	99.5	99.4	97.9	97.4	97.1	96.7	95.3	95.1	96.0	95.6
Cd	112.6	140.1	169.8	279.9	353.5	386.9	392.7	440.8	475.4	429.0	425.3
Cr	103.6	111.6	147.6	210.0	220.4	223.4	251.1	283.3	298.1	275.3	285.8
Cu	99.5	99.7	98.7	95.1	93.8	93.4	92.6	90.8	90.9	92.6	89.9
Hg	100.1	100.4	101.2	101.8	102.1	102.8	102.6	102.8	103.5	103.2	103.6
Ni	100.0	100.1	100.2	100.2	100.2	100.3	100.3	100.2	100.3	100.4	100.0
Pb	99.2	99.6	98.7	93.9	90.0	89.4	88.1	84.6	85.2	86.9	84.1
Se	100.0	100.0	100.1	100.1	100.0	100.1	100.1	100.0	100.3	100.4	100.1
Zn	115.4	128.8	178.4	258.2	283.5	310.3	322.3	344.4	346.4	330.9	331.7
HCB	100.0	100.1	100.4	100.3	100.2	100.4	100.4	100.2	100.3	100.1	99.9
PCDD/F	102.0	103.3	99.7	99.9	99.8	100.0	100.1	100.0	100.2	100.2	97.7
Benzo(a)pyrene	100.0	99.9	100.1	100.0	100.0	100.0	100.0	100.0	100.0	99.9	97.6
Benzo(b)fluoranthene	111.4	110.2	99.2	99.9	99.7	100.0	100.0	99.8	100.1	99.6	97.3
Benzo(k)fluoranthene	113.4	111.8	99.3	100.0	100.0	100.1	100.1	100.1	100.1	99.7	97.2
Indeno(123cd)pyrene	113.8	112.4	99.1	99.9	99.7	100.0	100.0	99.8	100.1	99.7	97.0
PCB	100.0	99.8	100.0	100.0	99.9	100.0	100.0	99.9	100.0	99.9	97.3

### 3.2.9 Source specific planned improvements

A number of improvements are planned for the stationary combustion inventories:

1) Improved documentation for emission factors.

The reporting of, and references for, the applied emission factors will be further developed in future inventories.

2) Improved documentation for uncertainty estimates.

The uncertainty estimates have been improved this year and the documentation will be further improved in the next inventory.

3) Improved presentation of results for BC

In this inventory BC have been included in the inventory and in the reporting next year the presentation of results will be improved.

### 3.2.10 References

Berdowski, J.J.M., Veldt, C., Baas, J., Bloos, J.P.J. & Klein, A.E. 1995: Technical Paper to the OSPARCOM-HELCOM-UNECE Emission Inventory of heavy Metals and Persistent Organic Pollutants, TNO-report, TNO-MEP – R 95/247.

CEPMEIP, 2001: The Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance. Online data available at: <http://www.air.sk/tno/cepmeip/> (2015-01-28).

Christiansen, B.H., Evald, A., Baadsgaard-Jensen, J. Bülow, K. 1997: Fyring med biomassebaserede restprodukter, Miljøprojekt nr. 358, 1997, Miljøstyrelsen. Available at: <http://www2.mst.dk/Udgiv/publikationer/1997/87-7810-778-4/pdf/87-7810-778-4.pdf> (2015-01-27)

Danish Energy Agency (DEA) 2014a: The Danish Energy Statistics, Available at: <http://www.ens.dk/info/tal-kort/statistik-nogletal/arlig-energistatistik> (2015-01-28).

Danish Energy Agency (DEA) 2014b: The Danish energy statistics, Energi-producenttællingen 2013. Unpublished.

Danish Energy Agency (DEA) 2014c: Energy statistics for industrial subsectors. Unpublished.

Danish Environmental Protection Agency (DEPA) 1990: Bekendtgørelse om begrænsning af emissioner af svovldioxid, kvælstofoxider og støv fra store fyringsanlæg, Bekendtgørelse 689 af 15/10/1990, (Danish legislation). Available at: <https://www.retsinformation.dk/Forms/R0710.aspx?id=48656> (2015-01-28)

Danish Environmental Protection Agency (DEPA) 1998: Bekendtgørelse om begrænsning af svovlindholdet i visse flydende og faste brændstoffer, Be-

kendtgørelse 698 af 22/09/1998 (Danish legislation). Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=12076> (2015-01-28).

Danish Environmental Protection Agency (DEPA) 2001: Luftvejledningen, Begrænsning af luftforurening fra virksomheder, Vejledning fra Miljøstyrelsen nr. 2 2001 (Danish legislation). Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=12728> (2015-01-28).

Danish Environmental Protection Agency (DEPA) 2005: Bekendtgørelse om begrænsning af emission af nitrogenoxider, uforbrændte carbonhydrider og carbonmonooxid mv. fra motorer og turbiner. Bekendtgørelse 621 af 23/06/2005. Danish legislation. Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=12836> (2015-01-28).

Danish Environmental Protection Agency (DEPA) 2010a: Emissioner fra træfyrede brændeovne/kedler (Emissions from wood fired stoves/boilers). Danish Environmental Protection Agency, 2010. Available at:  
<http://www2.mst.dk/udgiv/publikationer/2010/978-87-92617-85-9/pdf/978-87-92617-86-6.pdf> (2015-01-28).

Danish Environmental Protection Agency (DEPA) 2010b: Bekendtgørelse om visse virksomheders afgivelse af miljøoplysninger. Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=128761> (2015-01-28).

Danish Environmental Protection Agency (DEPA) 2011: Bekendtgørelse om anlæg, der forbrænder affald. Bekendtgørelse 1356 af 21/12/2011. Danish legislation. Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=139530> (2015-01-28).

Danish Environmental Protection Agency (DEPA), 2013: Forudsætninger for fremskrivning af emissioner fra brændeovne (In Danish). Hessberg & Justesen Miljøteknologi, J.nr. MST-5261-00007, Ref. stjus, Den 5. november 2013. (Related to e-mail correspondance 2011-04-28 and 2013-11-05).

Danish Environmental Protection Agency (DEPA) 2014: Annual environmental reports. Available at:  
<http://www3.mst.dk/Miljoeoplysninger/PrtrPublicering/Index> (2015-01-28).

Danish Gas Technology Centre (DGC) 2001: Naturgas – Energi og miljø (In Danish).

Danish Ministry of Taxation (DMT) 2008: Lov om afgift af kvælstofoxider. Lov nr 472 af 17/06/2008. Danish legislation. Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=120340> (2015-01-28).

Denier van der Gon, H. & Kuenen, J. 2010: Improvements to metal emission estimates, 10th Joint TFEIP/EIONET meeting, 11-12 May 2010, Vienna, Austria.

Energinet.dk, 2012: Energinet.dk homepage. Available at:  
<http://energinet.dk/DA/GAS/Gasdata-og-kvalitet/Gaskvalitet/Sider/Vis-gaskvalitet.aspx?Visning=aarsgennemsnit> (2015-01-28).



European Environment Agency (EEA), 2013: EMEP/EEA air pollutant emission inventory guidebook 2013. Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (2015-01-28).

European Environment Agency (EEA) 2009: EMEP/EEA air pollutant emission inventory guidebook – 2009. Available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (2015-01-28).

Fausser, P., Nielsen, M., Winther, M., Plejdrup, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Hoffmann, L., Thomsen, M., Hjelgaard, K. & Nielsen, O.-K. 2013: Verification of the Danish 1990, 2000 and 2010 emission inventory data. Aarhus University, DCE – Danish Centre for Environment and Energy, 85 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 79. Available at: <http://dce2.au.dk/pub/SR79.pdf> (2015-01-28).

Finstad, A., Haakonsen, G., Kvingedal, E. & Rypdal, K. 2001: Utslipp til luft av noen miljøgifter i Norge, Dokumentasjon av metode og resultater, Statistics Norway Report 2001/17 (In Norwegian). Available at: [http://www.ssb.no/a/publikasjoner/pdf/rapp\\_200117/rapp\\_200117.pdf](http://www.ssb.no/a/publikasjoner/pdf/rapp_200117/rapp_200117.pdf)

Gruijthuijsen, L.v. & Jensen, J.K. 2000: Energi- og miljøoversigt, Danish Gas Technology Centre 2000 (In Danish). Available at: <http://www.dgc.dk/publikation/2000/energi-og-miljoeoversigt> (2015-01-28).

Gruijthuijsen, L.v. 2001: Metaller i naturgas, Målerapport April 2001, Dansk Gasteknisk Center (in Danish). Available at: [http://www.dgc.dk/sites/default/files/filer/publikationer/R0104\\_metaller\\_naturgas.pdf](http://www.dgc.dk/sites/default/files/filer/publikationer/R0104_metaller_naturgas.pdf) (2015-01-28).

Henriksen, T.C., Illerup, J.B. & Nielsen, O.-K. 2006: Dioxin Air Emission Inventory 1990-2004. National Environmental Research Institute, Denmark. 90 pp. – NERI Technical report no 602. Available at: <http://www2.dmu.dk/Pub/FR602.pdf> (2015-01-28).

Illerup, J.B., Geertinger, A., Hoffmann, L. & Christiansen, K. 1999: Emissionsfaktorer for tungmetaller 1990-1996. Danmarks Miljøundersøgelser. 66 s. – Faglig rapport fra DMU nr. 301. (In Danish) Available at: [http://www2.dmu.dk/1\\_viden/2\\_Publikationer/3\\_fagrappporter/rapporter/fr301.pdf](http://www2.dmu.dk/1_viden/2_Publikationer/3_fagrappporter/rapporter/fr301.pdf) (2015-01-28).

Illerup, J.B., Henriksen, T.C., Lundhede, T., Breugel, C.v., Jensen, N.Z. 2007: Brændeovne og små kedler – partikelemission og reduktionstiltag. Miljøstyrelsen, Miljøprojekt 1164, 2007. Available at: <http://www2.mst.dk/Udgiv/publikationer/2007/978-87-7052-451-3/pdf/978-87-7052-452-0.pdf> (2015-01-28).

IPCC, 1996: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reference Manual (Volume 3). Available at: <http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.html> (2015-01-28).

Jensen, J. 2001: Danish Gas Technology Centre, Personal communication e-mail 11-10-2001.

Jensen, L. & Nielsen, P.A., 1990: Emissioner fra halm- og flisfyr, dk-Teknik & Levnedsmiddelstyrelsen 1990 (In Danish).

Jensen, L. & Nielsen, P.B. 1996: Emissioner fra halm- og flisfyr, Arbejdsrapport fra Miljøstyrelsen nr. 5 1996, Bilagsrapport (In Danish).

Jeppesen, J.S. 2007: København energi (Copenhagen Energy), Jørgen Steen Jeppesen, personal communication.

Københavns Energi (KE) 2015: Københavns energi (Copenhagen Energy) fact sheet on town gas. Available at:  
<http://www.ke.dk/portal/pls/portal/docs/346012.PDF> (2015-01-27).

Kristensen, P.G. 2007: Danish Gas Technology Centre, Per Gravers Kristensen, personal communication.

Livbjerg, H. Thellefsen, M. Sander, B. Simonsen, P., Lund, C., Poulsen, K. & Fogh, C.L. 2001: Feltstudier af Forbrændingsaerosoler, EFP -98 Projekt, Aerosollaboratoriet DTU, FLS Miljø, Forskningscenter Risø, Elsam, Energi E2 (in Danish).

Nielsen, M. & Illerup, J.B., 2003: Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme. Eltra PSO projekt 3141. Kortlægning af emissioner fra decentrale kraftvarmeværker. Delrapport 6. Danmarks Miljøundersøgelser. 116 s. – Faglig rapport fra DMU nr. 442. (In Danish, with an English summary). Available at:  
[http://www2.dmu.dk/1\\_viden/2\\_Publikationer/3\\_fagrapporter/rapporter/FR442.pdf](http://www2.dmu.dk/1_viden/2_Publikationer/3_fagrapporter/rapporter/FR442.pdf) (2015-01-28).

Nielsen, M. Winther, M., Illerup, J.B. & Mikkelsen, M.H. 2003: Danish emission inventory for particulate matter (PM). National Environmental Research Institute, Denmark. 126 p. – Research Notes from NERI No. 189. Available at:  
[http://www2.dmu.dk/1\\_viden/2\\_Publikationer/3\\_arbrapporter/rapporter/AR189.pdf](http://www2.dmu.dk/1_viden/2_Publikationer/3_arbrapporter/rapporter/AR189.pdf) (2015-01-28).

Nielsen, M., Nielsen, O.-K., Plejdrup, M. & Hjelgaard, K. 2010: Danish Emission Inventories for Stationary Combustion Plants. Inventories until 2008. National Environmental Research Institute, Aarhus University, Denmark. 236 pp. – NERI Technical Report No. 795. Available at:  
<http://www.dmu.dk/Pub/FR795.pdf> (2015-01-28).

Nielsen, M., Nielsen, O.-K. & Plejdrup, M. 2014. Danish emission inventories for stationary combustion plants. Inventories until 2011. Aarhus University, DCE – Danish Centre for Environment and Energy, 188 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 102  
<http://www.dce.au.dk/Pub/SR102.pdf> (2015-01-28)

Nielsen, M., Nielsen, O.-K. & Thomsen, M. 2010: Emissions from decentralised CHP plants 2007 - Energinet.dk Environmental project no. 07/1882. Project report 5 – Emission factors and emission inventory for decentralised CHP production. National Environmental Research Institute, Aarhus University. 113 pp. – NERI Technical report No. 786. Available at:  
<http://www.dmu.dk/Pub/FR786.pdf> (2015-01-28)

Nielsen, O.-K., Plejdrup, M.S., Winther, M., Gyldenkerne, S., Thomsen, M., Fauser, P., Nielsen, M. Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K.,

Hoffmann, L. & Bruun, H.G. 2013: Quality manual for the Danish greenhouse gas inventory. Version 2. Aarhus University, DCE – Danish Centre for Environment and Energy, 44 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 47. Available at: <http://www.dmu.dk/Pub/SR47.pdf> (2015-01-27).

Nielsen, M., Nielsen, O.-K. & Hoffmann, L. 2013. Improved inventory for heavy metal emissions from stationary combustion plants. 1990-2009. Aarhus University, DCE – Danish Centre for Environment and Energy, 111 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 68. <http://dce2.au.dk/pub/SR68.pdf> (2015-01-28)

Nikolaisen, L., Nielsen, C., Larsen, M.G., Nielsen, V. Zielke, U., Kristensen, J.K. & Holm-Christensen, B. 1998: Halm til energiformål, Teknik – Miljø – Økonomi, 2. udgave, 1998, Videncenter for halm og flisfyring (In Danish). [http://www.videncenter.dk/gule%20halm%20haefte/Gul\\_Dansk/halm-DK01.pdf](http://www.videncenter.dk/gule%20halm%20haefte/Gul_Dansk/halm-DK01.pdf)

Petterson, E., Boman, C., Westerholm, R., Boström, D., Nordin, A. 2011: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 2: Wood Stove. Energy Fuels 2011, 25, 315–323. Available at: <http://pubs.acs.org/doi/pdfplus/10.1021/ef1007787> (2015-01-28).

Plejdrup, M.S., Nielsen, O.-K. & Nielsen, M. 2009: Emission Inventory for Fugitive Emissions in Denmark. National Environmental Research Institute, Aarhus University, Denmark. 47 pp. – NERI Technical Report no. 739. Available at: <http://www.dmu.dk/pub/FR739.pdf> (2015-01-28).

Serup, H., Falster, H., Gamborg, C., Gundersen, P., Hansen, L. Heding, N., Jacobsen, H.H., Kofman, P., Nikolaisen, L. & Thomsen, I.M. 1999: Træ til energiformål, Teknik – Miljø – Økonomi, 2. udgave, 1999, Videncenter for halm og flisfyring (In Danish).

Sternhufvud, C., Karvosenoja, N., Illerup, J., Kindbom, K., Lükewille, A., Johansson, M. & Jensen, D. 2004: Particulate matter emissions and abatement options in residential wood burning in the Nordic countries.

Wit, J.d & Andersen, S.D. 2003: Emission fra større gasfyrede kedler, Dansk Gasteknisk Center, 2003. Available at: <http://www.dgc.dk/publikation/2003/emission-fra-stoerre-gasfyrede-kedler> (2015-01-28).

### 3.3 Transport and other mobile sources (NFR sector 1A2, 1A3, 1A4 and 1A5)

The emission inventory basis for mobile sources is fuel consumption information from the Danish energy statistics. In addition, background data for road transport (fleet and mileage), air traffic (aircraft type, flight numbers, origin and destination airports), national sea transport (fuel surveys, ferry technical data, number of return trips, sailing time) and non-road machinery (engine no., engine size, load factor and annual working hours) are used to make the emission estimates sufficiently detailed. Emission data mainly comes from the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2013). However, for railways, measurements specific to Denmark are used.

In the Danish emission database, all activity rates and emissions are defined in SNAP sector categories (Selected Nomenclature for Air Pollution), according to the CollectER system. The emission inventories are prepared from a complete emission database based on the SNAP sectors. The aggregation to the sector codes used for both the UNFCCC and UNECE Conventions is based on a correspondence list between SNAP and CRF/NFR classification codes shown in Table 3.3.1 below (mobile sources only).

Table 3.3.1 SNAP – CRF/NFR correspondence table for transport.

SNAP classification	CRF/NFR classification
07 Road transport	1A3bi Road transport: Passenger cars 1A3bii Road transport: Light duty vehicles 1A3biii Road transport: Heavy duty vehicles 1A3biv Road transport: Mopeds & motorcycles
0801 Military	1A5b Other, Mobile
0802 Railways	1A3c Railways
0803 Inland waterways	1A5b Other, Mobile
080402 National sea traffic	1A3dii National navigation (Shipping)
080403 National fishing	1A4ciii Agriculture/Forestry/Fishing: National fishing
080404 International sea traffic	1A3di (i) International navigation (Shipping)
080501 Dom. airport traffic (LTO < 1000 m)	1A3aai (i) Civil aviation (Domestic, LTO)
080502 Int. airport traffic (LTO < 1000 m)	1A3ai (i) Civil aviation (International, LTO)
080503 Dom. cruise traffic (> 1000 m)	1A3aai (ii) Civil aviation (Domestic, Cruise)
080504 Int. cruise traffic (> 1000 m)	1A3ai (ii) Civil aviation (International, Cruise)
0806 Agriculture	1A4cii Agriculture/Forestry/Fishing: Off-road agriculture/forestry
0807 Forestry	1A4cii Agriculture/Forestry/Fishing: Off-road agriculture/forestry
0808 Industry	1A2gvii Manufacturing industries/Construction (mobile)
0809 Household and gardening	1A4bii Residential: Household and gardening (mobile)
0811 Commercial and institutional	1A4aai Commercial/Institutional: Mobile

Military transport activities (land and air) refer to the CRF/NFR sector Other (1A5), the latter sector also including recreational craft (SNAP code 0803).

For aviation, Landing and Take Off ((LTO)<sup>1</sup> refers to the part of flying, which is below 1000 m. This part of the aviation emissions (SNAP codes 080501 and 080502) are included in the national emissions total as prescribed by the UNECE reporting rules. According to UNFCCC the national emissions for aviation comprise the emissions from domestic LTO (0805010) and domestic cruise (080503). The fuel consumption and emission development for aviation explained in the following are based on UNFCCC categorization, in order to be consistent with the Danish NIR report.

For mobile sources, internal database models for road transport, air traffic, sea transport and non-road machinery have been set up at DCE, Aarhus University, in order to produce the emission inventories. The output results from the DCE models are calculated in a SNAP format, as activity rates (fuel consumption) and emission factors, which are then exported directly to the central Danish CollectER database.

Apart from national inventories, the DCE models are used also as a calculation tool in research projects, environmental impact assessment studies, and to produce basic emission information which requires various aggregation levels.

### 3.3.1 Source category description

The following description of source categories explains the development in fuel consumption and emissions for road transport and other mobile sources.

#### Fuel consumption

Table 3.3.2 Fuel consumption (PJ) for domestic transport in 2013 in NFR sectors.

NFR ID	Fuel consumption (PJ)
Manufacturing industries/Construction (mobile)	14.0
Civil aviation (Domestic)	1.9
Road transport: Passenger cars	95.3
Road transport: Light duty vehicles	22.4
Road transport: Heavy duty vehicles	39.8
Road transport: Mopeds & motorcycles	0.9
Railways	3.3
National navigation (Shipping)	5.2
Commercial/Institutional: Mobile	2.3
Residential: Household and gardening (mobile)	0.9
Agriculture/Forestry/Fishing: Off-road agriculture/forestry	15.5
Agriculture/Forestry/Fishing: National fishing	6.9
Other, Mobile	3.3
Road transport total	158.3
Other mobile total	53.3
Domestic total	211.7
Civil aviation (International)	34.4
Navigation (international)	24.7

<sup>1</sup> A LTO cycle consists of the flying modes approach/descent, taxiing, take off and climb out. In principle the actual times-in-modes rely on the actual traffic circumstances, the airport configuration, and the aircraft type in question.

Table 3.3.2 shows the fuel consumption for domestic transport based on DEA statistics for 2013 in NFR sectors. The fuel consumption figures in time series 1985-2013 are given in Annex 2.B.16 (NFR format) and are shown for 2013 in Annex 2.B.15 (CollectER format). Road transport has a major share of the fuel consumption for domestic transport. In 2013 this sector's fuel consumption share is 75 %, while the fuel consumption shares for Off road agriculture/forestry and Manufacturing industries (mobile) are 7 %, in both cases. For the remaining sectors the total fuel consumption share is 11 %.




Figure 3.3.1 Fuel consumption per fuel type for domestic transport 1985-2013.




Figure 3.3.2 Fuel consumption share pr fuel type for domestic transport in 2013.

From 1985 to 2013, diesel (sum of diesel and biodiesel) and gasoline (sum of gasoline and E5) fuel consumption has changed by 56 % and - 10 %, respectively (Figure 3.3.1), and in 2013 the fuel consumption shares for diesel and gasoline were 66 % and 27 %, respectively (not shown). Other fuels only have a 7 % share of the domestic transport total (Figure 3.3.2). Almost all gasoline is used in road transportation vehicles. Gardening machinery and recreational craft are merely small consumers. Regarding diesel, there is considerable fuel consumption in most of the domestic transport categories,

whereas a more limited use of residual oil and jet fuel is being used in the navigation sector and by aviation (civil and military flights), respectively<sup>2</sup>.

### Road transport

As shown in Figure 3.3.3, the fuel consumption for road transport<sup>3</sup> has generally increased until 2007, except from a small fuel consumption decline noted in 2000. The impact of the global financial crisis on fuel consumption for road transport becomes visible for 2008 and 2009. The fuel consumption development is due to a decreasing trend in the use of gasoline fuels from 1999 onwards combined with a steady growth in the use of diesel until 2007. Within sub-sectors, passenger cars represent the most fuel-consuming vehicle category, followed by heavy-duty vehicles, light duty vehicles and 2-wheelers, in decreasing order (Figure 3.3.4).




Figure 3.3.3 Fuel consumption pr fuel type and as totals for road transport 1985-2013.

<sup>2</sup> Biofuels are sold at gas filling stations and are assumed to be used by road transport vehicles.

<sup>3</sup> The sum share of bioethanol and biodiesel in the gasoline and diesel fuel blends for road transport is 5.5 %, in 2013.




Figure 3.3.4 Total fuel consumption pr vehicle type for road transport 1985-2013.

As shown in Figure 3.3.5 fuel consumption for gasoline passenger cars dominates the overall gasoline consumption trend. The development in diesel fuel consumption in recent years (Figure 3.3.6) is characterised by increasing fuel consumption for diesel passenger cars, while declines in the fuel consumption for trucks and buses (heavy-duty vehicles) and light duty vehicles are noted for 2008- 2009, 2012, and 2008-2012, respectively.




Figure 3.3.5 Gasoline fuel consumption pr vehicle type for road transport 1985-2013.




Figure 3.3.6 Diesel fuel consumption pr vehicle type for road transport 1985-2013.



In 2013, fuel consumption shares for gasoline passenger cars, diesel passenger cars, heavy-duty vehicles and gasoline light duty vehicles were 33, 27, 25, 13 and 1 %, respectively (Figure 3.3.7).




Figure 3.3.7 Fuel consumption share (PJ) pr vehicle type for road transport in 2013.

### Other mobile sources

It must be noted that the fuel consumption figures behind the Danish inventory for mobile equipment in the agriculture, forestry, industry, household and gardening (residential), and inland waterways (part of navigation) sectors, are less certain than for other mobile sectors. For these types of machinery, the DEA statistical figures do not directly provide fuel consumption information, and fuel consumption totals are subsequently estimated from activity data and fuel consumption factors. For recreational craft the latest historical year is 2004.

As seen in Figure 3.3.8, classified according to CRF the most important sectors are Agriculture/forestry (1A4cii), Industry-other (mobile machinery part of 1A2g) and Navigation (1A3d). Minor fuel consuming sectors are Civil Aviation (1A3a), Railways (1A3c), Other (military mobile and recreational craft: 1A5b), Commercial/institutional (1A4a) and Residential (1A4b).

The 1985-2013 time series are shown pr fuel type in Figures 3.3.9-3.3.12 for diesel, gasoline and jet fuel, respectively.




Figure 3.3.8 Total fuel consumption in CRF sectors for other mobile sources 1985-2013.




Figure 3.3.9 Diesel fuel consumption in CRF sectors for other mobile sources 1985-2013.




Figure 3.3.10 Gasoline fuel consumption in CRF sectors for other mobile source 1985-2013.




Figure 3.3.11 Residual oil fuel consumption in CRF sectors for other mobile sources 1985-2013.




Figure 3.3.12 Jet fuel consumption in CRF sectors for other mobile sources 1985-2013.

In terms of diesel, the fuel consumption decreases for agricultural machines until 2000, due to a decline in the number of tractors and harvesters. After 2000, the increase in the engine sizes of new sold machines makes the total fuel consumption grow until 2008, whereas from 2008 onwards the turnover of old less fuel efficient machinery has more than outbalanced the prevailing trend towards smaller total stock numbers. The fuel consumption for industry has increased from the beginning of the 1990's, due to an increase in the activities for construction machinery. The fuel consumption increase has been very pronounced in 2005-2008, for 2009; however, the global financial crisis has a significant impact on the building and construction activities. For fisheries, the development in fuel consumption reflects the activities in this sector.

The Navigation sector comprises national sea transport (fuel consumption between two Danish ports including sea travel directly between Denmark and Greenland/Faroe Islands). For national sea transport, the diesel fuel consumption curve reflects the combination of traffic and ferries in use for regional ferries. From 1998 to 2000, a significant decline in fuel consumption is apparent. The most important explanation here is the closing of ferry service routes in connection with the opening of the Great Belt Bridge in 1997. The fuel consumption decreases in 2011 and 2012 are due to reductions in the number of round trips made by regional ferries. For railways, the gradual shift towards electrification explains the lowering trend in diesel fuel consumption and the emissions for this transport sector. The fuel consumed (and associated emissions) to produce electricity is accounted for in the stationary combustion part of the Danish inventories.

The largest gasoline fuel consumption is found for household and gardening machinery in the Commercial/Institutional (1A4a) and Residential (1A4b) sectors. Especially from 2001-2006, a significant fuel consumption increase is apparent due to considerable growth in the machinery stock. The decline in gasoline fuel consumption for Agriculture/forestry/fisheries (1A4c) is due to the gradual phasing out of gasoline-fuelled agricultural tractors.

In terms of residual oil there has been a substantial decrease in the fuel consumption for regional ferries. The fuel consumption decline is most significant from 1990-1992 and from 1997-1999.

The considerable variations from one year to another in military jet fuel consumption are due to planning and budgetary reasons, and the passing de-

mand for flying activities. Consequently, for some years, a certain amount of jet fuel stock-building might disturb the real picture of aircraft fuel consumption. Civil aviation has decreased until 2004, since the opening of the Great Belt Bridge in 1997, both in terms of number of flights and total jet fuel consumption. From 2011 to 2012 the total consumption of jet fuel decreased significantly due to a drop in the number of domestic flights.

### Bunkers

The residual oil and diesel oil fuel consumption fluctuations reflect the quantity of fuel sold in Denmark to international ferries, international warships, other ships with foreign destinations, transport to Greenland and the Faroe Islands, tank vessels and foreign fishing boats. For jet petrol, the sudden fuel consumption drop in 2002 is explained by the recession in the air traffic sector due to the events of September 11, 2001 and structural changes in the aviation business. In 2009, the impact of the global financial crisis on flying activities becomes very visible.




Figure 3.3.13 Bunker fuel consumption 1985-2013.

### Emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC

In Table 3.3.3 the SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC emissions for road transport and other mobile sources are shown for 2013 in NFR sectors. The emission figures in the time series 1985-2013 are given in Annex 2.B.16 (NFR format) and are shown for 2013 in Annex 2.B.15 (CollectER format).

From 1985 to 2013, the road transport emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, PM (exhaust emissions; all size fractions) and BC have decreased by 99, 57, 88, 85, 74 and 54 %, respectively (Figures 3.3.14-3.3.18), whereas the NH<sub>3</sub> emissions have increased by 1916 % during the same time period (Figure 3.3.19).

For other mobile sources, the emission changes for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO and PM (all size fractions) are -90, -29, -40, -5, -73 and -68 %, respectively (Figures 3.3.21-3.3.25). The NH<sub>3</sub> emissions have increased by 15 % during the same time period (Figure 3.3.26).

Table 3.3.3 Emissions of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC in 2013 for road transport and other mobile sources.

	SO <sub>2</sub> tonnes	NO <sub>x</sub> tonnes	NMVOC tonnes	CO tonnes	NH <sub>3</sub> tonnes	TSP tonnes	PM <sub>10</sub> tonnes	PM <sub>2.5</sub> tonnes	BC tonnes
Manufacturing industries/Construction (mobile)	6 7 454	1 079	6 164	3	613	613	613	613	433
Civil aviation (Domestic)	45	724	94	614	0	3	3	3	1
Road transport: Passenger cars	42 18 094	5 640	69 286	1 218	503	503	503	503	364
Road transport: Light duty vehicles	10 6 230	546	3 887	28	344	344	344	344	274
Road transport: Heavy duty vehicles	17 16 185	389	5 897	25	258	258	258	258	178
Road transport: Mopeds & motorcycles	0	144	1 322	7 550	1	26	26	26	4
Road transport: Gasoline evaporation	0	0	1 410	0	0	0	0	0	0
Road transport: Brake wear	0	0	0	0	0	497	487	194	13
Road transport: Tyre wear	0	0	0	0	0	877	526	369	134
Road transport: Road abrasion	0	0	0	0	0	1 080	540	292	0
Railways	2	2 361	164	372	1	70	70	70	46
National navigation (Shipping)	1 243	8 059	271	908	0	163	161	160	32
Commercial/Institutional: Mobile	1	219	3 601	72 587	0	67	67	67	3
Residential: Household and gardening (mobile)	0	92	1 924	26 477	0	15	15	15	1
Agriculture/Forestry/Fishing: Off-road agriculture/forestry	7	7 257	1 545	16 294	3	584	584	584	353
Agriculture/Forestry/Fishing: National fishing	323	9 139	400	1 318	0	149	147	147	46
Other, Mobile	72	1 513	376	3 881	1	99	99	99	39
Road transport exhaust total	69 40 654	9 307	86 619	1 272	1 131	1 131	1 131	1 131	821
Road transport non exhaust total	0	0	0	0	0	2 454	1 554	854	147
Other mobile sources total	1 700	36 819	9 454	128 615	8	1 763	1 760	1 758	954
Domestic total	1 769	77 473	18 761	215 234	1 280	5 348	4 444	3 743	1 922
Civil aviation (International)	790	12 636	163	1 898	0	40	40	40	19
Navigation (International)	7 140	46 358	1 510	4 982	0	835	827	822	146

### Road transport

The step-wise lowering of the sulphur content in diesel fuel has given rise to a substantial decrease in the road transport emissions of SO<sub>2</sub> (Figure 3.3.14). In 1999, the sulphur content was reduced from 500 ppm to 50 ppm (reaching gasoline levels), and for both gasoline and diesel the sulphur content was reduced to 10 ppm in 2005. Since Danish diesel and gasoline fuels have the same sulphur percentages, at present, the 2013 shares for SO<sub>2</sub> emissions and fuel consumption for passenger cars, heavy-duty vehicles, light-duty vehicles and 2-wheelers are the same in each case: 60, 25, 14 and 1 %, respectively (Figure 3.3.20).




Figure 3.3.14 SO<sub>2</sub> emissions (tonnes) pr vehicle type for road transport 1985-2013.

Historically, the emission totals of NMVOC and CO have been very dominated by the contributions coming from private cars, as shown in Figures 3.3.16- 3.3.17. However, the NMVOC and CO (and NO<sub>x</sub>) emissions from this vehicle type have shown a steady decreasing tendency since the introduction of private catalyst cars in 1990 (EURO I) and the introduction of even more emission-efficient EURO II, III, IV and V private cars (introduced in 1997, 2001, 2006 and 2011, respectively).

For NO<sub>x</sub> the emission decrease for passenger cars is composed of a significant drop in emissions from gasoline cars driven by technology improvements, and an increase in emissions from diesel cars due to the dieselization of the Danish vehicle fleet, and almost unchanged emission factors for diesel passenger cars throughout the period regardless of EU emission legislation demands. For light duty vehicles the NO<sub>x</sub> emission trend is also the result of a technology driven emission reduction for gasoline vehicles, and a traffic induced emission increase for diesel vehicles; the emission factors for the latter vehicle category have been relatively constant over the years just as for diesel cars.

For heavy duty vehicles the real traffic emissions are not reduced in the order as intended by the EU emission legislation. Most markedly for Euro II engines the emission factors are even higher than for Euro I due to the so-called engine cycle-beating effect. Outside the legislative test cycle stationary measurement points, the electronic engine control for heavy duty Euro II engines switches to a fuel efficient engine running mode, thus leading to increasing NO<sub>x</sub> emissions (Figure 3.3.15). However, the reduction in transport activities due to the global financial crisis causes the NO<sub>x</sub> emissions for heavy duty vehicles to decrease significantly in 2008 and 2009.

Exhaust particulate emissions from road transportation vehicles are well below PM<sub>2.5</sub>. The emissions from light- and heavy-duty vehicles have significantly decreased since the mid-1990s due to gradually stricter EURO emission standards. In recent years until 2008 the environmental benefit of introducing gradually cleaner diesel private cars has been somewhat outbalanced by an increase in sales of new vehicles. After 2008 the PM emissions gradually become lower due to the increasing number of Euro V cars equipped with particulate filter sold in Denmark from 2006 onwards (Figure 3.3.18).

An undesirable environmental side effect of the introduction of catalyst cars is the increase in the emissions of NH<sub>3</sub> from the first two generations of catalyst cars (Euro I and II) compared to conventional cars. The emission factors for later catalytic converter technologies are considerably lower than the ones for Euro I and II, thus causing the emissions to decrease from 2001 onwards (Figure 3.3.19).

The 2013 emission shares for heavy-duty vehicles, passenger cars, light-duty vehicles and 2-wheelers for NO<sub>x</sub> (40, 45, 15 and 0 %), NMVOC (4, 62, 7 and 13 %), CO (6, 80, 6 and 8 %), PM (21, 45, 32 and 2 %) and NH<sub>3</sub> (1, 97, 2 and 0 %), are also shown in Figure 3.3.20.




Figure 3.3.15 NO<sub>x</sub> emissions (tonnes) pr vehicle type for road transport 1985-2013.




Figure 3.3.16 NMVOC emissions (tonnes) pr vehicle type for road transport 1985-2013.




Figure 3.3.17 CO emissions (tonnes) pr vehicle type for road transport 1985-2013.




Figure 3.3.18 PM emissions (tonnes) pr vehicle type for road transport 1985-2013.




Figure 3.3.19 NH<sub>3</sub> emissions (tonnes) pr vehicle type for road transport 1985-2013.






Figure 3.3.20 SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, PM and BC emission shares per vehicle type for road transport in 2013.

### Non-exhaust emissions of TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC

Apart from the exhaust emission estimates of particulate matter (PM), the Danish emission inventories also comprise the non-exhaust PM emissions coming from road transport brake and tyre wear, and road abrasion.

In Table 3.3.3, the non-exhaust TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC emissions for road transport are shown for 2013 in NFR sectors. The activity data and emission factors are also shown in Annex 2.B.15.

The respective source category distributions for TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions are identical for each of the non-exhaust emission type's brake wear, tyre wear and road abrasion, and, hence, only the PM<sub>10</sub> distributions are shown in Figure 3.3.28. Passenger cars caused the highest emissions in 2013, followed by trucks, light-duty vehicles, buses and 2-wheelers.




Figure 3.3.28 Brake and tyre wear and road abrasion PM<sub>10</sub> emission shares and PM and BC exhaust/non-exhaust distributions for road traffic in 2013.

Figure 3.3.28 also shows the exhaust/non-exhaust distribution of the total particulate emissions from road transport, for each of the size classes TSP, PM<sub>10</sub> and PM<sub>2.5</sub> and for BC. The exhaust emission shares of total road transport TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and BC are 26, 35, 50 and 85 %, respectively, in 2013. For brake and tyre wear and road abrasion the TSP shares are 15, 27 and 33 %, respectively. The same three sources have PM<sub>10</sub> shares of 20, 22 and 22 %, respectively, PM<sub>2.5</sub> shares of 11, 22 and 17 %, and BC shares of 1, 14 and 0 %, respectively. In general, the non-exhaust shares of total particulate emissions are expected to increase in the future as total exhaust emissions decline. The latter emission trend is due to the stepwise strengthening of exhaust emission standards for all vehicle types.

#### Other mobile sources

For SO<sub>2</sub> the trends in the Navigation (1A3d) emissions shown in Figure 3.3.21 mainly follow the development of the heavy fuel oil consumption (Figure 3.3.11). Though, from 1993 to 1995 relatively higher contents of sulphur in the fuel (estimated from sales) cause a significant increase in the emissions of SO<sub>2</sub>. The SO<sub>2</sub> emissions for Fisheries (1A4c) correspond with the development in the consumption of marine gas oil. The main explanation for the development of the SO<sub>2</sub> emission curves for Railways (1A3c) and non-road machinery in Agriculture/forestry (1A4c) and Industry (1A2f), are the stepwise sulphur content reductions for diesel used by machinery in these sectors.




Figure 3.3.21 SO<sub>2</sub> emissions (ktonnes) in NFR sectors for other mobile sources 1985-2013.

In general, the emissions of NO<sub>x</sub>, NMVOC and CO from diesel-fuelled working equipment and machinery in agriculture, forestry and industry have decreased slightly since the end of the 1990s due to gradually strengthened emission standards given by the EU emission legislation directives. For industry, the emission impact from the global financial crisis becomes very visible for 2009.

NO<sub>x</sub> emissions mainly come from diesel machinery, and the most important sources are Agriculture/forestry/fisheries (1A4c), Navigation (1A3d), Industry (1A2f) and Railways (1A3c), as shown in Figure 3.3.22. The 2013 emission shares are 45, 22, 20 and 6 %, respectively (Figure 3.3.27). Minor emissions come from the sectors, Civil Aviation (1A3a), Other (1A5) and Residential (1A4b).

The NO<sub>x</sub> emission trend for Navigation, Fisheries and Agriculture is determined by fuel consumption fluctuations for these sectors, and the development of emission factors. For ship engines the emission factors tend to increase for new engines until mid-1990s. After that, the emission factors gradually reduce until 2000, bringing them to a level comparable with the emission limits for new engines in this year. For agricultural machines, there have been somewhat higher NO<sub>x</sub> emission factors for 1991-stage I machinery, and an improved emission performance for stage I and II machinery since the late 1990s.

The emission development from 1985 to 2008 for industry NO<sub>x</sub> is the product of a fuel consumption increase, most pronounced from 2005-2008, and a development in emission factors as explained for agricultural machinery. For railways, the gradual shift towards electrification explains the declining trend in diesel fuel consumption and NO<sub>x</sub> emissions for this transport sector until 2001.




Figure 3.3.22 NO<sub>x</sub> emissions (tonnes) in NFR sectors for other mobile sources 1985-2013.

The 1985-2013 time series of NMVOC and CO emissions are shown in Figures 3.3.23 and 3.3.24 for other mobile sources. The 2013 sector emission shares are shown in Figure 3.3.27. For NMVOC, the most important sectors are Commercial/Institutional (1A4a), Agriculture/forestry/-fisheries (1A4c), Residential (1A4b) and Industry (1A2g) with 2013 emission shares of 38, 20, 20 and 12 %, respectively. The same five sectors also contribute with most of the CO emissions. For Commercial/Institutional (1A4a), Residential (1A4b), Agriculture/forestry/fisheries (1A4c) and Industry (1A2g) the emission shares are 56, 21, 14 and 5 %, respectively. Minor NMVOC and CO emissions come from Navigation (1A3d), Railways (1A3c), Civil Aviation (1A3a) and Other (1A5).

For NMVOC and CO, the significant emission increases for the commercial/institutional and residential sectors after 2000 are due to the increased number of gasoline working machines. Improved NMVOC emission factors for diesel machinery in agriculture and gasoline equipment in forestry (chain saws) are the most important explanations for the NMVOC emission decline in the Agriculture/forestry/fisheries sector. This explanation also applies for the industrial sector, which is dominated by diesel-fuelled machinery. From 1997 onwards, the NMVOC emissions from Navigation decrease due to the gradually phase-out of the 2-stroke engine technology for recreational craft. The main reason for the significant 1985-2006 CO emission decrease for Agriculture/forestry/-fisheries is the phasing out of gasoline tractors.

As shown in Figure 3.3.27, for other mobile sources the largest TSP contributors in 2013 are Agriculture/forestry/fisheries (1A4c), Industry (1A2f) and Navigation (1A3d), with emission shares of 41, 35 and 9 %, respectively. The remaining sectors: Railways (1A3c), Civil aviation (1A3a), Other (1A5) and Residential (1A4b) represent only minor emission sources.

The 1985-2013 TSP emissions for navigation and fisheries are determined by the fuel consumption fluctuations in these years, and the development of the emission factors, which to a major extent is a function of the fuel sulphur content. The emission development for Agriculture/forestry is determined by the generally decreasing total diesel fuel consumption and gradually reducing emission factors over the time period.

The TSP emission development for industrial non-road machinery is the product of a fuel consumption increase from 1985 to 2008 and a development in emission factors, as explained for agricultural machinery. The TSP emission explanations for railways are the same as for NO<sub>x</sub> (Figure 3.3.22).

The amounts of NH<sub>3</sub> emissions calculated for other mobile sources are very small. The largest emission sources are Agriculture-/forestry/fisheries (1A4c), Industry (1A2f), Other (1A5b) and Railways (1A3c), with emission shares of 44, 32, 10 and 9 %, respectively.




Figure 3.3.23 NMVOC emissions (tonnes) in NFR sectors for other mobile sources 1985-2013.




Figure 3.3.24 CO emissions (tonnes) in NFR sectors for other mobile sources 1985-2013.




Figure 3.3.25 TSP emissions (tonnes) in NFR sectors for other mobile sources 1985-2013.




Figure 3.3.26 NH<sub>3</sub> emissions (tonnes) in NFR sectors for other mobile sources 1985-2013.




Figure 3.3.27 SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, PM and BC emission shares per vehicle type for other mobile sources in 2013.

### Heavy metals

In Table 3.3.4, the heavy metal emissions for road transport and other mobile sources are shown for 2013 in NFR sectors. The emission figures in the time series 1990-2013 are given in Annex 2.B.16 (NFR format) and are shown for 1990 and 2013 in Annex 2.B.15 (CollectER format).

Table 3.3.4 Heavy metal emissions in 2013 for road transport and other mobile sources.

	Arsenic kg	Cadmium kg	Chromium kg	Copper kg	Mercury kg	Nickel kg	Lead kg	Selenium kg	Zinc kg
Manufacturing industries/Construction (mobile)	0	3	9	7	2	3	15	0	513
Civil aviation (Domestic)	0	0	0	0	0	0	876	0	3
Road transport: Passenger cars	0	28	60	86	15	30	122	0	5 578
Road transport: Light duty vehicles	0	5	16	12	3	5	28	0	960
Road transport: Heavy duty vehicles	0	6	21	15	5	6	34	0	1 138
Road transport: Mopeds & motorcycles	0	0	0	0	0	0	0	0	20
Road transport: Gasoline evaporation	0	0	0	0	0	0	0	0	0
Road transport: Brake wear	5	4	56	39 913	0	55	5 169	10	8 464
Road transport: Tyre wear	1	2	3	14	0	22	71	18	9 595
Road transport: Road abrasion	0	0	21	11	0	17	51	0	82
Railways	0	1	2	2	0	1	4	0	125
National navigation (Shipping)	31	2	14	31	5	1 664	18	36	84
Commercial/Institutional: Mobile	0	1	1	2	0	1	2	0	118
Residential: Household and gardening (mobile)	0	0	0	1	0	0	1	0	43
Agriculture/Forestry/Fishing: Off-road agriculture/forestry	0	3	10	7	2	3	17	0	584
Agriculture/Forestry/Fishing: National fishing	8	2	6	8	8	11	16	32	81
Other, Mobile	0	0	1	1	0	0	34	0	92
Road transport exhaust total	1	38	97	113	22	41	184	0	7 695
Road transport non exhaust total	6	6	81	39 937	0	95	5 291	27	18 141
Other mobile sources total	39	11	44	59	18	1 683	983	68	1 644
Domestic total	46	56	222	40 110	40	1 819	6 457	96	27 480
Civil aviation (International)	0	0	0	0	0	0	25	0	0
Navigation (International)	178	12	77	178	20	9 948	92	184	428

The heavy metal emission estimates for road transport are based on a national research study made by Winther and Slentø (2010). The latter study calculate the exhaust related emissions from fuel and engine oil as well as the wear related emissions from tyre, brake and road wear. Apart from Pb, the emission factors only deviate to a less extent due to changes in fleet and mileage composition over the years, which bring relative changes in fuel consumption per fuel type, engine oil use and aggregated emission factors for brake, tyre and road wear.

The most important exhaust related emissions for road transport are Cd, Cr, Hg and Zn. the most important wear related emissions are Cu and Pb almost solely coming from tyre wear, and Zn from brake and tyre wear. For other mobile sources, the most important emission contributions are calculated for Ni, Se and As, coming from the use of marine diesel oil in fisheries and navigation and residual oil in navigation.

The Figures 3.3.29 and 3.3.30 show the heavy metal emission distributions for all road transport sources split into vehicle categories, and for other mobile sectors, respectively.

For non-road mobile machinery in agriculture, forestry, industry, commercial/institutional and recreational, as well as military and railways, fuel related emission factors from road transport are used derived for the year 2009.



For civil aviation jet fuel no emissions are estimated due to lack of emission data, whereas for aviation gasoline fuel related emission factors for road transport gasoline is used derived for the year 2009, except for Pb where national data exist.

For navigation and fisheries, the heavy metal emission factors are fuel related, and are taken from the EMEP/EEA guidebook.




Figure 3.3.29 Heavy metal emission shares for road transport in 2013.




Figure 3.3.30 Heavy metal emission shares for other mobile sources in 2013.




Figure 3.3.31 Pb emissions (kg) pr vehicle type for road transport 1985-2013.




Figure 3.3.32 Pb emissions (kg) in NFR sectors for other mobile sources 1990-2013.

### Dioxin and PAH

In Table 3.3.5, the dioxin, PAH, HCB and PCB emissions for road transport and other mobile sources are shown for 2013 in NFR sectors. The emission figures in the time series 1990-2013 are given in Annex 2.B.16 (NFR format) and are shown for 1990 and 2013 in Annex 2.B.15 (CollectER format).

Table 3.3.5 Dioxin, PAH, HCB and PCB emissions in 2013 for road transport and other mobile sources.

	HCB	Dioxins/Furans	Flouranthene	Benzo(b) flouranthene	Benzo(k) flouranthene	Benzo(a) pyrene	Benzo- (g,h,i) perylene	Indeno (1,2,3-c,d) pyrene	PCB
	g	g	kg	kg	kg	kg	kg	kg	g
Manufacturing industries/Construction (mobile)	0,079	0,010	57	7	6	3	6	3	7
Civil aviation (Domestic)	0,000	0,000	0	0	0	0	0	0	0
Road transport: Passenger cars	0,265	0,048	669	50	38	44	88	44	1
Road transport: Light duty vehicles	0,130	0,012	209	15	11	13	26	12	0
Road transport: Heavy duty vehicles	0,244	0,042	83	21	24	4	3	5	21
Road transport: Mopeds & motorcycles	0,000	0,014	8	0	0	0	1	0	0
Road transport: Gasoline evaporation	0,000	0,000	0	0	0	0	0	0	0
Road transport: Brake wear	0,000	0,000	0	0	0	0	0	0	0
Road transport: Tyre wear	0,000	0,000	0	0	0	0	0	0	0
Road transport: Road abrasion	0,000	0,000	0	0	0	0	0	0	0
Railways	0,021	0,002	5	1	1	0	0	0	2
National navigation (Shipping)	0,014	0,066	33	2	1	0	4	4	0
Commercial/Institutional: Mobile	0,000	0,012	10	0	0	0	2	1	0
Residential: Household and gardening (mobile)	0,000	0,004	4	0	0	0	1	0	0
Agriculture/Forestry/Fishing: Off-road agriculture/forestry	0,092	0,013	67	8	7	4	7	4	8
Agriculture/Forestry/Fishing: National fishing	0,013	0,083	51	4	2	1	10	8	0
Other, Mobile	0,012	0,003	10	1	1	1	1	1	1
Road transport exhaust total	0,639	0,116	969	86	74	61	118	62	22
Road transport non exhaust total	0,000	0,000	0	0	0	0	0	0	0
Other mobile sources total	0,230	0,194	237	24	19	10	31	21	17
Domestic total	0,869	0,310	1206	110	93	71	149	82	39
Civil aviation (International)	0,000	0,000	0	0	0	0	0	0	0
Navigation (International)	0,069	0,315	138	10	5	3	19	16	0

For mobile sources, road transport displays the largest emission of dioxins and PAH. The dioxin emission share for road transport is 37 % of all mobile emissions in 2013, whereas Agriculture/forestry-/fisheries and Navigation have smaller shares of 31 and 21 %. For the different PAH components, road transport shares are around 80 % of total emissions for mobile sources. The remaining emissions almost solely come from Agriculture/forestry-/fisheries, Navigation and Industry with Agriculture/forestry-/fisheries as the largest source.

Figures 3.3.33 and 3.3.34 show the dioxin and PAH emission distributions into vehicle categories and other mobile sectors, respectively.




Figure 3.3.33 Dioxin, PAH, HCB and PCB emission shares for road transport in 2013.




Figure 3.3.34 Dioxin, PAH, HCB and PCB emission shares for other mobile sources in 2013.


### **Bunkers**

The most important emissions from bunker fuel consumption (fuel consumption for international transport) are SO<sub>2</sub> and NO<sub>x</sub>. The bunker emission totals are shown in Table 3.3.3 for 2013, split into sea transport and civil aviation. All emission figures in the 1985-2013 time series are given in Annex 2.B.16 (NFR format). In Annex 2.B.15, the emissions are also given in CollectER format for 2013.


The differences in emissions between navigation and civil aviation are much larger than the differences in fuel consumption and display a poor emission performance for international sea transport. In broad terms, the emission trends shown in Figure 3.3.35 are similar to the fuel consumption development.

However, for navigation minor differences occur for the emissions of SO<sub>2</sub> and NO<sub>x</sub> due to varying amounts of marine gas oil and residual oil, and for SO<sub>2</sub> and NO<sub>x</sub> the development in the emission factors also have an impact on the emission trends. For civil aviation, apart from the annual consumption of jet fuel, the development of the NO<sub>x</sub> emissions is also due to yearly variations in LTO/aircraft type (earlier than 2001) and city-pair statistics (2001 onwards).


### CO<sub>2</sub> emissions - international bunkers



### SO<sub>2</sub> emissions - international bunkers



### NO<sub>x</sub> emissions - international bunkers



### TSP emissions - international bunkers




Figure 3.3.35 CO<sub>2</sub>, SO<sub>2</sub>, NO<sub>x</sub> and TSP emissions for international transport 1985-2013.



### 3.3.2 Methodological issues

The description of methodologies and references for the transport part of the Danish inventory is given in two sections: one for road transport and one for the other mobile sources.

#### **Methodology and references for Road Transport**

For road transport, the detailed methodology is used to make annual estimates of the Danish emissions, as described in the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2013). The actual calculations are made with a model developed by ENVIS, using the European COPERT IV model methodology explained by EMEP/EEA (2013). In COPERT, fuel consumption and emission simulations can be made for operationally hot engines, taking into account gradually stricter emission standards and emission degradation due to catalyst wear. Furthermore, the emission effects of cold-start and evaporation are simulated.

#### **Vehicle fleet and mileage data**

Corresponding to the COPERT IV fleet classification, all present and future vehicles in the Danish fleet are grouped into vehicle classes, sub-classes and layers. The layer classification is a further division of vehicle sub-classes into groups of vehicles with the same average fuel consumption and emission behaviour, according to EU emission legislation levels. Table 3.3.6 gives an overview of the different model classes and sub-classes, and the layer level with implementation years are shown in Annex 2.B.1.

Fleet and annual mileage data are provided by DTU Transport for the vehicle categories present in COPERT IV (Jensen, 2014). DTU Transport use data from the Danish vehicle register kept by Statistics Denmark. The vehicle register data consist of vehicle type (passenger cars, vans, trucks, buses, mopeds, motorcycles), fuel type, vehicle weight, gross vehicle weight, engine size (passenger cars registered from 2005+), Euro class (trucks and buses registered from 1997+), NEDC type approval fuel efficiency value (passenger cars registered from 1997+) and vehicle first registration year.

In order to establish engine size data for passenger cars registered before 2005, a weight class-engine size transformation key is used examined by Cowi (2008) for new Danish cars from 1998. For the years before 1998, data for 1998 is used, and for the years 1999-2004 a linear interpolation between 1998 and 2005 weight class-engine size relations is used. For trucks, truck driver registration notes gathered by Statistics Denmark are used to split the fleet figures of ordinary trucks into number of solo trucks and truck-trailer combinations. Further the registration notes make it possible to assume the average total vehicle weight of the truck trailer combination. For articulated trucks also, the registration notes make it possible to assume the average total vehicle weight of the full articulated truck.

Danish mileage data comes from the Danish Road Directorate based on the Danish vehicle inspection program. Total mileage per year and vehicle category are derived for the years 1985-2013, together with a more detailed mileage matrix examined for the year 2008 (based on detailed vehicle inspection data analysis). The detailed mileage matrix contains annual mileage per vehicle subcategory for new vehicles and for every vintage back in time, which determines the yearly mileage reduction percentages as a function of vehicle age. In a first step, the detailed mileage matrix is combined with corresponding fleet numbers in order to estimate intermediate total mileages for each

year on a detailed fleet level. Next, each year's detailed (intermediate) mileage figures are scaled according to the difference between true and intermediate total mileage per vehicle subcategory.

DTU Transport (Jensen, 2014) also provides information of the mileage split between urban, rural and highway driving based on traffic monitoring data. The respective average speeds come from The Danish Road Directorate (e.g. Winther & Ekman, 1998). Additional data for the moped fleet and motorcycle fleet disaggregation is given by The National Motorcycle Association (Markamp, 2013).

Table 3.3.6 Model vehicle classes and sub-classes, trip speeds and mileage split.

Vehicle classes	Fuel type	Engine size/weight	Trip speed [km pr h]		
			Urban	Rural	Highway
PC	Gasoline	< 1.4 l.	40	70	100
PC	Gasoline	1.4 – 2 l.	40	70	100
PC	Gasoline	> 2 l.	40	70	100
PC	Diesel	< 2 l.	40	70	100
PC	Diesel	> 2 l.	40	70	100
PC	LPG		40	70	100
PC	2-stroke		40	70	100
LDV	Gasoline		40	65	80
LDV	Diesel		40	65	80
LDV	LPG		40	65	80
Trucks	Gasoline		35	60	80
Trucks	Diesel	Rigid 3,5 - 7,5t	35	60	80
Trucks	Diesel	Rigid 7,5 - 12t	35	60	80
Trucks	Diesel	Rigid 12 - 14 t	35	60	80
Trucks	Diesel	Rigid 14 - 20t	35	60	80
Trucks	Diesel	Rigid 20 - 26t	35	60	80
Trucks	Diesel	Rigid 26 - 28t	35	60	80
Trucks	Diesel	Rigid 28 - 32t	35	60	80
Trucks	Diesel	Rigid >32t	35	60	80
Trucks	Diesel	TT/AT 14 - 20t	35	60	80
Trucks	Diesel	TT/AT 20 - 28t	35	60	80
Trucks	Diesel	TT/AT 28 - 34t	35	60	80
Trucks	Diesel	TT/AT 34 - 40t	35	60	80
Trucks	Diesel	TT/AT 40 - 50t	35	60	80
Trucks	Diesel	TT/AT 50 - 60t	35	60	80
Trucks	Diesel	TT/AT >60t	35	60	80
Urban buses	Gasoline		30	50	70
Urban buses	Diesel	< 15 tonnes	30	50	70
Urban buses	Diesel	15-18 tonnes	30	50	70
Urban buses	Diesel	> 18 tonnes	30	50	70
Coaches	Gasoline		35	60	80
Coaches	Diesel	< 15 tonnes	35	60	80
Coaches	Diesel	15-18 tonnes	35	60	80
Coaches	Diesel	> 18 tonnes	35	60	80
Mopeds	Gasoline		30	30	-
Motorcycles	Gasoline	2 stroke	40	70	100
Motorcycles	Gasoline	< 250 cc.	40	70	100
Motorcycles	Gasoline	250 – 750 cc.	40	70	100
Motorcycles	Gasoline	> 750 cc.	40	70	100

In addition data from a survey made by the Danish Road Directorate (Hansen, 2010) has given information of the total mileage driven by foreign trucks on Danish roads in 2009. This mileage contribution has been added to the total mileage for Danish trucks on Danish roads, for trucks > 16 tonnes of gross vehicle weight. The data has been further processed by DTU Transport; by using appropriate assumptions the mileage have been back-casted to 1985 and forecasted to 2013.




Figure 3.3.36 Number of vehicles in sub-classes in 1985-2013.

For passenger cars, the engine size differentiation is less certain for the years before 2005. The increase in the total number of passenger cars is mostly due to a growth in the number of diesel cars between 1.4 and 2 litres (from the 2000's up to now). Until 2005, there has been a decrease in the number of gasoline cars with an engine size between 0.8 and 1.4 litres. These cars, however, have also increased in numbers during the later years, while the number of 1.4-2 litres gasoline cars has decreased. Since the late 1990's small cars (< 0.8 l gasoline and <1.4 l. diesel) has slowly begun to penetrate the fleet.

There has been a considerable growth in the number of diesel light-duty vehicles from 1985 to 2006; the number of vehicles has however decreased somewhat after 2006.

For the truck-trailer and articulated truck combinations there is a tendency towards the use of increasingly larger trucks throughout the time period. The decline in fleet numbers for many of the truck categories in 2007/2008 and until 2009 is caused by the impact of the global financial crisis and the reflagging of Danish commercial trucks to companies based in the neighbouring countries.

The number of urban buses has been almost constant between 1985 and 2013. The sudden change in the level of coach numbers from 1994 to 1995 is due to uncertain fleet data.

The reason for the significant growth in the number of mopeds from 1994 to 2002 is the introduction of the so-called Moped 45 vehicle type. From 2004 onwards there is a gradual switch from 2-stroke to 4-stroke in new sales for this vehicle category. For motorcycles, the number of vehicles has grown in general throughout the entire 1985-2013 period. The increase is, however, most visible from the mid-1990s and onwards.

The vehicle numbers are summed up in layers for each year (Figure 3.3.37) by using the correspondence between layers and first year of registration:

$$N_{j,y} = \sum_{i=FYear(j)}^{LYear(j)} N_{i,y} \quad (1)$$

Where N = number of vehicles, j = layer, y = year, i = first year of registration.

Weighted annual mileages pr layer are calculated as the sum of all mileage driven pr first registration year divided by the total number of vehicles in the specific layer.

$$M_{j,y} = \frac{\sum_{i=FYear(j)}^{LYear(j)} N_{i,y} \cdot M_{i,y}}{\sum_{i=FYear(j)}^{LYear(j)} N_{i,y}} \quad (2)$$

Since 2006 economical incitements have been give to private vehicle owners to buy Euro 5 diesel passenger cars and vans in order to bring down the particulate emissions from diesel vehicles. The estimated sales between 2006 and 2010 have been examined by the Danish EPA and are included in the fleet data behind the Danish inventory (Winther, 2011).

For heavy duty trucks, there is a slight deviation from the strict correspondence between EU emission layers and first registration year.

In this case, specific Euro class information for most of the vehicles from 2001 onwards is incorporated into the fleet and mileage data model developed by Jensen (2014). For inventory years before 2001, and for vehicles with no Euro information the normal correspondence between layers and first year of registration is used.

Vehicle numbers and weighted annual mileages pr layer are shown in Annex 2.B.1 and 3.B.2 for 1985-2013. The trends in vehicle numbers pr layer are also shown in Figure 3.3.37. The latter figure shows how vehicles complying with the gradually stricter EU emission levels (EURO I, II, III, IV, V etc.) have been introduced into the Danish motor fleet.




Figure 3.3.37 Layer distribution of vehicle numbers pr vehicle type in 1985-2013.

### Emission legislation

For Euro 1-4 passenger cars and light duty trucks, the chassis dynamometer test cycle used in the EU for emission approval is the NEDC (New European Driving Cycle), see Nørgaard and Hansen (2004). The test cycle is also used also for fuel consumption measurements. The NEDC cycle consists of two parts, the first part being a 4-time repetition (driving length: 4 km) of the ECE test cycle. The latter test cycle is the so-called urban driving cycle<sup>4</sup> (average speed: 19 km pr h). The second part of the test is the run-through of the EUDC (Extra Urban Driving Cycle) test driving segment, simulating the fuel consumption under rural and highway driving conditions. The driving length of EUDC is 7 km at an average speed of 63 km pr h. More information regarding the fuel measurement procedure can be found in the EU-directive [80/1268/EØF](#).

For NO<sub>x</sub>, VOC (NMVOC + CH<sub>4</sub>), CO and PM, the emissions from road transport vehicles have to comply with the different EU directives listed in Table 3.3.7. The emission directives distinguish between three vehicle classes according to vehicle reference mass<sup>5</sup>: Passenger cars and light duty trucks (<1305 kg), light duty trucks (1305-1760 kg) and light duty trucks (>1760 kg). The specific emission limits are shown in Annex 2.B.3.

In practice, the emissions from vehicles in traffic are different from the legislation limit values and, therefore, the latter figures are considered to be too inaccurate for total emission calculations. A major constraint is that the emission approval test conditions reflect only to a small degree the large variety of emission influencing factors in the real traffic situation, such as cumulated mileage driven, engine and exhaust after treatment maintenance levels and driving behaviour.

Therefore, in order to represent the Danish fleet and to support average national emission estimates, emission factors must be chosen which derive from numerous emissions measurements, using a broad range of real world driving patterns and a sufficient number of test vehicles. It is similar im-

<sup>4</sup> For Euro 3 and on, the emission approval test procedure was slightly changed. The 40 s engine warm up phase before start of the urban driving cycle was removed.

<sup>5</sup> Reference mass: net vehicle weight + mass of fuel and other liquids + 100 kg.

portant to have separate fuel consumption and emission data for cold-start emission calculations and gasoline evaporation (hydrocarbons).

For heavy-duty vehicles (trucks and buses), the emission limits are given in g pr kWh and the measurements are carried out for engines in a test bench, using the EU ESC (European Stationary Cycle) and ETC (European Transient Cycle) test cycles, depending on the Euro norm and exhaust gas after-treatment system installed. A description of the test cycles is given by Nørgaard and Hansen, 2004). Measurement results in g pr kWh from emission approval tests cannot be directly used for inventory work. Instead, emission factors used for national estimates must be transformed into g pr km, and derived from a sufficient number of measurements which represent the different vehicle size classes, Euro engine levels and real world variations in driving behaviour.

In terms of the sulphur content in the fuels used by road transportation vehicles, the EU directive 2003/17/EF describes the fuel quality standards agreed by the EU. In Denmark, the sulphur content in gasoline and diesel was reduced to 10 ppm in 2005, by means of a fuel tax reduction for fuels with 10 ppm sulphur contents.

Table 3.3.7 Overview of the existing EU emission directives for road transport vehicles.

Vehicle category	Emission layer	EU directive	First reg.
Passenger cars (gasoline)	PRE ECE	-	-
	ECE 15/00-01	70/220 - 74/290	1972 <sup>a</sup>
	ECE 15/02	77/102	1981 <sup>b</sup>
	ECE 15/03	78/665	1982 <sup>c</sup>
	ECE 15/04	83/351	1987 <sup>d</sup>
	Euro I	91/441	1.10.199
	Euro II	94/12	1.1.1997
	Euro III	98/69	1.1.2001
	Euro IV	98/69	1.1.2006
	Euro V	715/2007(692/2008)	1.1.2011
	Euro VI	715/2007(692/2008)	1.9.2015
Passenger cars (diesel and LPG)	Conventional	-	-
	ECE 15/04	83/351	1987 <sup>d</sup>
	Euro I	91/441	1.10.199
	Euro II	94/12	1.1.1997
	Euro III	98/69	1.1.2001
	Euro IV	98/69	1.1.2006
	Euro V	715/2007(692/2008)	1.1.2011
	Euro VI	715/2007(692/2008)	1.9.2015
Light duty trucks (gasoline and diesel)	Conventional	-	-
	ECE 15/00-01	70/220 - 74/290	1972 <sup>a</sup>
	ECE 15/02	77/102	1981 <sup>b</sup>
	ECE 15/03	78/665	1982 <sup>c</sup>
	ECE 15/04	83/351	1987 <sup>d</sup>
	Euro I	93/59	1.10.199
	Euro II	96/69	1.10.199
	Euro III	98/69	1.1.2002
	Euro IV	98/69	1.1.2007
	Euro V	715/2007(692/2008)	1.1.2011
	Euro VI	715/2007(692/2008)	1.9.2015
Heavy duty vehicles	Euro 0	88/77	1.10.199
	Euro I	91/542	1.10.199
	Euro II	91/542	1.10.199
	Euro III	1999/96	1.10.200
	Euro IV	1999/96	1.10.200
	Euro V	1999/96	1.10.200
	Euro VI	595/2009	1.10.201
Mopeds	Conventional	-	-
	Euro I	97/24	2000
	Euro II	2002/51	2004
	Euro III	2002/51	2014 <sup>f</sup>
	Euro IV	168/2013	2017
Motor cycles	Euro V	168/2013	2021
	Conventional	-	0
	Euro I	97/24	2000
	Euro II	2002/51	2004

Continued

Euro III	2002/51	2007
Euro IV	168/2013	2017
Euro V	168/2013	2021

a,b,c,d: Expert judgement suggest that Danish vehicles enter into the traffic before EU directive first registration dates. The effective inventory starting years are a: 1970; b: 1979; c: 1981; d: 1986.

e: The directive came into force in Denmark in 1991 (EU starting year: 1993).

### Fuel consumption and emission factors

The fuel consumption and emission factors used in the Danish inventory come from the COPERT IV (version 11) model. The source for these data is various European measurement programmes. In general the COPERT data are transformed into trip-speed dependent fuel consumption and emission factors for all vehicle categories and layers by using trip speeds as shown in Table 3.3.8. The factors are listed in Annex 2.B.4.

### Adjustment for fuel efficient vehicles

In the Danish fleet and mileage database kept by DTU Transport, the type approval fuel efficiency value based on the NEDC driving cycle ( $TA_{NEDC}$ ) is registered for each single car. Further, a modified fuel efficiency value ( $TA_{inuse}$ ) is calculated using  $TA_{NEDC}$ , vehicle weight and engine size as input parameters. The  $TA_{inuse}$  value better reflects the fuel consumption associated with the NEDC driving cycle under real (“inuse”) traffic conditions (Emisia, 2012).

From 2006 up to last historical year represented by fleet data, the average CO<sub>2</sub> emission factor (by fleet number) is calculated for each year’s new sold cars, based on the registered  $TA_{NEDC}$  values. Using the average CO<sub>2</sub> emission factor for the last historical year as starting point, the average emission factor for each year’s new sold cars are linearly reduced, until the emission factor reaches 95 g CO<sub>2</sub> /km in 2020.

From 2006 up to last historical year, the average CO<sub>2</sub> emission factor (by fleet number) is also calculated for each year’s new sold cars, and for each fuel type/engine size combination, based on  $TA_{NEDC}$  and  $TA_{inuse}$ .

The linear reduction of the average emission factor for each year’s new sold cars is then used to reduce the CO<sub>2</sub> emission factors for new sold cars based on  $TA_{inuse}$ , between last historical year and 2020, for each of the fuel type/engine size fleet segments.

Subsequently for each layer and inventory year, CO<sub>2</sub> emission factors are calculated based on  $TA_{inuse}$  and weighted by total mileage. On the same time corresponding layer specific CO<sub>2</sub> factors from COPERT IV are set up valid for Euro 4+ vehicles in the COPERT model. The COPERT IV CO<sub>2</sub> factors are derived from fuel consumption factors included in the COPERT IV model (EMEP/EEA, 2013) that represent the COPERT test vehicles under the NEDC driving cycle in real world traffic ( $TA_{COPERT IV, inuse}$ ).

In a final step the ratio between the layer specific CO<sub>2</sub> emission factors for the Danish fleet and the COPERT Euro IV vehicles under  $TA_{inuse}$  are used to scale the trip speed dependent fuel consumption factors provided by COPERT IV for Euro 4 layers onwards.



### **Adjustment for EGR, SCR and filter retrofits**

In COPERT IV emission factors are available for Euro V heavy duty vehicles using EGR and SCR exhaust emission aftertreatment systems, respectively. The estimated new sales of Euro V diesel trucks equipped with EGR and SCR during the 2006-2010 time periods has been examined by Hjelgaard and Winther (2011). These inventory fleet data are used in the Danish inventory to calculate weighted emission factors for Euro V trucks in different size categories.

During the 2000's urban environmental zones have been established in Danish cities in order to bring down the particulate emissions from diesel fuelled heavy duty vehicles. Driving in these environmental zones prescribe the use of diesel particulate filters. The Danish EPA has provided the estimated number of Euro I-III urban buses and Euro II-III trucks and tourist buses which have been retrofitted with filters during the 2000's. These retrofit data are included in the Danish inventory by assuming that particulate emissions are lowered by 80 % compared with the emissions from the same Euro technology with no filter installed (Winther, 2011).

For all vehicle categories/technology levels not represented by measurements, the emission factors are produced by using reduction factors. The latter factors are determined by assessing the EU emission limits and the relevant emission approval test conditions, for each vehicle type and Euro class.

### **Adjustment for biofuel usage**

A literature review carried out in the Danish research project REBECA revealed no significant changes in emission factors between neat gasoline and E5 gasoline-ethanol blends for the combustion related emission components; NO<sub>x</sub>, CO and VOC (Winther et al., 2012). Hence, due to the current low ethanol content in today's road transport gasoline, no modifications of the neat gasoline based COPERT emission factors are made in the inventories in order to account for ethanol usage.

REBECA results published by Winther (2009) have shown that the emission impact of using diesel-biodiesel blends is very small at low biodiesel blend ratios. Consequently no bio fuel emission factor adjustments are needed for diesel vehicles as well. However, adjustment of the emission factors for diesel vehicles will be made if the biodiesel content of road transport diesel fuel increases to a more significant level in the future.

### **Deterioration factors**

For three-way catalyst cars the emissions of NO<sub>x</sub>, NMVOC and CO gradually increase due to catalyst wear and are, therefore, modified as a function of total mileage by the so-called deterioration factors. Even though the emission curves may be serrated for the individual vehicles, on average, the emissions from catalyst cars stabilise after a given cut-off mileage is reached due to OBD (On Board Diagnostics) and the Danish inspection and maintenance programme.

For each year, the deterioration factors are calculated pr first registration year by using deterioration coefficients and cut-off mileages, as given in EMEP/EEA (2013), for the corresponding layer. The deterioration coefficients are given for the two driving cycles: "Urban Driving Cycle" (UDF) and "Extra Urban Driving Cycle" (EUDF: urban and rural), with trip speeds of 19 and 63 km pr h, respectively.

Firstly, the deterioration factors are calculated for the corresponding trip speeds of 19 and 63 km pr h in each case determined by the total cumulated mileage less than or exceeding the cut-off mileage. The Formulas 3 and 4 show the calculations for the "Urban Driving Cycle":

$$UDF = U_A \cdot MTC + U_B, MTC < U_{MAX} \quad (3)$$

$$UDF = U_A \cdot U_{MAX} + U_B, MTC \geq U_{MAX} \quad (4)$$

where UDF is the urban deterioration factor,  $U_A$  and  $U_B$  the urban deterioration coefficients, MTC = total cumulated mileage and  $U_{MAX}$  urban cut-off mileage.

In the case of trip speeds below 19 km pr h the deterioration factor, DF, equals UDF, whereas for trip speeds exceeding 63 km pr h, DF=EUDF. For trip speeds between 19 and 63 km pr h the deterioration factor, DF, is found as an interpolation between UDF and EUDF. Secondly, the deterioration factors, one for each of the three road types, are aggregated into layers by taking into account vehicle numbers and annual mileage levels pr first registration year:

$$DF_{j,y} = \frac{\sum_{i=FYear(j)}^{LYear(j)} DF_{i,y} \cdot N_{i,y} \cdot M_{i,y}}{\sum_{i=FYear(j)}^{LYear(j)} DF_{i,y} \cdot N_{i,y}} \quad (5)$$

where DF is the deterioration factor.

For  $N_2O$  and  $NH_3$ , COPERT IV takes into account deterioration as a linear function of mileage for gasoline fuelled EURO 1-4 passenger cars and light duty vehicles. The level of emission deterioration also relies on the content of sulphur in the fuel. The deterioration coefficients are given in EMEP/EEA (2013), for the corresponding layer. A cut-off mileage of 250.000 km is behind the calculation of the modified emission factors, and for the Danish situation the low sulphur level interval is assumed to be most representative.

#### Emissions and fuel consumption for hot engines

Emissions and fuel-use results for operationally hot engines are calculated for each year and for layer and road type. The procedure is to combine fuel consumption and emission factors (and deterioration factors for catalyst vehicles), number of vehicles, annual mileage levels and the relevant road-type shares given in Table 3.3.8. For non-catalyst vehicles this yields:

$$E_{j,k,y} = EF_{j,k,y} \cdot S_k \cdot N_{j,y} \cdot M_{j,y} \quad (6)$$

Here E = fuel consumption/emission, EF = fuel consumption/emission factor, S = road type share and k = road type.

For catalyst vehicles the calculation becomes:

$$E_{j,k,y} = DF_{j,k,y} \cdot EF_{j,k,y} \cdot S_k \cdot N_{j,y} \cdot M_{j,y} \quad (7)$$

### Extra emissions and fuel consumption for cold engines

Extra emissions of NO<sub>x</sub>, VOC, CH<sub>4</sub>, CO, PM, NH<sub>3</sub> and fuel consumption from cold start are simulated separately. For SO<sub>2</sub>, the extra emissions are derived from the cold start fuel consumption results.

Each trip is associated with a certain cold-start emission level and is assumed to take place under urban driving conditions. The number of trips is distributed evenly across the months. First, cold emission factors are calculated as the hot emission factor times the cold:hot emission ratio. Secondly, the extra emission factor during cold start is found by subtracting the hot emission factor from the cold emission factor. Finally, this extra factor is applied on the fraction of the total mileage driven with a cold engine (the  $\beta$ -factor) for all vehicles in the specific layer.

The cold:hot ratios depend on the average trip length and the monthly ambient temperature distribution. The Danish temperatures for 2013 are given in Cappelen et al. (2013). For previous years, temperature data are taken from similar reports available from The Danish Meteorological Institute ([www.dmi.dk](http://www.dmi.dk)). The cold:hot ratios are equivalent for gasoline fuelled conventional passenger cars and vans and for diesel passenger cars and vans, respectively, see EMEP/EEA (2013). For conventional gasoline and all diesel vehicles the extra emissions become:

$$CE_{j,y} = \beta \cdot N_{j,y} \cdot M_{j,y} \cdot EF_{U,j,y} \cdot (CEr - 1) \quad (8)$$

Where CE is the cold extra emissions,  $\beta$  = cold driven fraction, CEr = Cold:Hot ratio.

For catalyst cars, the cold:hot ratio is also trip speed dependent. The ratio is, however, unaffected by catalyst wear. The Euro I cold:hot ratio is used for all future catalyst technologies. However, in order to comply with gradually stricter emission standards, the catalyst light-off temperature must be reached in even shorter periods of time for future EURO standards. Correspondingly, the  $\beta$ -factor for gasoline vehicles is reduced step-wise for Euro II vehicles and their successors.

For catalyst vehicles the cold extra emissions are found from:

$$CE_{j,y} = \beta_{red} \cdot \beta_{EUROI} \cdot N_{j,y} \cdot M_{j,y} \cdot EF_{U,j,y} \cdot (CEr_{EUROI} - 1) \quad (9)$$

where  $\beta_{red}$  = the  $\beta$  reduction factor.

For CH<sub>4</sub>, specific emission factors for cold driven vehicles are included in COPERT IV. The  $\beta$  and  $\beta_{red}$  factors for VOC are used to calculate the cold driven fraction for each relevant vehicle layer. The NMVOC emissions during cold start are found as the difference between the calculated results for VOC and CH<sub>4</sub>.

For NH<sub>3</sub>, specific cold start emission factors are also proposed by COPERT IV. For catalyst vehicles, however, just like in the case of hot emission factors, the emission factors for cold start are functions of cumulated mileage (emission deterioration). The level of emission deterioration also relies on the content of sulphur in the fuel. The deterioration coefficients are given in EMEP/EEA (2013), for the corresponding layer. For cold start, the cut-off

mileage and sulphur level interval for hot engines are used, as described in the deterioration factors paragraph.

### Evaporative emissions from gasoline vehicles

For each year, evaporative emissions of hydrocarbons are simulated in the model as hot and warm running losses, hot and warm soak loss and diurnal emissions. The calculation approach is the same as in COPERT III. All emission types depend on RVP (Reid Vapour Pressure) and ambient temperature. The emission factors are shown in Ntziachristos et al. (2000).

Running loss emissions originate from vapour generated in the fuel tank while the vehicle is running. The distinction between hot and warm running loss emissions depends on engine temperature. In the model, hot and warm running losses occur for hot and cold engines, respectively. The emissions are calculated as annual mileage (broken down into cold and hot mileage totals using the  $\beta$ -factor) times the respective emission factors. For vehicles equipped with evaporation control (catalyst cars), the emission factors are only one tenth of the uncontrolled factors used for conventional gasoline vehicles.

$$R_{j,y} = N_{j,y} \cdot M_{j,y} \cdot ((1 - \beta) \cdot HR + \beta \cdot WR) \quad (10)$$

Where R is running loss emissions and HR and WR are the hot and warm running loss emission factors, respectively.

In the model, hot and warm soak emissions for carburettor vehicles also occur for hot and cold engines, respectively. These emissions are calculated as number of trips (broken down into cold and hot trip numbers using the  $\beta$ -factor) times respective emission factors:

$$S_{j,y}^c = N_{j,y} \cdot \frac{M_{j,y}}{l_{trip}} \cdot ((1 - \beta) \cdot HS + \beta \cdot WS) \quad (11)$$

Where  $S^c$  is the soak emission,  $l_{trip}$  = the average trip length, and HS and WS are the hot and warm soak emission factors, respectively. Since all catalyst vehicles are assumed to be carbon canister controlled, no soak emissions are estimated for this vehicle type. Average maximum and minimum temperatures per month are used in combination with diurnal emission factors to estimate the diurnal emissions from uncontrolled vehicles  $E^d(U)$ :

$$E_{j,y}^d(U) = 365 \cdot N_{j,y} \cdot e^d(U) \quad (12)$$

Each year's total is the sum of each layer's running loss, soak loss and diurnal emissions.

### Fuel consumption balance

The calculated fuel consumption in COPERT IV must equal the statistical fuel sale totals according to the UNFCCC and UNECE emissions reporting format. The statistical fuel sales for road transport are derived from the Danish Energy Authority data (see DEA, 2014). The DEA data are further processed for gasoline in order to account for e.g. non-road and recreational craft fuel consumption, which are not directly stated in the statistics, please refer to paragraph 1.1.4 for further information regarding the transformation of DEA fuel data.

The standard approach to achieve a fuel balance in annual emission inventories is to multiply the annual mileage with a fuel balance factor derived as the ratio between simulated and statistical fuel figures for gasoline and diesel, respectively. This method is also used in the present model.




Figure 3.3.38 DEA:DCE Fuel ratios (mileage adjustment factors) based on DEA fuel sales data and DCE fuel consumption estimates.




Figure 3.3.39 DEA:DCE Fuel ratios and (mileage adjustment factors) based on DEA fuel consumption data and DCE fuel consumption estimates.

In the figures 3.3.38 and 3.3.39 the COPERT IV:DEA gasoline and diesel fuel consumption ratios are shown for fuel sales and fuel consumption from 1985-2011. The data behind the figures are also listed in Annex 2.B.8. The fuel consumption figures are related to the traffic on Danish roads by Danish vehicles and foreign trucks.

Pr fuel type, all mileage numbers are equally scaled in order to obtain fuel equilibrium, and hence the mileage factors used are the reciprocal values of the COPERT IV:DEA fuel consumption: fuel sales ratio.

The reasons for the differences between DEA sales figures and bottom-up fuel estimates are mostly due to a combination of the uncertainties related to

COPERT IV fuel consumption factors, allocation of vehicle numbers in sub-categories, annual mileage, trip speeds and mileage splits for urban, rural and highway driving conditions.

The final fuel consumption and emission factors are shown in Annex 2.B.7 for 1985-2013. The total fuel consumption and emissions are shown in Annex 2.B.8, pr vehicle category and as grand totals, for 1985-2013 (and NFR format in Annex 2.B.16. In Annex 2.B.15, fuel consumption and emission factors as well as total emissions are given in CollectER format for 2013.

In Table 3.3.8, the aggregated emission factors for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP and BC are shown in CollectER format for Danish road transport.

Table 3.3.8 Fuel-based emission factors for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP and BC for road transport in Denmark (2013).

SNAP ID	Category	ModeFuel type		Emission factors <sup>1</sup> [g pr GJ]						
				SO <sub>2</sub>	NO <sub>x</sub>	NMVOC	CO	NH <sub>3</sub>	TSP	BC
070101	Passenger cars	Highway	Bio ethanol	0.00	114.38	27.44	641.76	29.29	1.00	0.15
070101	Passenger cars	Highway	Biodiesel	0.00	306.53	3.50	8.95	0.76	9.93	7.93
070101	Passenger cars	Highway	Diesel	0.47	306.53	3.50	8.95	0.76	9.93	7.93
070101	Passenger cars	Highway	Gasoline	0.46	114.38	27.44	641.76	29.29	1.00	0.15
070101	Passenger cars	Highway	LPG	0.00	163.05	25.53	1136.25	8.84	10.04	1.73
070102	Passenger cars	Rural	Bio ethanol	0.00	96.85	31.78	510.56	29.37	0.97	0.15
070102	Passenger cars	Rural	Biodiesel	0.00	279.98	4.89	20.89	0.81	8.30	6.53
070102	Passenger cars	Rural	Diesel	0.47	279.98	4.89	20.89	0.81	8.30	6.53
070102	Passenger cars	Rural	Gasoline	0.46	96.85	31.78	510.56	29.37	0.97	0.15
070102	Passenger cars	Rural	LPG	0.00	178.53	37.22	468.65	3.95	14.45	2.49
070103	Passenger cars	Urban	Bio ethanol	0.00	125.48	271.46	3049.34	6.96	0.84	0.13
070103	Passenger cars	Urban	Biodiesel	0.00	278.29	14.54	57.14	0.55	15.12	11.70
070103	Passenger cars	Urban	Diesel	0.47	278.29	14.54	57.14	0.55	15.12	11.70
070103	Passenger cars	Urban	Gasoline	0.46	125.48	271.46	3049.34	6.96	0.84	0.13
070103	Passenger cars	Urban	LPG	0.00	93.12	98.48	803.92	2.80	11.78	2.03
070201	Light duty vehicles	Highway	Bio ethanol	0.00	154.24	18.39	530.11	22.79	1.28	0.13
070201	Light duty vehicles	Highway	Biodiesel	0.00	289.05	14.62	93.24	0.37	15.85	12.75
070201	Light duty vehicles	Highway	Diesel	0.47	289.05	14.62	93.24	0.37	15.85	12.75
070201	Light duty vehicles	Highway	Gasoline	0.46	154.24	18.39	530.11	22.79	1.28	0.13
070201	Light duty vehicles	Highway	LPG	0.00	183.50	28.44	1175.04	0.00	10.04	1.46
070202	Light duty vehicles	Rural	Bio ethanol	0.00	135.07	26.97	401.29	21.88	1.13	0.12
070202	Light duty vehicles	Rural	Biodiesel	0.00	294.29	16.32	78.87	0.39	12.76	10.19
070202	Light duty vehicles	Rural	Diesel	0.47	294.29	16.32	78.87	0.39	12.76	10.19
070202	Light duty vehicles	Rural	Gasoline	0.46	135.07	26.97	401.29	21.88	1.13	0.12
070202	Light duty vehicles	Rural	LPG	0.00	199.99	44.15	465.82	0.00	14.45	2.10
070203	Light duty vehicles	Urban	Bio ethanol	0.00	124.24	191.86	3643.81	4.31	0.69	0.08
070203	Light duty vehicles	Urban	Biodiesel	0.00	271.04	34.45	108.63	0.26	22.06	17.67
070203	Light duty vehicles	Urban	Diesel	0.47	271.04	34.45	108.63	0.26	22.06	17.67
070203	Light duty vehicles	Urban	Gasoline	0.46	124.24	191.86	3643.81	4.31	0.69	0.08
070203	Light duty vehicles	Urban	LPG	0.00	102.60	96.19	611.21	0.00	11.86	1.72
070301	Heavy duty vehicles	Highway	Bio ethanol	0.00	1037.78	474.61	7610.35	0.28	55.35	2.77
070301	Heavy duty vehicles	Highway	Biodiesel	0.00	339.30	6.32	131.97	0.75	5.91	4.20
070301	Heavy duty vehicles	Highway	Diesel	0.47	339.30	6.32	131.97	0.75	5.91	4.20
070301	Heavy duty vehicles	Highway	Gasoline	0.46	1037.78	474.61	7610.35	0.28	55.35	2.77
070302	Heavy duty vehicles	Rural	Bio ethanol	0.00	1141.55	820.40	8371.39	0.30	60.88	3.04
070302	Heavy duty vehicles	Rural	Biodiesel	0.00	416.58	8.69	130.36	0.63	6.38	4.48
070302	Heavy duty vehicles	Rural	Diesel	0.47	416.58	8.69	130.36	0.63	6.38	4.48
070302	Heavy duty vehicles	Rural	Gasoline	0.46	1141.55	820.40	8371.39	0.30	60.88	3.04
070303	Heavy duty vehicles	Urban	Bio ethanol	0.00	456.62	696.09	7102.99	0.20	40.59	2.03
070303	Heavy duty vehicles	Urban	Biodiesel	0.00	546.32	13.01	145.90	0.38	7.65	5.27
070303	Heavy duty vehicles	Urban	Diesel	0.47	546.32	13.01	145.90	0.38	7.65	5.27
070303	Heavy duty vehicles	Urban	Gasoline	0.46	456.62	696.09	7102.99	0.20	40.59	2.03
070400	Mopeds	Urban	Bio ethanol	0.00	152.17	4161.90	6825.31	1.07	69.69	9.04
070400	Mopeds	Urban	Gasoline	0.46	152.17	4161.90	6825.31	1.07	69.69	9.04
070501	Motorcycles	Highway	Bio ethanol	0.00	271.13	626.52	10268.62	1.27	15.21	2.59
070501	Motorcycles	Highway	Gasoline	0.46	271.13	626.52	10268.62	1.27	15.21	2.59
070502	Motorcycles	Rural	Bio ethanol	0.00	192.44	628.85	9496.68	1.55	18.51	3.16
070502	Motorcycles	Rural	Gasoline	0.46	192.44	628.85	9496.68	1.55	18.51	3.16
070503	Motorcycles	Urban	Bio ethanol	0.00	118.69	776.38	9155.40	1.51	18.04	3.08
070503	Motorcycles	Urban	Gasoline	0.46	118.69	776.38	9155.40	1.51	18.04	3.08

<sup>1</sup> References. SO<sub>2</sub>: Country specific; NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, PM and BC: COPERT IV.

### **Non-exhaust particulate emissions from road transport**

The TSP, PM<sub>10</sub> and PM<sub>2.5</sub> emissions arising from tyre and brake wear (SNAP 0707) and road abrasion (SNAP 0708) are estimated for the years 2000-2013 as prescribed by the UNECE convention reporting format. The emissions are calculated by multiplying the total annual mileage per vehicle category with the correspondent average emission factors for each source type. The calculation procedure is consistent with the COPERT IV model approach used to estimate the Danish national emissions coming from exhaust. A more thorough explanation of the calculations is given by Winther and Slentø (2010). Emission factors are taken from EMEP/EEA (2013) and specific Danish tyre wear data are gathered by Winther and Slentø (2010). The emission factors and total emissions for 2013 are shown in Annex 2.B.15.

### **Methodologies and references for other mobile sources**

Other mobile sources are divided into several sub-sectors: sea transport, fishery, air traffic, railways, military, and working machinery and material in the industry, forestry, agriculture and household and gardening sectors. The emission calculations are made using the detailed method as described in the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2013) for air traffic, off-road working machinery and equipment, and ferries, while for the remaining sectors the simple method is used.

### **3.3.3 Activity data**

#### **Air traffic**

The activity data for air traffic consists of air traffic statistics provided by the Danish Transport Authority and Copenhagen Airport. Fuel statistics for jet fuel consumption and aviation gasoline are obtained from the Danish energy statistics (DEA, 2014).

For 2001 onwards, per flight records are provided by the Danish Transport Authority as data codes for aircraft type, and origin and destination airports (city-pairs).

Subsequently the aircraft types are separated by DCE into larger aircraft using jet fuel (jet engines, turbo props, helicopters) and small aircraft types with piston engines using aviation gasoline. This is done by using different aircraft dictionaries, internet look-ups and by communication with the Danish Transport Authority. Each of the larger aircraft type is then matched with a representative type for which fuel consumption and emission data are available from the EMEP/EEA databank. Relevant for this selection is aircraft maximum take off mass, engine types, and number of engines. A more thorough explanation is given in Winther (2001a, b).

Annex 2.B.10 shows the correspondence table between the actual aircraft type codes and representative aircraft types behind the Danish inventory. Annex 2.B.10 also show the number of LTO's per representative aircraft type for domestic and international flights starting from Copenhagen Airport and other airports, respectively<sup>6</sup>, in a time series from 2001-2013. The airport split is necessary to make due to the differences in LTO emission factors (c.f. section 3.3.4).

<sup>6</sup> Excluding flights for Greenland and the Faroe Islands. These flights are separately listed in Annex 2.B.10



The same type of LTO activity data for the flights for Greenland and the Faroe Islands are shown in Annex 2.B.10 also, further detailed into an origin-destination airport matrix and having flight distances attached. This level of detail satisfies the demand from UNFCCC to provide precise documentation for the part of the inventory for the Kingdom of Denmark being outside the Danish mainland.

The ideal flying distance (great circle distance) between the city-pairs is calculated by DCE in a separate database. The calculation algorithm uses a global latitude/altitude coordinate table for airports. In cases when airport coordinates are not present in the DCE database, these are looked up on the internet and entered into the database accordingly.

For inventory years prior to 2001, detailed LTO/aircraft type statistics are obtained from Copenhagen Airport (for this airport only), while information of total take-off numbers for other Danish airports is provided by the Danish Transport Authority. The assignment of representative aircraft types for Copenhagen Airport is done as described above. For the remaining Danish airports representative aircraft types are not directly assigned. Instead appropriate average assumptions are made relating to the fuel consumption and emission data part.




Figure 3.3.40 Most frequent domestic flying routes for large aircraft in Denmark.

Copenhagen Airport is the starting or end point for most of the domestic aviation made by large aircraft in Denmark (Figure 3.3.40; routes to Greenland/Faroe Islands are not shown). Even though many domestic flights not touching Copenhagen Airport are also reported in the flight statistics kept by the Danish Transport Authority, these flights, however, are predominantly made with small piston engine aircraft using aviation gasoline. Hence, the consumption of jet fuel by flights not using Copenhagen is merely marginal.




Figure 3.3.41 No. of LTO's for the most important airports in Denmark 2001-2013.

Figure 3.3.41 shows the number of domestic and international LTO's for Danish airports<sup>7</sup>, in a time series from 2001-2013.

### Non-road working machinery and equipment

Non-road working machinery and equipment are used in agriculture, forestry and industry, for household/gardening purposes and for sailing purposes (recreational craft). Information on the number of different types of machines, their respective load factors, engine sizes and annual working hours has been provided by Winther et al. (2006) for the years until 2004. For later inventory years, supplementary stock data are annually provided by the Association of Danish Agricultural Machinery Dealers and the Association of Producers and Distributors of Fork Lifts in Denmark. The stock development from 1985-2013 for the most important types of machinery are shown in Figures 3.3.42-3.3.49 below. The stock data are also listed in Annex 2.B.11, together with figures for load factors, engine sizes and annual working hours. As regards stock data for the remaining machinery types, please refer to (Winther et al., 2006).

<sup>7</sup> Flights for Greenland and the Faroe Islands are included under domestic in the figure.

It is important to note that key experts in the field of industrial non-road activities assumes a significant decrease in the activities for 2009 due to the global financial crisis. This reduction is in the order of 25 % for 2009 for industrial non-road in general (pers. comm. Per Stjernqvist, Volvo Construction Equipment 2010). For fork lifts 5 % and 20 % reductions are assumed for 2008 and 2009, respectively (pers. comm. Peter H. Møller, Rocla A/S).

For agriculture, the total number of agricultural tractors and harvesters per year are shown in the Figures 3.3.42-3.3.43, respectively. The figures clearly show a decrease in the number of small machines, these being replaced by machines in the large engine-size ranges.




Figure 3.3.42 Total numbers in kW classes for tractors from 1985 to 2013.





Figure 3.3.43 Total numbers in kW classes for harvesters from 1985 to 2013.

The tractor and harvester developments towards fewer vehicles and larger engines, shown in Figure 3.3.44, are very clear. From 1985 to 2013, tractor and harvester numbers decrease by around 39 % and 60 %, respectively, whereas the average increase in engine size for tractors is 45 %, and 202 % for harvesters, in the same time period.




Figure 3.3.44 Total numbers and average engine size for tractors and harvesters from 1985 to 2013.

The most important machinery types for industrial use are different types of construction machinery and fork lifts. The Figures 3.3.45 and 3.3.46 show the 1985-2013 stock development for specific types of construction machinery and diesel fork lifts. Due to lack of data, the construction machinery stock for 1990 is used also for 1985-1989. For most of the machinery types there is an increase in machinery numbers from 1990 onwards, due to increased construction activities. It is assumed that track type excavators/wheel type loaders (0-5 tonnes), and telescopic loaders first enter into use in 1991 and 1995, respectively.





Figure 3.3.45 1985-2013 stock development for specific types of construction machinery.




Figure 3.3.46 Total numbers of diesel fork lifts in kW classes from 1985 to 2013.

The emission level shares for tractors, harvesters, construction machinery and diesel fork lifts are shown in Figure 3.3.47, and present an overview of the penetration of the different pre-Euro engine classes, and engine stages complying with the gradually stricter EU stage I and II emission limits. The average lifetimes of 30, 25, 20 and 10 years for tractors, harvesters, fork lifts and construction machinery, respectively, influence the individual engine technology turn-over speeds.

The EU emission directive Stage I and II implementation years relate to engine size, and for all four machinery groups the emission level shares for the specific size segments will differ slightly from the picture shown in Figure

3.3.47. Due to scarce data for construction machinery, the emission level penetration rates are assumed to be linear and the general technology turnover pattern is as shown in Figure 3.3.47.




Figure 3.3.47 Emission level shares for tractors, harvesters, construction machinery and diesel fork lifts (1985 to 2013).

The 1985-2013 stock development for the most important household and gardening machinery types is shown in Figure 3.3.48.

For lawn mowers and cultivators, the machinery stock remains approximately the same for all years, whereas the stock figures for riders, chain saws, shrub clearers, trimmers and hedge cutters increase from 1990 onwards. The yearly stock increases, in most cases, become larger after 2000. The lifetimes for gasoline machinery are short and, therefore, there new emission levels (not shown) penetrate rapidly.




Figure 3.3.48 Stock developments 1985-2013 for the most important household and gardening machinery types.

Figure 3.3.49 shows the development in numbers of different recreational craft from 1985-2013. The 2004 stock data for recreational craft are repeated for 2005+, since no new fleet information has been obtained.

For diesel boats, increases in stock and engine size are expected during the whole period, except for the number of motor boats (< 27 ft.) and the engine sizes for sailing boats (<26 ft.), where the figures remain unchanged. A decrease in the total stock of sailing boats (<26 ft.) by 21 % and increases in the total stock of yawls/cabin boats and other boats (<20 ft.) by around 25 % are expected. Due to a lack of information specific to Denmark, the shifting rate from 2-stroke to 4-stroke gasoline engines is based on a German non-road study (IFEU, 2004).






Figure 3.3.49 1985-2013 Stock and engine size development for recreational craft.

### National sea transport

The methodology used to estimate the fuel consumption figures for national sea transport, based on fleet activity estimates for regional ferries, local ferries and other national sea transport is described by Winther (2008).

Table 3.3.11 lists the most important domestic ferry routes in Denmark in the period 1990-2013. For these ferry routes and the years 1990-2005, the following detailed traffic and technical data have been gathered by Winther (2008): Ferry name, year of service, engine size (MCR), engine type, fuel type, average load factor, auxiliary engine size and sailing time (single trip).

For 2006-2013, the above mentioned traffic and technical data for specific ferries have been provided by Kristensen (2014) in the case of Mols-Linien (Sjællands Odde-Ebeltoft, Sjællands Odde-Århus, Kalundborg-Århus), by Jørgensen (2014) for Færgen A/S (Køge-Rønne, Tårs-Spødsbjerg). For Esbjerg/Hanst-holm/Hirtshals-Torshavn traffic and technical data have been provided by Dávastovu (2010).

Table 3.3.9 Domestic ferry routes comprised in the Danish inventory.

Ferry service	Service period
Esbjerg-Torshavn	1990-1995, 2009+
Halsskov-Knudshoved	1990-1999
Hanstholm-Torshavn	1991-1992, 1999+
Hirtshals-Torshavn	2010
Hundested-Grenaa	1990-1996
Kalundborg-Juelsminde	1990-1996
Kalundborg-Samsø	1990+
Kalundborg-Århus	1990+
Korsør-Nyborg, DSB	1990-1997
Korsør-Nyborg, Vognmandsruten	1990-1999
København-Rønne	1990-2004
Køge-Rønne	2004+
Sjællands Odde-Ebeltoft	1990+
Sjællands Odde-Århus	1999+
Tårs-Spødsbjerg	1990+




Figure 3.3.50 Domestic regional ferry routes in Denmark (2013).

The number of round trips pr ferry route from 1990 to 2013 is provided by Statistics Denmark (2014), see Figure 3.3.51 (Esbjerg/Hanstholm/Hirtshals-Torshavn not shown). The traffic data are also listed in Annex 3.B.12, together with different ferry specific technical and operational data.

For each ferry, Annex 3.B.12 lists the relevant information as regards ferry route, name, year of service, engine size (MCR), engine type, fuel type, average load factor, auxiliary engine size and sailing time (single trip). There is a lack of historical traffic data for 1985-1989, and hence, data for 1990 is used for these years, to support the fuel consumption and emission calculations.




Figure 3.3.51 No. of round trips for the most important ferry routes in Denmark 1990-2013.

It is seen from Table 3.3.11 (and Figure 3.3.51) that several ferry routes were closed in the time period from 1996-1998, mainly due to the opening of the Great Belt Bridge (connecting Zealand and Funen) in 1997. Hundested-Grenaa and Kalundborg-Juelsminde was closed in 1996, Korsør-Nyborg (DSB) closed in 1997, and Halsskov-Knudshoved and Korsør-Nyborg (Vognmandsruten) was closed in 1998. The ferry line København-Rønne was replaced by Køge-Rønne in 2004 and from 1999 a new ferry connection was opened between Sjællands Odde and Århus.

For the local ferries, a bottom-up estimate of fuel consumption for 1996 has been taken from the Danish work in Wismann (2001). The latter project calculated fuel consumption and emissions for all sea transport in Danish waters in 1995/1996 and 1999/2000. In order to cover the entire 1990-2012 inventory period, the fuel figure for 1996 has been adjusted according to the developments in local ferry route traffic shown in Annex 3.B.12.

Fuel sold for freight transport by Royal Arctic Line between Aalborg (Denmark) and Greenland and by Eim Skip - East route between Aarhus (Denmark) and Torshavn (Faroe Islands) are included under other national sea transport in the Danish inventories. In both cases all fuel is being bought in Denmark (Rasmussen, 2014 and Thorarensen, 2014).

For the remaining part of the traffic between two Danish ports, other national sea transport, bottom-up estimates for fuel consumption have been calculated for the years 1995 and 1999 by Wismann (2007). These fuel consumption estimates are used as activity data for the inventory years until 1995 and 1999 onwards. Interpolated figures are used for the inventory years 1996-1998.

The calculations use the database set up for Denmark in the Wismann (2001) study, with actual traffic data from the Lloyd's LMIS database (not including ferries). The database was split into three vessel types: bulk carriers, container ships, and general cargo ships; and five size classes: 0-1000, 1000-3000, 3000-10000, 10000-20000 and >20000 DTW. The calculations assume that bulk carriers and container ships use heavy fuel oil, and that general cargo ships use gas oil. For further information regarding activity data for local ferries and other national sea transport, please refer to Winther (2008).

The fleet activity data for regional ferries and the fleet activity based fuel consumption estimates for local ferries and other national sea transport replace the fuel based activity data, which originated directly from the DEA statistics.

#### **Other sectors**

The activity data for military, railways, international sea transport and fishery consists of fuel consumption information from DEA (2014). For international sea transport, the basis is in principle fuel sold in Danish ports for vessels with a foreign destination, as prescribed by the IPCC guidelines.

However, it must be noted that fuel sold for sailing activities between Denmark and Greenland/Faroe Islands are reported as international in the DEA energy statistics. Hence, for inventory purposes in order to follow the IPCC guidelines the bottom-up fuel estimates for the ferry routes Esbjerg/Hanstholm/Hirshals-Torshavn, and fuel reports from Royal Arctic Line and Eim Skip is being subtracted from the fuel sales figures for international sea transport prior to inventory fuel input.

For fisheries, the calculation methodology described by Winther (2008) remains fuel based. However, the input fuel data differ from the fuel sales figures previously used. The changes are the result of further data processing of the DEA reported gas oil sales for national sea transport and fisheries, prior to inventory input. For years when the fleet activity estimates of fuel consumption for national sea transport are smaller than reported fuel sold, fuel is added to fisheries in the inventory. Conversely, lower fuel sales in relation to bottom-up estimates for national sea transport means that fuel is being subtracted from the original fisheries fuel sales figure in order to make up the final fuel consumption input for fisheries.

The updated fuel consumption time series for national sea transport lead, in turn, to changes in the energy statistics for fisheries (gas oil), industry (heavy fuel oil), and international sea transport, so the national energy balance can remain unchanged.

For all sectors, fuel-use figures are given in Annex 2.B.14 for 2013 in CollectER format.

### Emission legislation

For non-road working machinery and equipment, and recreational craft and railway locomotives/motor cars, the emission directives list specific emission limit values (g pr kWh) for CO, VOC, NO<sub>x</sub> (or VOC + NO<sub>x</sub>) and TSP, depending on engine size (kW for diesel, ccm for gasoline) and date of implementation (referring to engine market date).

For diesel, the directives 97/68 and 2004/26 (Table 3.3.10) relate to non-road machinery other than agricultural and forestry tractors and the directives have different implementation dates for machinery operating under transient and constant loads. The latter directive also comprises emission limits for railway machinery (Table 3.3.14). For tractors the relevant directives are 2000/25 and 2005/13 (Table 3.3.10). For gasoline, the directive 2002/88 distinguishes between hand-held (SH) and not hand-held (NS) types of machinery (Table 3.3.11).

For engine type approval, the emissions (and fuel consumption) are measured using various test cycles (ISO 8178). Each test cycle consists of a number of measurement points for specific engine loads during constant operation. The specific test cycle used depends on the machinery type in question and the test cycles are described in more details in the directives.

Table 3.3.10 Overview of EU emission directives relevant for diesel fuelled non-road machinery.

Stage/Engine size [kW]	CO	VOC	NO <sub>x</sub> VOC+NO <sub>x</sub>		PM	Diesel machinery			Tractors	
			[g pr kWh]			EU Directive	Implement. date	Implement. date	EU Directive	Implement. date
						Transient	Constant			
<b>Stage I</b>										
37<=P<75	6.5	1.3	9.2	-	0.85	97/68	1/4 1999	-	2000/25	1/7 2001
<b>Stage II</b>										
130<=P<560	3.5	1	6	-	0.2	97/68	1/1 2002	1/1 2007	2000/25	1/7 2002
75<=P<130	5	1	6	-	0.3		1/1 2003	1/1 2007		1/7 2003
37<=P<75	5	1.3	7	-	0.4		1/1 2004	1/1 2007		1/1 2004
18<=P<37	5.5	1.5	8	-	0.8		1/1 2001	1/1 2007		1/1 2002
<b>Stage IIIA</b>										
130<=P<560	3.5	-	-	4	0.2	2004/26	1/1 2006	1/1 2011	2005/13	1/1 2006
75<=P<130	5	-	-	4	0.3		1/1 2007	1/1 2011		1/1 2007
37<=P<75	5	-	-	4.7	0.4		1/1 2008	1/1 2012		1/1 2008
19<=P<37	5.5	-	-	7.5	0.6		1/1 2007	1/1 2011		1/1 2007
<b>Stage IIIB</b>										
130<=P<560	3.5	0.19	2	-	0.025	2004/26	1/1 2011	-	2005/13	1/1 2011
75<=P<130	5	0.19	3.3	-	0.025		1/1 2012	-		1/1 2012
56<=P<75	5	0.19	3.3	-	0.025		1/1 2012	-		1/1 2012
37<=P<56	5	-	-	4.7	0.025		1/1 2013	-		1/1 2013
<b>Stage IV</b>										
130<=P<560	3.5	0.19	0.4	-	0.025	2004/26	1/1 2014		2005/13	1/1 2014
56<=P<130	5	0.19	0.4	-	0.025		1/10 2014			1/10 2014

Table 3.3.11 Overview of the EU Emission Directive 2002/88 for gasoline fuelled non-road machinery.

	Category	Engine size [ccm]	CO [g pr kWh]	HC [g pr kWh]	NO <sub>x</sub> [g pr kWh]	HC+NO <sub>x</sub> [g pr kWh]	Implement. date
Stage I							
Hand held	SH1	S<20	805	295	5.36	-	1/2 2005
	SH2	20=<S<50	805	241	5.36	-	1/2 2005
	SH3	50=<S	603	161	5.36	-	1/2 2005
Not hand held	SN3	100=<S<225	519	-	-	16.1	1/2 2005
	SN4	225=<S	519	-	-	13.4	1/2 2005
Stage II							
Hand held	SH1	S<20	805	-	-	50	1/2 2008
	SH2	20=<S<50	805	-	-	50	1/2 2008
	SH3	50=<S	603	-	-	72	1/2 2009
Not hand held	SN1	S<66	610	-	-	50	1/2 2005
	SN2	66=<S<100	610	-	-	40	1/2 2005
	SN3	100=<S<225	610	-	-	16.1	1/2 2008
	SN4	225=<S	610	-	-	12.1	1/2 2007

For recreational craft, Directive 2003/44 comprises the emission legislation limits for diesel engines, and for 2-stroke and 4-stroke gasoline engines, respectively. The CO and VOC emission limits depend on engine size (kW) and the inserted parameters presented in the calculation formulas in Table 3.3.12. For NO<sub>x</sub>, a constant limit value is given for each of the three engine types. For TSP, the constant emission limit regards diesel engines only.

In Table 3.3.13 the Stage II emission limits are shown for recreational craft. CO and HC+NO<sub>x</sub> limits are provided for gasoline engines depending on the rated engine power and the engine type (stern-drive vs. outboard) while CO, HC+NO<sub>x</sub>, and particulate emission limits are defined for Compression Ignition (CI) engines depending on the rated engine power and the swept volume.

Table 3.3.12 Overview of the EU Emission Directive 2003/44 for recreational craft.

Engine type	Impl. date	CO=A+B/P <sup>n</sup>			HC=A+B/P <sup>n</sup>			NO <sub>x</sub>	TSP
		A	B	n	A	B	n		
2-stroke gasoline	1/1 2007	150.0	600.0	1.0	30.0	100.0	0.75	10.0	-
4-stroke gasoline	1/1 2006	150.0	600.0	1.0	6.0	50.0	0.75	15.0	-
Diesel	1/1 2006	5.0	0.0	0	1.5	2.0	0.5	9.8	1.0

Table 3.3.13 Overview of the EU Emission Directive 2013/53 for recreational craft.

Diesel engines					
Swept Volume, SV l/cyl.	Rated Engine Power, P <sub>N</sub> kW	Impl. Date	CO g/kWh	HC + NO <sub>x</sub> g/kWh	PM g/kWh
SV < 0.9	P <sub>N</sub> < 37				
	37 ≤ P <sub>N</sub> < 75 (*)	18/1 2017	5	4.7	0.30
	75 ≤ P <sub>N</sub> < 3 700	18/1 2017	5	5.8	0.15
0.9 ≤ SV < 1.2	P <sub>N</sub> < 3 700	18/1 2017	5	5.8	0.14
1.2 ≤ SV < 2.5		18/1 2017	5	5.8	0.12
2.5 ≤ SV < 3.5		18/1 2017	5	5.8	0.12
3.5 ≤ SV < 7.0		18/1 2017	5	5.8	0.11
Gasoline engines					
Engine type	Rated Engine Power, P <sub>N</sub> kW		CO g/kWh	HC + NO <sub>x</sub> g/kWh	PM g/kWh
Stern-drive and inboard engines	P <sub>N</sub> ≤ 373	18/1 2017	75	5	-
	373 ≤ P <sub>N</sub> ≤ 485	18/1 2017	350	16	-
	P <sub>N</sub> > 485	18/1 2017	350	22	-
Outboard engines and PWC engines (**)	P <sub>N</sub> ≤ 4.3	18/1 2017	500 – (5.0 x P <sub>N</sub> )	15.7 + (50/PN <sup>0.9</sup> )	-
	4.3 ≤ P <sub>N</sub> ≤ 40	18/1 2017	500 – (5.0 x P <sub>N</sub> )	15.7 + (50/PN <sup>0.9</sup> )	-
	P <sub>N</sub> > 40	18/1 2017	300		-

(\*) Alternatively, this engine segment shall not exceed a PM limit of 0.2 g/kWh and a combined HC + NO<sub>x</sub> limit of 5.8 g/kWh.

(\*\*) Small and medium size manufacturers making outboard engines ≤ 15 kW have until 18/1 2020 to comply.

Table 3.3.14 Overview of the EU Emission Directive 2004/26 for railway locomotives and motorcars.

Engine size [kW]			CO [g pr kWh]	HC [g pr kWh]	NO <sub>x</sub> [g pr kWh]	HC+NO <sub>x</sub> [g pr kWh]	PM [g pr kWh]	Implement. date
Locomotives Stage IIIA								
130 ≤ P < 560	RL A		3.5	-	-	4	0.2	1/1 2007
560 < P	RH A		3.5	0.5	6	-	0.2	1/1 2009
2000 ≤ P and piston displacement ≥ 5 l/cyl.	RH A		3.5	0.4	7.4	-	0.2	1/1 2009
	Stage IIIB	RB	3.5	-	-	4	0.025	1/1 2012
Motor cars Stage IIIA								
130 < P	RC A		3.5	-	-	4	0.2	1/1 2006
	Stage IIIB							
130 < P	RC B		3.5	0.19	2	-	0.025	1/1 2012

Aircraft engine emissions of NO<sub>x</sub>, CO, VOC and smoke are regulated by ICAO (International Civil Aviation Organization). The engine emission certification standards are contained in Annex 16 — Environmental Protection, Volume II — Aircraft Engine Emissions to the Convention on International Civil Aviation (ICAO Annex 16, 2008, plus amendments). The emission standards relate to the total emissions (in grams) from the so-called LTO (Landing and Take Off) cycle divided by the rated engine thrust (kN). The ICAO LTO cycle contains the idealised aircraft movements below 3000 ft (915 m) during approach, landing, airport taxiing, take off and climb out.

For smoke all aircraft engines manufactured from 1 January 1983 have to meet the emission limits agreed by ICAO. For NO<sub>x</sub>, CO, VOC The emission legislation is relevant for aircraft engines with a rated engine thrust larger

than 26.7 kN. In the case of CO and VOC, the ICAO regulations apply for engines manufactured from 1 January 1983.

For  $\text{NO}_x$ , the emission regulations fall in five categories

- 1) For engines of a type or model for which the date of manufacture of the first individual production model was before 1 January 1996, and for which the production date of the individual engine was before 1 January 2000.
- 2) For engines of a type or model for which the date of manufacture of the first individual production model is on or after 1 January 1996, or for individual engines with a production date on or after 1 January 2000.
- 3) For engines of a type or model for which the date of manufacture of the first individual production model is on or after 1 January 2004.
- 4) For engines of a type or model for which the date of manufacture of the first individual production model is on or after 1 January 2008, or for individual engines with a production date on or after 1 January 2013.
- 5) For engines of a type or model for which the date of manufacture of the first individual production model is on or after 1 January 2014.

The regulations published by ICAO are given in the form of the total quantity of pollutants ( $D_p$ ) emitted in the LTO cycle divided by the maximum sea level thrust ( $F_{oo}$ ) and plotted against engine pressure ratio at maximum sea level thrust.

The limit values for  $\text{NO}_x$  are given by the formulae in Table 3.3.15.



Table 3.3.15 Current certification limits for NO<sub>x</sub> for turbo jet and turbo fan engines.

	Engines first produced before 1.1.1996 & for engines manufactured before 1.1.2000	Engines first produced on or after 1.1.1996 & for engines manufactured on or after 1.1.2000	Engines for which the date of manufacture of the first individual production model was on or after 1 January 2004	Engines first produced on or after 1.1.2008 & for engines manufactured on or after 1.1.2013	Engines for which the date of manufacture of the first individual production model was on or after 1.1.2014
Applies to engines >26.7 kN	$D_p/F_{oo} = 40 + 2\pi_{oo}$	$D_p/F_{oo} = 32 + 1.6\pi_{oo}$			
Engines of pressure ratio less than 30					
Thrust more than 89 kN			$D_p/F_{oo} = 19 + 1.6\pi_{oo}$	$D_p/F_{oo} = 16.72 + 1.4080\pi_{oo}$	$7.88 + 1.4080\pi_{oo}$
Thrust between 26.7 kN and not more than 89 kN			$D_p/F_{oo} = 37.572 + 1.6\pi_{oo} - 0.208F_{oo}$	$D_p/F_{oo} = 38.54862 + (1.6823\pi_{oo}) - (0.2453F_{oo}) - (0.00308\pi_{oo}F_{oo})$	$D_p/F_{oo} = 40.052 - 1.5681\pi_{oo} - 0.3615F_{oo} - 0.0018\pi_{oo} \times F_{oo}$
Engines of pressure ratio more than 30 and less than 62.5 (104.7)					
Thrust more than 89 kN			$D_p/F_{oo} = 7 + 2.0\pi_{oo}$	$D_p/F_{oo} = -1.04 + (2.0 \times \pi_{oo})$	
Thrust between 26.7 kN and not more than 89 kN			$D_p/F_{oo} = 42.71 + 1.4286\pi_{oo} - 0.4013F_{oo} + 0.00642\pi_{oo}F_{oo}$	$D_p/F_{oo} = 46.1600 + (1.4286\pi_{oo}) - (0.5303F_{oo}) - (0.00642\pi_{oo}F_{oo})$	
Engines with pressure ratio 62.5 or more					
Engines with pressure ratio 82.6 or more			$D_p/F_{oo} = 32 + 1.6\pi_{oo}$	$D_p/F_{oo} = 32 + 1.6\pi_{oo}$	
Engines of pressure ratio more than 30 and less than (104.7)					
Thrust more than 89 kN					$D_p/F_{oo} = -9.88 + 2.0\pi_{oo}$
Thrust between 26.7 kN and not more than 89 kN					$D_p/F_{oo} = 41.9435 + 1.505\pi_{oo} - 0.5823F_{oo} + 0.005562\pi_{oo} \times F_{oo}$
Engines with pressure ratio 104.7 or more					
					$D_p/F_{oo} = 32 + 1.6\pi_{oo}$

Source: International Standards and Recommended Practices, Environmental Protection, ICAO Annex 16 Volume II , 3rd edition July 2008, plus amendments: Amendment 7 (17 November 2011), Amendment 8 (July 2014)

where:

$D_p$  = the sum of emissions in the LTO cycle in g.

$F_{oo}$  = thrust at sea level take-off (100 %).

$\pi_{oo}$  = pressure ratio at sea level take-off thrust point (100 %).

The equivalent limits for HC and CO are  $D_p/F_{oo} = 19.6$  for HC and  $D_p/F_{oo} = 118$  for CO (ICAO Annex 16 Vol. II paragraph 2.2.2). Smoke is limited to a regulatory smoke number =  $83 (F_{oo})^{-0.274}$  or a value of 50, whichever is the lower.

A further description of the technical definitions in relation to engine certification as well as actual engine exhaust emission measurement data can be found in the ICAO Engine Exhaust Emission Database. The latter database is accessible from <http://www.caa.co.uk>, hosted by the UK Civil Aviation Authority.

Marpol 73/78 Annex VI agreed by IMO (International Maritime Organisation) concerns the control of NO<sub>x</sub> emissions (Regulation 13 plus amendments) and SO<sub>x</sub> and particulate emissions (Regulation 14 plus amendments) from ships (DNV, 2009).

The baseline NO<sub>x</sub> emission regulation of Annex VI apply for diesel engines with a power output higher than 130 kW, which are installed on a ship constructed on or after 1 January 2000 and diesel engines with a power output higher than 130 kW which undergo major conversion on or after 1 January 2000.

The baseline NO<sub>x</sub> emission limits for ship engines in relation to their rated engine speed (n) given in RPM (Revolutions Pr Minute) are the following:

- 17 g pr kWh, n < 130 RPM
- 45 x n-0.2 g pr kWh, 130 ≤ n < 2000 RPM
- 9.8 g pr kWh, n ≥ 2000 RPM

The further amendment of Annex VI Regulation 13 contains a three tiered approach in order to strengthen the emission standards for NO<sub>x</sub>. The three tier approach comprises the following:

- Tier I: Diesel engines (> 130 kW) installed on a ship constructed on or after 1 January 2000 and prior to 1 January 2011 (initial regulation).
- Tier II: Diesel engines (> 130 kW) installed on a ship constructed on or after 1 January 2011.
- Tier III<sup>8</sup>: Diesel engines (> 130 kW) installed on a ship constructed on or after 1 January 2016.

The three tier NO<sub>x</sub> emission limit functions are shown in Table 3.3.16.

Table 3.3.16 Tier I-III NO<sub>x</sub> emission limits for ship engines in MARPOL Annex VI.

	NO <sub>x</sub> limit	RPM (n)
Tier I	17 g pr kWh	n < 130
	45 · n-0.2 g pr kWh	130 ≤ n < 2000
	9,8 g pr kWh	n ≥ 2000
Tier II	14.4 g pr kWh	n < 130
	44 · n-0.23 g pr kWh	130 ≤ n < 2000
	7.7 g pr kWh	n ≥ 2000
Tier III	3.4 g pr kWh	n < 130
	9 · n-0.2 g pr kWh	130 ≤ n < 2000
	2 g pr kWh	n ≥ 2000

Further, the NO<sub>x</sub> Tier I limits are to be applied for existing engines with a power output higher than 5000 kW and a displacement per cylinder at or above 90 litres, installed on a ship constructed on or after 1 January 1990 but prior to 1 January 2000.

In relation to the sulphur content in heavy fuel and marine gas oil used by ship engines, Table 3.3.17 shows the EU and IMO (Regulation 14 plus amendments) legislation in force for SECA (Sulfur Emission Control Area) areas and outside SECA's.

<sup>8</sup> For ships operating in a designated Emission Control Area. Outside a designated Emission Control Area, Tier II limits apply.

Table 3.3.17 Current legislation in relation to marine fuel quality.

Legislation	Heavy fuel oil		Gas oil	
	S- %	Implement. date (day/month/year)	S- %	Implement. date (day/month/year)
EU-directive 93/12	None		0.2 <sup>1</sup>	01.10.1994
EU-directive 1999/32	None		0.2	01.01.2000
EU-directive 2005/33 <sup>2</sup>	SECA - Baltic sea	1.5	11.08.2006	01.01.2008
	SECA - North sea	1.5	11.08.2007	01.01.2008
	Outside SECA's	None		0.1
MARPOL Annex VI	SECA – Baltic sea	1.5	19.05.2006	
	SECA – North sea	1.5	21.11.2007	
	Outside SECA	4.5	19.05.2006	
MARPOL Annex VI amendments	SECA's	1	01.03.2010	
	SECA's	0.1	01.01.2015	
	Outside SECA's	3.5	01.01.2012	
	Outside SECA's	0.5	01.01.2020 <sup>3</sup>	

<sup>1</sup> Sulphur content limit for fuel sold inside EU.

<sup>2</sup> From 1.1.2010 fuel with a sulphur content higher than 0.1 % must not be used in EU ports for ships at berth exceeding two hours.

<sup>3</sup> Subject to a feasibility review to be completed no later than 2018. If the conclusion of such a review becomes negative the effective date would default 1 January 2025.

For non-road machinery, the EU directive 2003/17/EC gives a limit value of 10 ppm sulphur in diesel (from 2011).

#### Emission factors

The SO<sub>2</sub> emission factors are fuel related, and rely on the sulphur contents given in the relevant EU fuel directives or in the Danish legal announcements. However, for jet fuel the default factor from IPCC (1996) is used. Road transport diesel is assumed to be used by engines in military and railways, and road transport gasoline is assumed to be used by non-road working machinery and recreational craft. Hence, these types of machinery have the same SO<sub>2</sub> emission factors, as for road transport.

For all mobile sources, the emission factor source for BC, NH<sub>3</sub>, heavy metals and PAH is the EMEP/EEA guidebook (EMEP/EEA, 2013). The heavy metal emission factors for road transport and other mobile sources except national sea transport and fisheries originate from Winther and Slentø (2010). For civil aviation jet fuel, no heavy metal emission factors are proposed due to lack of data.

In the case of military ground equipment, aggregated emission factors for gasoline and diesel are derived from road traffic emission simulations. For piston engine aircraft using aviation gasoline, aggregated emission factors for conventional cars are used.

For railways, specific Danish measurements from the Danish State Railways (DSB) (Delvig, 2014) are used to calculate the emission factors of NO<sub>x</sub>, VOC, CO and TSP, and a NMVOC/CH<sub>4</sub> split is made based on expert judgment.

For agriculture, forestry, industry, household gardening and recreational craft, the NO<sub>x</sub>, VOC, CO and TSP emission factors are derived from various European measurement programmes; see IFEU (2004, 1999) and Winther et al. (2006). The NMVOC/CH<sub>4</sub> split is taken from IFEU (1999).

For national sea transport and fisheries, the NO<sub>x</sub> emission factors predominantly come from the engine manufacturer MAN Diesel, as a function of engine production year. The CO, VOC and TSP emission factors come from the Danish TEMA2010 emission model (Trafikministeriet, 2010), whereas the PM<sub>10</sub> and PM<sub>2.5</sub> size fractions are obtained from MAN Diesel.

Specifically for the ferries used by Mols Linjen new NO<sub>x</sub>, VOC and CO emission factors are provided by Kristensen (2008), originating from measurement results by Hansen et al. (2004), Wismann (1999) and PHP (1996). Kristensen (2013) has provided complimentary emission factor data for new ferries.

For ship engines VOC/CH<sub>4</sub> splits are taken from EMEP/EEA (2013), and all emission factors are shown in Annex 2.B.13.

The source for aviation (jet fuel) emission factors is the EMEP/EEA guidebook (EMEP/EEA, 2013). For a number of different representative aircraft types, the EMEP/EEA guidebook comprises fuel flow and NO<sub>x</sub>, CO and VOC emission indices for the four LTO modes and distance based emission factors for cruise. VOC/CH<sub>4</sub> splits for aviation are taken from EMEP/EEA (2013).

For all sectors, emission factors are given in CollectER format in Annex 2.B.15 for 2013. Table 3.3.19 shows the emission factors for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP and BC in CollectER format used to calculate the emissions from other mobile sources in Denmark.

#### **Factors for deterioration, transient loads and gasoline evaporation for non-road machinery**

The emission effects of engine wear are taken into account for diesel and gasoline engines by using the so-called deterioration factors. For diesel engines alone, transient factors are used in the calculations, to account for the emission changes caused by varying engine loads. The evaporative emissions of NMVOC are estimated for gasoline fuelling and tank evaporation. The factors for deterioration, transient loads and gasoline evaporation are taken from IFEU (2004, 1999, 2014), and are shown in Annex 2.B.10. For more details regarding the use of these factors, please refer to paragraph 3.1.4 or Winther et al. (2006).

Table 3.3.18 Fuel based emission factors for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP and BC for other mobile sources in Denmark (2013).

SNAP ID	Category	Fuel type	Emission factors <sup>1</sup> [g pr GJ]						
			SO <sub>2</sub>	NO <sub>x</sub>	NMVOC	CO	NH <sub>3</sub>	TSP	BC
080100	Military	AvGas	22.99	859.00	1242.60	6972.00	1.60	10.00	1.50
080100	Military	Diesel	0.44	332.41	10.40	80.77	0.62	10.12	7.79
080100	Military	Gasoline	0.44	112.69	150.85	1434.44	22.16	1.48	0.21
080100	Military	Jet fuel	22.99	250.57	24.94	229.89	0.00	1.16	0.56
080200	Railways	Diesel	0.47	705.00	49.11	111.00	0.20	21.00	13.65
080300	Recreational craft	Diesel	46.84	764.00	138.41	392.19	0.17	85.15	31.51
080300	Recreational craft	Gasoline	0.46	577.88	597.66	9513.05	0.11	10.93	0.55
080402	National sea traffic	Diesel	46.84	1276.29	43.54	148.56	0.00	21.55	6.68
080402	National sea traffic	Residual oil	489.00	1910.03	63.36	209.02	0.00	43.98	5.28
080403	Fishing	Diesel	46.84	1323.46	57.86	190.89	0.00	21.55	6.68
080404	International sea traffic	Diesel	46.84	1588.52	58.07	191.57	0.00	21.55	6.68
080404	International sea traffic	Residual oil	489.00	2120.24	63.89	210.76	0.00	43.98	5.28
080501	Air traffic, Dom. < 3000 ft.	AvGas	22.83	859.00	1242.60	6972.00	1.60	10.00	1.50
080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22.99	330.14	9.51	107.31	0.00	1.16	0.56
080502	Air traffic, Int. < 3000 ft.	AvGas	22.83	859.00	1242.60	6972.00	1.60	10.00	1.50
080502	Air traffic, Int. < 3000 ft.	Jet fuel	22.99	352.63	9.77	105.59	0.00	1.16	0.56
080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22.99	356.58	7.07	102.73	0.00	1.16	0.56
080504	Air traffic, Int. > 3000 ft.	Jet fuel	22.99	312.14	6.47	76.02	0.00	1.16	0.56
080600	Agriculture	Diesel	0.47	484.78	48.03	297.49	0.18	37.96	23.72
080600	Agriculture	Gasoline	0.46	109.75	1172.11	21868.55	1.49	30.50	1.52
080700	Forestry	Diesel	0.47	327.36	27.43	213.51	0.18	23.10	17.25
080700	Forestry	Gasoline	0.46	54.79	3754.36	17915.98	0.09	82.19	4.11
080800	Industry	Diesel	0.47	476.00	53.75	298.69	0.18	47.06	33.66
080800	Industry	Gasoline	0.46	213.59	1571.80	14228.73	0.10	22.10	1.10
080800	Industry	LPG	0.00	1328.11	146.09	104.85	0.21	4.89	0.24
080900	Household and gardening	Gasoline	0.46	108.21	2256.56	31048.01	0.09	17.36	0.87
081100	Commercial and institutional	Gasoline	0.46	93.28	1533.43	30913.57	0.09	28.53	1.43
080501	Air traffic, Dom. < 3000 ft.	AvGas	22.83	859.00	1242.60	6972.00	1.60	10.00	0.00
080501	Air traffic, Dom. < 3000 ft.	Jet fuel	22.99	327.09	12.49	141.70	0.00	1.16	0.56
080502	Air traffic, Int. < 3000 ft.	AvGas	22.83	859.00	1242.60	6972.00	1.60	10.00	0.00
080502	Air traffic, Int. < 3000 ft.	Jet fuel	22.99	359.17	10.56	130.33	0.00	1.16	0.56
080503	Air traffic, Dom. > 3000 ft.	Jet fuel	22.99	370.16	4.98	51.13	0.00	1.16	0.56
080504	Air traffic, Int. > 3000 ft.	Jet fuel	22.99	376.37	3.71	43.02	0.00	1.16	0.56

<sup>1</sup> References. SO<sub>2</sub>: Country-specific; Military: Aggregated emission factors for road transport; Railways (NO<sub>x</sub>, CO, NMVOC and TSP): Danish State Railways; Agriculture, forestry, industry, household gardening and inland waterways (NO<sub>x</sub>, CO, VOC and TSP): IFEU (2004, 1999, 2014); National sea transport and fishing: MAN B&W (NO<sub>x</sub>) and TE-MA2000 (CO, NMVOC, TSP); Aviation - jet fuel (NO<sub>x</sub>, CO, NMVOC): EMEP/EEA; Aviation - av.gasoline: Aggregated emission factors for conventional gasoline cars.

### 3.3.4 Calculation method

#### Air traffic

For aviation, the domestic and international estimates are made separately for landing and take-off (LTOs < 3000 ft), and cruising (> 3000 ft).

By using the LTO mode specific fuel flow and emission indices from EMEP/EEA (2013), the fuel consumption and emission factors for the full LTO cycle can be estimated for each of the representative aircraft types used in the Danish inventory.

The fuel consumption for one LTO cycle is calculated according to the following sum formula:

$$FC_{LTO}^a = \sum_{m=1}^4 t_m \cdot ff_{a,m} \quad (13)$$

Where FC = fuel consumption (kg), m = LTO mode (approach/landing, taxiing, take off, climb out), t = times in mode (s), ff = fuel flow (kg pr s), a = representative aircraft type.

The emissions for one LTO cycle are estimated as follows:

$$E_{LTO}^a = \sum_{m=1}^4 FC_{a,m} \cdot EI_{a,m} \quad (14)$$

Due to lack of specific airport data, for approach/descent, take off and climb out, standardised times-in-modes of 4, 0.7 and 2.2 mins are used as defined by ICAO (ICAO, 1995), whereas for taxiing the appropriate time interval is 13 mins in Copenhagen Airport and 5 mins in other airports present in the Danish inventory.

For each representative aircraft type, the calculated fuel consumption and emission factors per LTO are shown in Annex 2.B.10 for Copenhagen Airport and other airports.

The calculations for cruise use the distance specific fuel consumption and emissions given by EMEP/EEA (2013) per representative aircraft type. Data interpolations or extrapolations are made – in each case determined by the great circle distance between the origin and the destination airports.

If the great circle distance, y, is smaller than the maximum distance for which fuel consumption and emission data are given in the EMEP/EEA data bank the fuel consumption or emission E (y) becomes:

$$E(y) = E_{x_i} + \frac{(y - x_i)}{x_{i+1} - x_i} \cdot (E_{x_{i+1}} - E_{x_i}) \quad y < x_{\max}, i = 0, 1, 2, \dots, \max-1 \quad (15)$$

In (15)  $x_i$  and  $x_{\max}$  denominate the separate distances and the maximum distance, respectively, with known fuel consumption and emissions. If the flight distance y exceeds  $x_{\max}$  the maximum figures for fuel consumption and emissions must be extrapolated and the equation then becomes:

$$E(y) = E_{x_{\max}} + \frac{(y - x_{\max})}{x_{\max} - x_{\max-1}} \cdot (E_{x_{\max}} - E_{x_{\max-1}}) \quad y > x_{\max} \quad (16)$$

Total results are summed up and categorised according to each flight's destination airport code in order to distinguish between domestic and international flights.

Annex 2.B.10 shows the average fuel consumption and emission factors per representative aircraft type for cruise flying, as well as total distance flown, for 2013<sup>9</sup>. The factors are split between Copenhagen Airport and other airports and distinguish between domestic and international flights.

<sup>9</sup> Excluding flights for Greenland and the Faroe Islands.

Specifically for flights between Denmark and Greenland or the Faroe Islands, for each representative aircraft type, the flight distances are directly shown in Annex 2.B.10, which go into the cruise calculation expressions 15 and 16.

The overall fuel precision in the model is 0.92 in 2013, derived as the fuel ratio between model estimates and statistical sales. The fuel difference is accounted for by adjusting cruising fuel consumption and emissions in the model according to domestic and international cruising fuel shares.

For inventory years before 2001, the calculation procedure is to estimate each year's fuel consumption and emissions for LTO based on LTO/aircraft type statistics from Copenhagen Airport, and total take off numbers for other airports provided by the Danish Transport Authority. Due to lack of aircraft type specific LTO data, fuel consumption and emission factors derived for domestic LTO's in Copenhagen Airport is used for all LTO's in other airports. In a next step, the total fuel consumption for cruise (true cruise fuel consumption) is found year by year as the statistical fuel consumption total minus the calculated fuel consumption for LTO.

For each inventory year, intermediate cruise fuel consumption figures split into four parts (Copenhagen/Other airports; domestic/international) are found as proportional values between part specific LTO fuel consumption values estimated as described previously, and part specific cruise:LTO fuel consumption ratios for 1998. In the latter ratio, the cruise fuel contribution is taken from a Danish city-pair emission inventory in 1998 (Winther, 2001a), whereas the LTO fuel consumption is the 1998 figures estimated as explained above.

Each inventory year's true cruise fuel consumption is finally split into four parts by using the intermediate cruise fuel consumption values as a distribution key. As emission factor input data for cruise, aggregated fuel related emission factors are derived from the present model and the first historical year 2001.

#### **Non-road working machinery and recreational craft**

Prior to adjustments for deterioration effects and transient engine operations, the fuel consumption and emissions in year X, for a given machinery type, engine size and engine age, are calculated as:

$$E_{Basis}(X)_{i,j,k} = N_{i,j,k} \cdot HRS_{i,j,k} \cdot P \cdot LF_i \cdot EF_{y,z} \quad (17)$$

where  $E_{Basis}$  = fuel consumption/emissions in the basic situation, N = number of engines, HRS = annual working hours, P = average rated engine size in kW, LF = load factor, EF = fuel consumption/emission factor in g pr kWh, i = machinery type, j = engine size, k = engine age, y = engine-size class and z = emission level. The basic fuel consumption and emission factors are shown in Annex 2.B.11.

The deterioration factor for a given machinery type, engine size and engine age in year X depends on the engine-size class (only for gasoline), y, and the emission level, z. The deterioration factors for diesel and gasoline 2-stroke engines are found from:

$$DF_{i,j,k}(X) = \frac{K_{i,j,k}}{LT_i} \cdot DF_{y,z} \quad (18)$$

where DF = deterioration factor, K = engine age, LT = lifetime, i = machinery type, j = engine size, k = engine age, y = engine-size class and z = emission level.

For gasoline 4-stroke engines the deterioration factors are calculated as:

$$DF_{i,j,k}(X) = \sqrt{\frac{K_{i,j,k}}{LT_i}} \cdot DF_{y,z} \quad (19)$$

The deterioration factors inserted in (18) and (19) are shown in Annex 2.B.11. No deterioration is assumed for fuel consumption (all fuel types) or for LPG engine emissions and, hence, DF = 1 in these situations.

The transient factor for any given machinery type, engine size and engine age in year X, relies only on emission level and load factor, and is denominated as:

$$TF_{i,j,k}(X) = TF_z \quad (20)$$

Where i = machinery type, j = engine size, k = engine age and z = emission level.

The transient factors inserted in (20) are shown in Annex 2.B.11. No transient corrections are made for gasoline and LPG engines and, hence, TF<sub>z</sub> = 1 for these fuel types.

The final calculation of fuel consumption and emissions in year X for a given machinery type, engine size and engine age, is the product of the expressions 17-20:

$$E(X)_{i,j,k} = E_{Basis}(X)_{i,j,k} \cdot TF(X)_{i,j,k} \cdot (1 + DF(X)_{i,j,k}) \quad (21)$$

The evaporative hydrocarbon emissions from fuelling are calculated as:

$$E_{Evap, fueling, i} = FC_i \cdot EF_{Evap, fueling} \quad (22)$$

Where E<sub>Evap, fueling, i</sub> = hydrocarbon emissions from fuelling, i = machinery type, FC = fuel consumption in kg, EF<sub>Evap, fueling</sub> = emission factor in g NMVOC pr kg fuel.

For tank evaporation, the hydrocarbon emissions are found from:

$$E_{Evap, tank, i} = N_i \cdot EF_{Evap, tank, i} \quad (23)$$

Where E<sub>Evap, tank, i</sub> = hydrocarbon emissions from tank evaporation, N = number of engines, i = machinery type and EF<sub>Evap, fueling</sub> = emission factor in g NMVOC pr year.

#### **Ferries, other national sea transport and fisheries**

The fuel consumption and emissions in year X, for regional ferries are calculated as:



$$E(X) = \sum_i N_i \cdot T_i \cdot S_{i,j} \cdot P_i \cdot LF_j \cdot EF_{k,l,y} \quad (24)$$

Where E = fuel consumption/emissions, N = number of round trips, T = sailing time pr round trip in hours, S = ferry share of ferry service round trips, P = engine size in kW, LF = engine load factor, EF = fuel consumption/emission factor in g pr kWh, i = ferry service, j = ferry, k = fuel type, l = engine type, y = engine year.

For the remaining navigation categories, the emissions are calculated using a simplified approach:

$$E(X) = \sum_i EC_{i,k} EF_{k,l,y} \quad (25)$$

Where E = fuel consumption/emissions, EC = energy consumption, EF = fuel consumption/emission factor in g pr kg fuel, i = category (local ferries, other national sea, fishery, international sea), k = fuel type, l = engine type, y = average engine year.

The emission factor inserted in (25) is found as an average of the emission factors representing the engine ages which are comprised by the average lifetime in a given calculation year, X:

$$EF_{k,l,y} = \frac{\sum_{year=X-LT}^{year=X} EF_{k,l}}{LT_{k,l}} \quad (26)$$

#### Other sectors

For military and railways, the emissions are estimated with the simple method using fuel-related emission factors and fuel consumption from the DEA:

$$E = FC \cdot EF \quad (27)$$

where E = emission, FC = fuel consumption and EF = emission factor. The calculated emissions for other mobile sources are shown in CollectER format in Annex 2.B.15 for the years 2013 and as time series 1985-2013 in Annex 2.B.16 (NFR format).

#### Energy balance between DEA statistics and inventory estimates

Following convention rules, the DEA statistical fuel sales figures are the basis for the full Danish inventory. However, in some cases for mobile sources the DEA statistical sectors do not fully match the inventory sectors. This is the case for non-road machinery, where relevant DEA statistical sectors also include fuel consumed by stationary sources.

In other situations, fuel consumption figures estimated by DCE from specific bottom-up calculations are regarded as more reliable than DEA reported sales. This is the case for national sea transport.

In the following the transferral of fuel consumption data from DEA statistics into inventory relevant categories is explained for national sea transport and fisheries, non-road machinery and recreational craft, and road transport. A full list of all fuel consumption data, DEA figures as well as intermediate fuel consumption data, and final inventory input figures is shown in Annex 2.B.14.

### **National sea transport and fisheries**

For national sea transport in Denmark, the fuel consumption estimates obtained by DCE (see 3.3.3 Activity data – national sea transport) are regarded as much more accurate than the DEA fuel sales data, since the large fluctuations in reported fuel sales cannot be explained by the actual development in the traffic between different national ports. As a consequence, the DCE bottom-up estimates are used in the Danish inventory for national sea transport.

There are different potential reasons for the differences between estimated fuel consumption and reported sales for national sea transport in Denmark. According to the DEA, the latter fuel differences are most likely explained by inaccurate costumer specifications made by the oil suppliers. This inaccuracy can be caused by a sector misallocation in the sales statistics between national sea transport and fisheries for gas oil, and between national sea transport and industry for heavy fuel oil (Peter Dal, DEA, personal communication, 2007). Further, fuel sold for vessels sailing between Denmark and Greenland/Faroe Islands are reported as international in the DEA statistics, and this fuel categorisation is different from the IPCC guideline definitions (see following paragraph “Bunkers”).

Following this, for fisheries and industry the updated fuel consumption time series for national sea transport lead, in turn, to changes in the fuel activity data for fisheries (gas oil), industry (heavy fuel oil) and international sea transport, so the national energy balance can remain unchanged.

For fisheries, fuel investigations made prior to the initiation of the work made by Winther (2008) have actually pointed out a certain area of inaccuracy in the DEA statistics. No engines installed in fishing vessels use heavy fuel oil, even though a certain amount of heavy fuel oil is listed in the DEA numbers for some statistical years (H. Amdissen, Danish Fishermen's Association, personal communication, 2006). Hence, for fisheries small amounts of fuel oil are transferred to national sea transport, and in addition small amounts of gasoline and diesel are transferred to recreational craft.

### **Non-road machinery and recreational craft**

For diesel and LPG, the non-road fuel consumption estimated by DCE is partly covered by the fuel-use amounts in the following DEA sectors: agriculture and forestry, market gardening, and building and construction. The remaining quantity of non-road diesel and LPG is taken from the DEA industry sector.

For gasoline, the DEA residential sector, together with the DEA sectors mentioned for diesel and LPG, contribute to the non-road fuel consumption total. In addition, a certain amount of fuel from road transport is needed to reach the fuel-use goal.

The amount of diesel and LPG in DEA industry not being used by non-road machinery is included in the sectors, “Combustion in manufacturing industry” (0301) and “Non-industrial combustion plants” (0203) in the Danish emission inventory.

For recreational craft, the calculated fuel-use totals for diesel and gasoline are subsequently subtracted from the DEA fishery sector. For gasoline, the DEA reported fuel consumption for fisheries is far too small to fill the fuel

gap, and hence the missing fuel amount is taken from the DEA road transport sector.

### **Bunkers**

The distinction between domestic and international emissions from aviation and navigation should be in accordance with the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. For the national emission inventory this, in principle, means that fuel sold (and associated emissions) for flights/sea transportation starting from a seaport/airport in the Kingdom of Denmark, with destinations inside or outside the Kingdom of Denmark, are regarded as domestic or international, respectively.

### **Aviation**

As prescribed by the IPCC guidelines, for aviation, the fuel consumption and emissions associated with flights inside the Kingdom of Denmark are counted as domestic.

This report includes flights from airports in Denmark and associated jet fuel sales. Hence, the flights between airports in Denmark and flights from Denmark to Greenland and the Faroe Islands are classified as domestic and flights from Danish airports with destinations outside the Kingdom of Denmark are classified as international flights.

In Greenland and in the Faroe Islands, the jet fuel sold is treated as domestic. This decision becomes reasonable when considering that almost no fuel is bunkered in Greenland/the Faroe Islands by flights other than those going to Denmark.

### **Navigation**

In DEA statistics, the domestic fuel total consists of fuel sold to Danish ferries and other ships sailing between two Danish ports. The DEA international fuel total consists of the fuel sold in Denmark to international ferries, international warships, other ships with foreign destinations, transport to Greenland and the Faroe Islands, tank vessels and foreign fishing boats.

In order to follow the IPCC guidelines the bottom-up fuel estimates for the ferry routes between Denmark and the Faroe Islands, and freight transport between Denmark and Greenland/Faroe Islands are being subtracted from the fuel sales figures for international sea transport prior to inventory fuel input.

In Greenland, all marine fuel sales are treated as domestic. In the Faroe Islands, fuel sold in Faroese ports for Faroese fishing vessels and other Faroese ships is treated as domestic. The fuel sold to Faroese ships bunkering outside Faroese waters and the fuel sold to foreign ships in Faroese ports or outside Faroese waters is classified as international (Lastein and Winther, 2003).

Conclusively, the domestic/international fuel split (and associated emissions) for navigation is not determined with the same precision as for aviation. It is considered, however, that the potential of incorrectly allocated fuel quantities is only a small part of the total fuel sold for navigational purposes in the Kingdom of Denmark.

### 3.3.5 Uncertainties and time series consistency

Emission uncertainty estimates are made for road transport and other mobile sources using the guidelines for estimating uncertainties in the EMP/EEA guidebook (EMEP/EEA, 2013). However, for TSP the latter source indicates no uncertainty factor and, instead, this factor is based on expert judgement.

The activity data uncertainty factor is assumed to be 2 and 10 % for road transport and other mobile sources, respectively, based on expert judgement.

The uncertainty estimates should be regarded as preliminary only and may be subject to changes in future inventory documentation. The calculations are shown in Annex 2.B.17 for all emission components.

Table 3.3.19 Uncertainties for activity data, emission factors and total emissions in 2013 and as a trend.

Pollutant	Emission factor uncertainties [ %]		Emission uncertainties [ %]	
	Road	Other	Overall 2013	Trend
SO <sub>2</sub>	50	50	49	3
NO <sub>x</sub>	50	100	55	9
NMVOG	50	100	56	8
CO	50	100	63	17
NH <sub>3</sub>	1000	1000	994	1679
TSP	50	100	47	5
PM <sub>10</sub>	50	100	50	4
PM <sub>2.5</sub>	50	100	54	3
BC	50	100	56	4
Arsenic	1000	1000	873	62
Cadmium	1000	1000	824	160
Chromium	1000	1000	826	200
Copper	1000	1000	999	5
Mercury	1000	1000	713	106
Nickel	1000	1000	928	34
Lead	1000	1000	859	12
Selenium	1000	1000	767	137
Zinc	1000	1000	942	39
Dioxins	1000	1000	729	153
Flouranthene	1000	1000	827	84
Benzo(b) flouranthene	1000	1000	811	180
Benzo(k) flouranthene	1000	1000	818	269
Benzo(a) pyrene	1000	1000	873	288
Benzo(g,h,i) perylene	1000	1000	818	64
indeno(1,2,3-c,d) pyrene	1000	1000	789	164
HCB	1000	1000	782	235
PCB	1000	1000	713	130

As regards time series consistency, background flight data cannot be made available on a city-pair level from 2000 or earlier. However, aided by LTO/aircraft statistics for these years and the use of proper assumptions, a good level of consistency is still obtained in this part of the transport inventory.

The time series of emissions for mobile machinery in the agriculture, forestry, industry, household and gardening (residential), and inland waterways (part of navigation) sectors are less certain than time series for other sectors, since DEA statistical figures do not explicitly provide fuel consumption information for working equipment and machinery.

### **3.3.6 Quality assurance/quality control (QA/QC)**

It is the intention to publish every second year a sector report for road transport and other mobile sources. The last sector report concerned the 2010 inventory (Winther, 2012).

The QA/QC descriptions of the Danish emission inventories for transport are given in Nielsen et al. (2014).

### **3.3.7 Recalculations**

The following recalculations and improvements of the emission inventories have been made since the emission reporting in 2013.

#### **Road transport**

Based on the updated version of COPERT IV version 11 launched in 2014, fuel consumption and NO<sub>x</sub>, VOC, CO and PM emission factors for euro 5 and 6 gasoline and diesel passenger cars and light duty vehicles have been updated in the model. Updated Euro V and VI fuel consumption and NO<sub>x</sub>, VOC, CO and PM emission factors for heavy duty vehicles have been included in the calculations also.

For N<sub>2</sub>O and NH<sub>3</sub>, Euro 5/6 and V/VI emission factors are also updated for passenger cars/light duty vehicles (only N<sub>2</sub>O) and heavy duty vehicles.

Further a new Euro 6c technology class has been added for diesel passenger cars and light duty vehicles.

The amount of diesel sold for road transport reported by the Danish Energy Agency has been slightly changed in 2012.

Very small changes in mileage data has been made for the years 1985-2012 based on new information from DTU Transport.

The percentage emission change interval and year of largest percentage differences (low %; high %, year) for the different emission components are: SO<sub>2</sub> (0 %; 0.2 %, 2011), NO<sub>x</sub> (-1.4 %; 0 %, 2012), NMVOC (-0.5 %; 0 %, 2012), NH<sub>3</sub> (-2 %; 0.6 %, 2012) and TSP (-2.3 %; 0.1 %, 2012).

#### **Navigation**

Recreational craft have been regrouped in the Danish inventory. These vessels have now been removed from the navigation sector and included under Other (1A5b), the latter sector according to its sector subtitle also comprise recreational craft. Further, small amounts of LPG and kerosene previously included in the navigation sector has now been transferred to stationary sources.

The following largest percentage differences (in brackets) for domestic navigation are noted for: SO<sub>2</sub> (-5.6 %), NO<sub>x</sub> (-12.3 %), NMVOC (-80 %), NH<sub>3</sub> (-100 %) and TSP (-42 %).

#### **Agriculture/forestry**

The baseline emission factors of NO<sub>x</sub>, TSP, CO and VOC and the transient factors for fuel consumption, NO<sub>x</sub>, TSP, CO and VOC for diesel machinery has been slightly changed in the calculation model used to estimate the emissions from Danish non road mobile machinery. Further, the CH<sub>4</sub> fraction of VOC has been updated. The changes are made based on new emission knowledge published by IFEU (1999) for baseline emission factors and CH<sub>4</sub> fractions, and IFEU (2014) for transient factors.

The number of agricultural tractors has been regrouped into finer engine size intervals for all inventory years. The total number of agricultural tractors and harvesters has been updated for the years 2006-2012 based on new stock data from Statistics Denmark for the year 2013.

The following largest percentage differences (in brackets) for agriculture/forestry are noted for: SO<sub>2</sub> (-16.2 %), NO<sub>x</sub> (-17.7 %), NMVOC (-5.3 %), NH<sub>3</sub> (-13.4 %) and TSP (13.6 %).

#### **Fisheries**

Small amounts of LPG and kerosene previously included in the navigation sector has now been transferred to stationary sources.

The following largest percentage differences (in brackets) for fisheries are noted for: SO<sub>2</sub> (0.1 %), NO<sub>x</sub> (-0.8 %), NMVOC (-5.1 %), NH<sub>3</sub> (0.0 %) and TSP (-0.1 %).

#### **Industry**

The baseline emission factors of NO<sub>x</sub>, TSP, CO and VOC and the transient factors for fuel consumption, NO<sub>x</sub>, TSP, CO and VOC for diesel machinery has been slightly changed in the calculation model used to estimate the emissions from Danish non road mobile machinery. Further, the CH<sub>4</sub> fraction of VOC has been updated. The changes are made based on new emission knowledge published by IFEU (1999) for baseline emission factors and CH<sub>4</sub> fractions, and IFEU (2014) for transient factors.

The following largest percentage differences (in brackets) for industrial non road machinery are noted for: SO<sub>2</sub> (0.7 %), NO<sub>x</sub> (1.3 %), NMVOC (4.3 %), NH<sub>3</sub> (1.8 %) and TSP (5.7 %).

#### **Civil aviation**

The model used for calculating civil aviation emissions has been updated by replacing the previous fuel consumption and emission factors for representative aircraft types (46 types) with a new and more comprehensive list of aircraft types (79 types) provided by Eurocontrol and published in the EMEP/EEA guidebook (EMEP/EEA, 2014).

The following largest percentage differences (in brackets) for civil aviation are noted for: SO<sub>2</sub> (38 %), NO<sub>x</sub> (35 %), NMVOC (-39 %), and TSP (30 %).

### **Other (Military and recreational craft)**

Recreational craft have been regrouped in the Danish inventory. These vessels have now been removed from the navigation sector and included under Other (1A5b), the latter sector according to its sector subtitle also comprises recreational craft.

NO<sub>x</sub>, CO, VOC and TSP emission factors have been updated for diesel and gasoline boats complying with Directive 2003/44 based on reassessment of the emission factor data reported by IFEU (2004). The latter reference has all the time been the source of emission data for recreational craft in the Danish emission inventory.

The following largest percentage differences (in brackets) for military are noted for: SO<sub>2</sub> (651 %), NO<sub>x</sub> (218 %), NMVOC (2800 %), NH<sub>3</sub> (149 %) and TSP (1047 %).

### **3.3.8 Improvements**

Fuel consumption and emission factors for road transport vehicles will be updated by the time when new data becomes available from COPERT model updates. Emission factors for TSP and BC will be updated for the representative aircraft types used in the aviation emission calculation model, based on aircraft engine specific smoke numbers and VOC emission indices, and jet fuel sulphur content.

### **3.3.9 References for transport and other mobile sources**

Cappelen, J. 2014: The Climate of Denmark 2013 - with English summary, Technical report No 14-01, pp. 79, Danish Meteorological Institute.

Cowi, 2008: Model til beregning af vej og banetransportens CO<sub>2</sub>-æquivalent emissioner, Technical note, pp. 42, Ministry of Transport (in Danish).

Dávastovu, J. 2010: Unpublished data material from Smyril Line.

Danish Energy Agency, 2014: The Danish energy statistics, Available at: <http://www.ens.dk/sites/ens.dk/files/info/tal-kort/statistik-noegletal/aarlig-energistatistik/energistatistik2013.pdf> (15-01-2015).

Delvig, P. 2014: Unpublished data material from the Danish State Railways.

DNV (2009): Marpol 73/78 Annex VI Regulations for the Prevention of Air Pollution from Ships - Technical and Operational implications, 32 pp.

Winther, M.; Ekman, B. (1998): Emissioner fra vejtrafikken i Danmark 1980-2010. Danmarks Miljøundersøgelser, Aarhus Universitet, 1998. 73 s. (Faglig rapport fra DMU, Vol. 256).

EMEP/EEA, 2013: Air Pollutant Emission Inventory Guidebook, prepared by the UNECE/EMEP Task Force on Emissions Inventories and Projections (TFEIP). Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (15-01-2015).

Hansen, C.O. 2010: Estimation af udenlandske bilers trafikarbejde i Danmark i 2009, Notatnr. 12032-005, 12 pp. Danish Road Directorate, 2010.

Hansen, K.F. & Jensen, M.G. 2004: MÅLING AF EMISSIONER FRA FREM-DRIVNINGSSANLÆG PÅ MADSS MOLS. Ruston 20RK270, Sagsnr.: 1076868, Documentation note, 5 pages (in Danish).

Hjelgaard, K. & Winther, M. 2011: The estimated new sales of Euro V diesel trucks equipped with EGR and SCR during the 2006-2010 time periods. Internal DCE note (unpublished). 2 p. (in Danish).

ICAO Annex 16 "International standards and recommended practices, Environmental protection", Volume II "Aircraft engine emissions", 3rd ed. (2008) plus amendments, 108 pp., ISBN 978-92-9231-123-0.

IFEU, 2004: Entwicklung eines Modells zur Berechnung der Luftschadstoffemissionen und des Kraftstoffverbrauchs von Verbrennungsmotoren in mobilen Geräten und Maschinen - Endbericht, UFOPLAN Nr. 299 45 113, pp. 122, Heidelberg.

IFEU 2009: Aktualisierung des Modells TREMOD - Mobile Machinery (TREMOM-MM), Endbericht; Institut für Energie- und Umweltforschung, pp. 48, Heidelberg.

IFEU 2014: Erarbeitung eines Konzepts zur Minderung der Umweltbelastung aus NRMM (non road mobile machinery) unter Berücksichtigung aktueller Emissionsfaktoren und Emissionsverminderungsoptionen für den Bestand, ISSN 1862-4804, Texte 24/2014, pp. 99, Heidelberg.

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories, IPCC, May 2000. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (06-07-2004).

Jensen, T.C., O. 2014: Dokumentation af konvertering af trafiktil emissionsopgørelser, arbejdsnotat 35187, 29 pp. DTU Transport, 2014.

Jørgensen, H. 2014: Unpublished data material from Færgen A/S.

Kristensen, F. 2008: Unpublished data material from Mols-Linjen.

Kristensen, F. 2013: Unpublished data material from Mols-Linjen.

Kristensen, F. 2014: Unpublished data material from Mols-Linjen.

Lastein, L. & Winther, M. 2003: Emission of greenhouse gases and long-range transboundary air pollutants in the Faroe Islands 1990-2001. National Environmental Research Institute. - NERI Technical Report 477. 62 pp. Available at: [http://www.dmu.dk/1\\_viden/2\\_Publikationer/3\\_fagrapporter/FR477.PDF](http://www.dmu.dk/1_viden/2_Publikationer/3_fagrapporter/FR477.PDF).

Markamp, H. 2013: Personal communication, Henrik Markamp, The National Motorcycle Association.

Ministry of Transport, 2010: TEMA2010 - et værktøj til at beregne transporters energiforbrug og emissioner i Danmark (TEMA2010 - a calculation tool for transport related fuel use and emissions in Denmark). Technical report. 135 pp.



Nielsen, O-K, Plejdrup, MS, Winther, M, Nielsen, M, Gyldenkærne, S, Mikkelsen, MH, Albrektsen, R, Thomsen, M, Hjelgaard, K, Hoffmann, L, Fauser, P, Bruun, HG, Johannsen, VK, Nord-Larsen, T, Vesterdal, L, Møller, IS, Caspersen, OH, Rasmussen, E, Petersen, SB, Baunbæk, L & Hansen, MG 2014, Denmark's National Inventory Report 2014: Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. Aarhus University, DCE - Danish Centre for Environment and Energy. Scientific Report from DCE - Danish Centre for Environment and Energy, nr. 101.

Ntziachristos, L. & Samaras, Z. 2000: COPERT III Computer Programme to Calculate Emissions from Road Transport - Methodology and Emission Factors (Version 2.1). Technical report No 49. European Environment Agency, November 2000, Copenhagen.

Nørgaard, T. & Hansen, K.F. 2004: Chiptuning af køretøjer - miljømæssig effekt, Miljøprojekt nr. 888, Miljøstyrelsen.

PHP, 1996: Research Report – Emission tests at Alpha, Mols 2 and Mols 4, 9L25MC mk6 engines #35031 and #35033, 22-23/10 1995 and 16/1 1996, DOK, PHP Basic Research, October 1996, 20 pages.

Rasmussen, H. 2014: Unpublished data material from Royal Arctic Line.

Statistics Denmark, 2014: Data from Statbank Denmark. Available at: <http://www.statistikbanken.dk/statbank5a/default.asp?w=1364>

Thorarensen, B. 2014: Unpublished data material from Eim Skip.

Winther, M. 2001a: 1998 Fuel Use and Emissions for Danish IFR Flights. Environmental Project no. 628, 2001. 112 p. Danish EPA. Prepared by the National Environmental Research Institute, Denmark. Available at: <http://www.mst.dk/udgiv/Publications/2001/87-7944-661-2/html/>

Winther, M. 2001b: Improving fuel statistics for Danish aviation. National Environmental Research Institute, Denmark. 56 p. – NERI Technical Report No. 387.

Winther, M. & Nielsen, O.K. 2006: Fuel use and emissions from non-road machinery in Denmark from 1985–2004 – and projections from 2005-2030. The Danish Environmental Protection Agency. - Environmental Project 1092: 238 pp. Available at: <http://www.dmu.dk/Udgivelser/Arbejdsrapporter/Nr.+200-249/>

Winther, M. 2008: Fuel consumption and emissions from navigation in Denmark from 1990-2005 - and projections from 2006-2030. Technical Report from NERI no. 650. 109 pp. Available at: <http://www2.dmu.dk/Pub/FR650.pdf>

Winther, M. 2009: Emission Differences between Petroleum based Diesel and different Biodiesel Blend Ratios for Road Transport Vehicles. / Winther, Morten. 2009. Transport and Air Pollution Symposium - 3rd Environment and Transport Symposium, nr. 17, Toulouse, France, 2.- 4. June 2009.

Winther, M. & Slentø, E. 2010: Heavy Metal Emissions for Danish Road Transport. National Environmental Research Institute, Aarhus University, Denmark. 99 pp. – NERI Technical Report no. 780. Available at: <http://www.dmu.dk/Pub/FR780.pdf>

Winther, M. 2011: Ændring i partikelemission som følge af fremskyndet salg af Euro 5 biler og retrofit af filtre på tunge køretøjer. Internal DCE note (unpublished). 2 p. (in Danish).

Winther, M. 2012: Danish emission inventories for road transport and other mobile sources. Inventories until the year 2010. National Environmental Research Institute, University of Aarhus. 283 pp. – DCE Scientific Report No. 24. <http://www.dmu.dk/Pub/SR24.pdf>.

Winther, M., Møller, F., Jensen, T.C. 2012: Emission consequences of introducing bio ethanol as a fuel for gasoline cars, Atmospheric Environment 55 (2012) 144-153

Wismann, T. 1999: MOLS-LINIEN, Mai Mols - Måling af emissioner fra fra hovedturbiner, dk-RAPPORT 14.901, 9 pages (in Danish).

Wismann, T. 2001: Energiforbrug og emissioner fra skibe i farvandene omkring Danmark 1995/1996 og 1999/2000 (Fuel consumption and emissions from ships in Danish coastal waters 1995/1996 and 1999/2000). The Danish Environmental Protection Agency. - Environmental Project 597: 88 pp. Available at: <http://www2.mst.dk/common/Udgivramme/Frame.asp?pg=http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-505-5/html/default.htm>.

Wismann, T. 2007: Energiforbrug for skibe i fart mellem danske havne (Fuel consumption by ships sailing between Danish ports), Internal note, September 2007, 3 pp.

### 3.4 Fugitive emissions (NFR sector 1B)

This chapter includes fugitive emissions in the NFR sector 1B.

#### 3.4.1 Source category description

According to the categorisation in the reporting format (NFR) for the UNECE CLRTAP fugitive emissions is a sub-category under the main-category Energy (Sector 1). Fugitive emissions (Sector 1B) is segmented into sub-categories covering emissions from solid fuels (1B1), oil (1B2a), natural gas (1B2b), from venting and flaring (1B2c), and other (1B2d). The sub-sectors relevant for the Danish emission inventory are shortly described below according to Danish conditions:

- 1B1c Fugitive emission from solid fuels: Coal mining is not occurring in Denmark. However, a large part of the electricity supply is based on coal fired power plants, and emissions from storage and handling of coal do occur. Emissions from solid fuels are therefore only relevant for the Danish national emission inventories in the case of particulate emissions. Emissions of other pollutants (e.g. CO) are not reported, as these emissions should be included in the inventory for the countries where the coal mining occurs.
- 1B2a Fugitive emissions from oil include emissions from exploration, extraction, storage, and transmission of crude oil, distribution of oil products and emissions from refineries.
- 1B2b Fugitive emissions from natural gas include emissions from exploration, extraction, transmission and distribution of natural gas.
- 1B2c Venting and flaring include activities onshore and offshore. Flaring occur both offshore and onshore in gas treatment and storage plants and in refineries. Venting occur in gas storage plants. Venting of gas is not occurring in extraction and in refineries as controlled venting enters the gas flare system.

Activity data, emission factors and emissions are stored in the Danish emission database on SNAP sector categories (Selected Nomenclature for Air Pollution). In Table 3.4.1 the corresponding SNAP codes and NFR sectors relevant to fugitive emissions are shown. Further, the table holds the SNAP names for the SNAP codes and the overall activity (e.g. oil and natural gas).

Table 3.4.1 List of NFR sectors relevant for fugitive emissions, and the corresponding SNAP codes and emission sources.

NFR sectors	SNAP Code	SNAP name	Fuel/source
1B1a	050103*	Storage of solid fuel	Coal mining and handling
1B2ai	050204	Exploration of oil	Oil
1B2ai	050201	Land-based activities	Oil
1B2ai	050202	Offshore activities	Oil
1B2aiv	040101	Petroleum products processing	Oil
1B2aiv	040103	Sulphur recovery plants	Oil
1B2av	050503	Service stations (including refuelling of cars)	Oil
1B2b	050304	Exploration of gas	Natural gas
1B2b	050303	Offshore activities	Natural gas
1B2b	050601	Transmissions and storage	Natural gas
1B2b	050603	Distribution	Natural gas
1B2c	050699	Venting in gas storage	Venting
1B2c	090203	Flaring in oil refinery	Flaring
1B2c	090206	Flaring in oil and gas extraction	Flaring
1B2c	090298	Flaring in gas storage	Flaring
1B2c	090299	Flaring in gas transmission and distribution	Flaring

<sup>1</sup> Emissions from gas extraction are included in 1B2a snap 050204, Exploration of oil.

Table 3.4.2 summarizes the Danish fugitive emissions in 2013. The methodologies, activity data and emission factors used for calculation are described in the following chapters.

Table 3.4.2 Summary of the Danish fugitive emission in 2013. P refers to point source and A to area source. Minor emission sources are excluded from the table.

IPCC code	SNAP code	Source	Pollutant	Emission	Unit
1B2a iv	040101	P	NMVOC	5575	Mg
1B2a iv	040103	P	SO <sub>2</sub>	517	Mg
1B2a iv	040104 <sup>1</sup>	A	Mg	---	Mg
1B1a	050103	A	TSP	743	Mg
1B1a	050103	A	PM <sub>10</sub>	297	Mg
1B1a	050103	A	PM <sub>2,5</sub>	30	Mg
1B1a	050103	A	BC	495	Mg
1B2a i	050201	A	NMVOC	1330	Mg
1B2a i	050202	A	NMVOC	1395	Mg
1B2a i	050204 <sup>2</sup>	A			
1B2b	050304 <sup>3</sup>	A			
1B2a v	050503	A	NMVOC	725	Mg
1B2b	050601	A	NMVOC	4	Mg
1B2b	050603	A	NMVOC	44	Mg
1B2c	050699	P	NMVOC	18	Mg
1B2c	090203	P	SO <sub>2</sub>	166	Mg
1B2c	090203	P	NO <sub>x</sub>	15	Mg
1B2c	090203	P	NMVOC	22	Mg
1B2c	090203	P	CO	52	Mg
1B2c	090203	P	TSP	0,3	Mg
1B2c	090203	P	PM <sub>10</sub>	0,3	Mg
1B2c	090203	P	PM <sub>2,5</sub>	0,3	Mg
1B2c	090203	P	BC	0,1	Mg
1B2c	090203	P	As	0,1	kg
1B2c	090203	P	Cd	0,2	kg
1B2c	090203	P	Cr	0,9	kg
1B2c	090203	P	Cu	0,6	kg
1B2c	090203	P	Hg	< 0,1	kg
1B2c	090203	P	Ni	1,2	kg
1B2c	090203	P	Pb	0,6	kg
1B2c	090203	P	Se	0,1	kg
1B2c	090203	P	Zn	7,6	kg
1B2c	090203	P	PCDD/F	< 0,01	g
1B2c	090203	P	Benzo(b)	< 0,01	kg
1B2c	090203	P	Benzo(k)	< 0,01	kg
1B2c	090203	P	Benzo(a)	< 0,01	kg
1B2c	090203	P	Indeno	< 0,01	kg
1B2c	090206	A	SO <sub>2</sub>	1	Mg
1B2c	090206	A	NO <sub>x</sub>	106	Mg
1B2c	090206	A	NMVOC	129	Mg
1B2c	090206	A	CO	161	Mg
1B2c	090206	A	TSP	4	Mg
1B2c	090206	A	PM <sub>10</sub>	4	Mg
1B2c	090206	A	PM <sub>2,5</sub>	4	Mg
1B2c	090206	A	BC	1	Mg

Continued

1B2c	090206	A	As	0,5	kg
1B2c	090206	A	Cd	< 0,1	kg
1B2c	090206	A	Cr	< 0,1	kg
1B2c	090206	A	Cu	< 0,1	kg
1B2c	090206	A	Hg	0,4	kg
1B2c	090206	A	Ni	< 0,1	kg
1B2c	090206	A	Pb	7,3	kg
1B2c	090206	A	Se	< 0,1	kg
1B2c	090206	A	Zn	< 0,1	kg
1B2c	090206	A	PCDD/F	< 0,01	g
1B2c	090206	A	Benzo(b)	< 0,01	kg
1B2c	090206	A	Benzo(k)	< 0,01	kg
1B2c	090206	A	Benzo(a)	< 0,01	kg
1B2c	090206	A	Indeno	< 0,01	kg
1B2c	090298 <sup>4</sup>	P	SO <sub>2</sub>	< 0,1	Mg
1B2c	090298 <sup>4</sup>	P	NO <sub>x</sub>	1,3	Mg
1B2c	090298 <sup>4</sup>	P	NMVOG	0,3	Mg
1B2c	090298 <sup>4</sup>	P	CO	1,5	Mg
1B2c	090299 <sup>5</sup>	A	SO <sub>2</sub>	< 0,1	Mg
1B2c	090299 <sup>5</sup>	A	NO <sub>x</sub>	0,1	Mg
1B2c	090299 <sup>5</sup>	A	NMVOG	0,1	Mg
1B2c	090299 <sup>5</sup>	A	CO	0,1	Mg

<sup>1</sup> Not occurring in 2013.

<sup>2</sup> Exploration of oil are excluded for all pollutants (main pollutants < 0.1 Mg, HM < 0.1 kg, PCDD/F < 0.01 g, and PAH < 0.01 kg).

<sup>3</sup> Exploration of gas are excluded for all pollutants (main pollutants < 0.1 Mg, HM < 0.1 kg, PCDD/F < 0.01 g, and PAH < 0.01 kg).

<sup>4</sup> Flaring in gas storage are excluded for PM, BC, HM, POPs (PM < 0.1 Mg, HM < 0.1 kg, PCDD/F < 0.01 g, and PAH < 0.01 kg).

<sup>5</sup> Flaring in gas transmission and distribution are excluded for PM, BC, HM and POPs (PM < 0.1 Mg, HM < 0.1 kg, PCDD/F < 0.01 g, and PAH < 0.01 kg).

### 3.4.2 Methodological issues

The following chapters give descriptions on the methods of calculation used in the Danish emission inventory. Further, the activity data and emission factors that form the basis for the calculations are described according to data source and values.

#### Fugitive emissions from solid fuels

The emissions of particulate matter from storage of coal are estimated on basis of the imported amount of coal (equation 3.4.1).

$$E_{coal\_storage} = EMF_{coal\_storage} \cdot I_{coal} \quad (\text{Eq. 3.4.1})$$

where  $EMF_{coal\_storage}$  is the emission factor for storage of coal in coal piles and  $I_{coal}$  is the amount of coal imported in the actual year.

### Fugitive emissions from oil

The emissions from oil derive from offshore activities, service stations and refineries. Emissions from offshore activities include emissions from exploration, production, onshore oil tanks and onshore and offshore loading of ships. In the case of service stations emissions from reloading of tankers and refuelling of vehicles are included. The emissions from refineries derive from petroleum products processing (oil refining). Emissions from flaring and venting are included in the chapters concerning flaring. The total fugitive emission from oil can be expressed as:

$$E_{total} = E_{exploration} + E_{extraction} + E_{ship} + E_{oil\ tanks} \quad (\text{Eq. 3.4.2})$$

### Fugitive emissions from exploration

Fugitive emissions from exploration of oil are based on amounts and composition per exploration drilling. As calorific values and densities are not available per drilling, data from a gas test in 1992 are used.

### Fugitive emissions from extraction

The methodology for calculation of emissions from extraction is based on standard emission factors from the 2006 IPCC Guidelines (IPCC, 2006).

### Loading of ships

Fugitive emissions of NMVOC from loading of ships include the transfer of oil from storage tanks or directly from the well into ships. The activity also includes losses during transport. When oil is loaded hydrocarbon vapour will be displaced by oil and new vapour will be formed, both leading to emissions. The emissions from ships are calculated by equation 3.4.3.

$$E_{ships} = EMF_{ships, onshore} \cdot L_{oil, onshore} + EMF_{ships, offshore} \cdot L_{oil, offshore} \quad (\text{Eq. 3.4.3})$$

where  $EMF_{ships}$  is the emission factor for loading of ships and  $L_{oil}$  is the amount of oil loaded.

### Oil tanks

The NMVOC emissions for storage of crude oil are given in the environmental reports for the raw oil terminal from DONG Oil Pipe for 2013 (DONG Oil Pipe A/S, 2014). An implied emission factor is calculated based on the amount of oil transported in pipelines according to equation 3.4.4.

$$IEF_{tanks} = \frac{E_{tanks}}{T_{oil}} \quad (\text{Eq. 3.4.4})$$

where  $IEF_{tanks}$  is the implied emission factor for storage of raw oil in tanks,  $E_{tanks}$  is the emission and  $T_{oil}$  is the amount of oil transported in pipelines.

### Service stations

NMVOC emissions from service stations are estimated as outlined in equation 3.4.5.

$$E_{service\ stations} = (EMF_{reloading} \cdot T_{fuel}) + (EMF_{refuelling} \cdot T_{fuel}) \quad (\text{Eq. 3.4.5})$$

where  $EMF_{reloading}$  is the emission factor for reloading of tankers to underground storage tanks at the service stations,  $EMF_{refuelling}$  is the emission factor for refuelling of vehicles and  $T_{fuel}$  is the amount of gasoline used for road transport.

### **Oil refining**

When oil is processed in the refineries, part of the volatile organic compounds (VOC) is emitted to the atmosphere. The VOC emissions from the oil refinery process include non-combustion emissions from handling and storage of feedstock (raw oil), from the petroleum product processing and from handling and storage of products. When only the total VOC emission is given by a refinery the emission of NMVOC is estimated based on the assumption that 10 % of VOC is CH<sub>4</sub> and the remaining 90 % is NMVOC (Hjerrild and Rasmussen, 2014).

Both the non-combustion processes, product processing and sulphur recovery plants emit SO<sub>2</sub>. The SO<sub>2</sub> emissions are calculated by the refineries and implemented in the emission inventory without further calculation.

Emissions from flaring in refineries are described under "Flaring". Emissions related to process furnaces in refineries are included in stationary combustion with the relevant emission factors.

### **Transmission and distribution of gas**

The fugitive emission from transmission, storage and distribution of natural gas is based on information from the gas companies. The transmission and distribution companies give data on the transported amount and length and material of the pipeline systems.

The fugitive losses from pipelines are only given for some companies, here among the transmission company. The available distribution data are used for the remaining companies too. Emissions are calculated from the fugitive losses from transmission and distribution pipelines due to the gas quality measured by Energinet.dk. The same approach is used for town gas, which is natural gas admixed ~ 50 % ambient air.

Calculations of emissions from distribution of town gas are based on data from the distribution companies on distribution losses. At present, there are two areas with town gas distribution and correspondingly distribution companies. Two others companies in other areas were closed in 2004 and 2006, and it have not been possible to collect data for all years in the time series. The emissions have been calculated for the years with available data and the distribution loss for the first year with data has been applied for the previous years in the time series. Data is missing for the later years (1996-2003) for one of the distribution companies. The distribution amount is assumed to decrease linearly to zero over these years, and the share ("distribution loss"/"distribution amount") are assumed equal to the value for 1995.

### **Venting and flaring**

Emissions from flaring are estimated from the amount of gas vented and flared offshore, in gas treatment/storage plants, in refineries, and in gas transmission and distribution, combined with corresponding emission factors. From 2006 data on offshore flaring is given in the reports for the European Union Greenhouse Gas Emission Trading System (EU ETS) and thereby flaring can be split to the individual production units. Before 2006 only the total flared amount is available.



### 3.4.3 Activity data

#### Coal storage

The activity data are the imported amount of coal converted using the calorific values of coal (Danish Energy Agency, 2014c). The coal import fluctuates between years in the time series with an overall decreasing trend. In 2013 the imported amount was 4 952 Gg (Figure 3.4.1) which is an increase since 2012.




Figure 3.4.1 Amount of imported coal.

#### Exploration and extraction of oil and gas, and loading of ships

Activity data used in the calculations of the emissions from oil and gas exploration and production, and loading of ships onshore and offshore are shown in Table 3.4.3. Data are based on information provided by the Danish Energy Agency (Andersen 2014, DEA 2014a) and from the environmental reports from DONG Oil Pipe A/S (2014).

Table 3.4.3 Activity data for 2013.

Activity	Value	Data source
Extracted gas, Sm <sup>3</sup>	1 242	Andersen, 2014
Extracted oil, Nm <sup>3</sup>	102	Andersen, 2014
Produced gas, 10 <sup>6</sup> Nm <sup>3</sup>	4 704	Danish Energy Agency, 2014a
Produced oil, 10 <sup>3</sup> m <sup>3</sup>	10 185	Danish Energy Agency, 2014a
Oil loaded, 10 <sup>3</sup> m <sup>3</sup>	1 549	Danish Energy Agency, 2014a
Oil loaded, 10 <sup>3</sup> m <sup>3</sup>	10 500	DONG Oil Pipe A/S, 2014
Density of crude oil, tonnes/m <sup>3</sup>	0,86	Danish Energy Agency, 2014a

Exploration of oil and gas vary between years. The largest oil amounts are seen for 1990, 2002 and 2005, while relatively large gas amounts are seen for more years of the time series. Explored amounts are shown in Figure 3.4.2.




Figure 3.4.2. Exploration of oil and gas in the Danish part of the North Sea.

As seen in Figure 3.4.3 the production of oil and gas in the North Sea has generally increased in the years 1990-2004, and since 2004 the production has decreased. Five major platforms were completed in 1997-1999, which is the main reason for the great increase in the oil production in the years 1998-2000.




Figure 3.4.3. Production of oil and gas in the Danish part of the North Sea.

The amount of oil loaded offshore on ships roughly follows the trend of the oil and gas production (Figure 3.4.4). In case of onshore loading of ships the trend is more smoothed.




Figure 3.4.4 Onshore and offshore loading of ships.

### Oil refining

Data on the amount of crude oil processed in the two Danish refineries are given by the refineries in their annual environmental report (A/S Dansk Shell, 2014 and Statoil A/S, 2014). Until 1996 a third refinery was in operation, leading to a decrease in the crude oil amount from 1996 to 1997. Data are shown in Figure 3.4.5. In 2013 the amount of crude oil being processed was 7 179 Gg.




Figure 3.4.5 Oil refineries. Processed crude oil in Danish refineries.

### Service stations

The Danish Energy statistics holds data on the sale of gasoline that is the basis for estimating emissions of NMVOC from service stations. The gasoline sales show an increase from 1990-1998 and a decreasing trend since 1999 as shown in Figure 3.4.6. In 2013 the gasoline sale was 1 339 Gg.




Figure 3.4.6 Gasoline sales in Denmark.

### Transmission, storage and distribution of gas

The activity data used in the calculation of the emissions from natural gas are shown in Table 3.4.4. Transmission rates for 1990-1998 refer to annual environmental reports of DONG Energy. In 1999-2006 transmission rates refer to the Danish Gas Technology Centre (Karll 2002, Karll 2003, Karll 2004, Karll 2005, Oertenblad 2006, Oertenblad 2007). From 2008 onwards transmission rates refer to Energinet.dk. Transmission losses for 1991-1999 are based on annual environmental report of DONG Energy. The average for 1991-1995 is applied for 1990. From 2005 onwards transmission losses are given by Energinet.dk. The average for 2005-2010 is applied for the years 2000-2004.

Distribution rates for 1990-1998 are estimated from the Danish energy statistics. Distribution rates are assumed to equal total Danish consumption rate minus the consumption rates of sectors that receive the gas at high pressure. The following consumers are assumed to receive high pressure gas: town gas production companies, production platforms and power plants. In 1999-2006 distribution rates refer to DONG Energy / Danish Gas Technology Centre / Danish gas distribution companies (Karll 2002, Karll 2003, Karll 2004, Karll 2005, Oertenblad 2006, Oertenblad 2007). Since 2007 the distribution rates are given by the companies. Distribution rates for town gas is based on the available data from the Danish town gas distribution companies of which more are closed down today. Distribution losses for 1990-2000 are based on annual environmental report of DONG Energy. For 2000-2006 the average losses-% for the gas distribution companies are used. From 2007 data on distribution losses available from the companies are used.

Table 3.4.4 Activity data on transmission and distribution of gas for selected years of the time series. Town gas is included in distribution.

	1990	1995	2000	2005	2010	2011	2012	2013
Transmission, Mm <sup>3</sup> <sup>1</sup>	2739	4689	7079	7600	7462	6181	5365	4886
Distribution of natural gas, Mm <sup>3</sup> <sup>2</sup>	1714	3054	3181	3265	3416	2933	2728	2634
Distribution of town gas, Mm <sup>3</sup> <sup>3</sup>	35	35	34	32	22	21	24	28

<sup>1</sup> Transmission rates for 1990-1998 refer to the annual environmental report of DONG Energy. In 1999-2006 transmission rates refer to the Danish Gas Technology Centre (Karll 2002, Karll 2003, Karll 2004, Karll 2005, Oertenblad 2006, Oertenblad 2007). From 2008 onwards transmission rates refer to Energinet.dk.

<sup>2</sup> In 1990-98 distribution rates are estimated from the Danish energy statistics. Distribution rates are assumed to equal total Danish consumption rate minus the consumption rates of sectors that receive the gas at high pressure. The following consumers are assumed to receive high pressure gas: town gas production companies, production platforms and power plants. In 1999-2006 distribution rates refer to DONG Energy / Danish Gas Technology Centre / Danish gas distribution companies (Karll 2002, Karll 2003, Karll 2004, Karll 2005, Oertenblad 2006, Oertenblad 2007). Since 2007 the distribution rates are given by the companies.

<sup>3</sup> The distribution of town gas is based on the available data from the Danish town gas distribution companies of which more are closed down today.

In 2013 the gas transmission rate was 4 886 Mm<sup>3</sup> and the distribution rate was 2 662 Mm<sup>3</sup>, hereof 28 Mm<sup>3</sup> town gas (Figure 3.4.7). The variation over the time series owes mainly to variations in the winter temperature and to the variation of import/export of electricity from Norway and Sweden. The transmission rate is less than the production rate, as part of the produced natural gas is exported through the NOGAT pipeline system.




Figure 3.4.7 Rates for transmission and distribution of gas. Distribution covers both natural gas and town gas.

Data on the transmission pipelines excluding offshore pipelines and on the distribution network are given by Energinet.dk, DGC and the distribution companies concerning length and material. The length of the transmission pipelines is approximately 900 km. Because the distribution system in Denmark is relatively new most of the distribution network is made of plastic (PE). In 2013 the length of the distribution network was around 20 000 km. The major part is made of plastic (approximately 90 %) and the remaining part is made of steel. For this reason the fugitive emission is negligible under normal operating conditions as the distribution system is basically tight with no fugitive losses. However, the plastic pipes are vulnerable and therefore most of the fugitive emissions from the pipes are caused by losses due to ex-

cavation damages and construction and maintenance activities performed by the gas companies. These losses are either measured or estimated by calculation in each case by the gas companies. About 5 % of the distribution network is used for town gas. This part of the network is older and the fugitive losses are larger. The fugitive losses from this network are associated with more uncertainty as it is estimated as a percentage (15 %) of the meter differential. This assumption is based on expert judgement from one of the town gas companies (Jensen, 2008). It must be noted that two town gas distribution companies have been closed in recent years (one in 2004 and another in 2006). There are only two town gas distribution companies left, and therefore the data availability is scarce.

## Venting and Flaring

### Venting

In Denmark there are two natural gas storage facilities. Both are obligated to make an environmental report on annual basis. Data on gas input and withdrawal are included and were 891 Mm<sup>3</sup> and 650 Mm<sup>3</sup> in 2013, respectively. Venting and flaring at the gas storage plants are included in the inventory. Venting of gas is assumed to be not occurring in extraction and in refineries as controlled venting enters the gas flare system. Venting rates in gas storage facilities are shown in Figure 3.4.8. As venting rates are not available for the years 1990-1994, the average for 1995-1998 is used.




Figure 3.4.8 Amount vented in gas treatment and storage plants.

### Flaring

Flaring rates for the two Danish refineries are given in their environmental reports and in additional data provided by the refineries directly to DCE. From 2006 flaring amounts are given in the EU ETS reporting. Data are not available for the years 1990-1993. The flaring amount for 1994 has been adopted for the previous years. Flaring rates are shown in Figure 3.4.9.




Figure 3.4.9 Amount flared in refineries (annual environmental reports from A/S Dansk Shell and Statoil A/).

Offshore flaring amounts are given in Denmark’s oil and gas production (Danish Energy Agency, 2014a) while flaring in treatment/storage plants are given in DONG Energy’s environmental reports (Dong Energy, 2014a; Dong Energy, 2014b; Energinet.dk, 2014). Flaring rates are shown in Figure 3.4.10 and 3.4.11.

Offshore flaring amounts have been decreasing over the last 10 years period in accordance with the decrease in production as seen in Figure 3.4.3. Further, there is focus on reduction of the amount being flared for environmental reasons.




Figure 3.4.10 Amounts of gas flared in offshore exploration.

Flaring rates in gas treatment and gas storage plants are not available before 1994. The mean value for 1994-1998 has been adopted as basis for the emission calculation for the years 1990-1993. The large amount of flared gas in 2007 owe to a larger maintenance work at the gas treatment plant. Flaring rates are shown in Figure 3.4.11.




Figure 3.4.11 Amount flared in gas treatment and storage plants.

### 3.4.4 Emission factors

#### Coal storage

Emissions of particulate matter (PM) from coal storage are estimated using emission factors from the Coordinated European Particulate Matter Emission Inventory Program, CEPMEIP (Visschedijk et al., 2004). The emission factors are listed in Table 3.4.5.

Table 3.4.5 Emission factors used to estimate particulate emissions from coal storage.

Emission factor	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>
Emission factor, g per Mg	150	60	6

#### Exploration of oil and gas

Emissions from exploration are calculated from the same emissions that are used for offshore flaring. The emission factors, which are based on DEPA 2008 and EMEP/EEA 2013, are listed in Table 3.4.13.

#### Extraction of oil and gas

Standard emission factors from the 2006 IPCC Guidelines (IPCC 2006) are used to calculate emissions from extraction of oil and gas. The emission factors are listed in Table 3.4.6.

#### Oil terminal

Implied emission factors are calculated annually for NMVOC based on emission provided in the environmental reports for the raw oil terminal (DONG Oil Pipe A/S, 2014). The implied emission factor for 2013 is listed in Table 3.5.6.

Table 3.4.6 Emission factors for extraction of oil and gas.

	NMVOC	Reference
Extraction of oil, Gg/1000m <sup>3</sup>	7,40E-07	IPCC 2006
Extraction of gas, Gg/Mm <sup>3</sup>	9,10E-05	IPCC 2006
Oil terminal, g/Mg oil extracted	62.0	IPCC 2006, DONG Oil Pipe A/S 2014

#### Loading of ships

In the EMEP/EEA Guidebook standard emission factors for different countries are given (EMEP/EEA, 2013). In the Danish emission inventory the



Norwegian emission factors are used for estimation of fugitive emissions from loading of ships onshore and offshore for the years 1990-2009. During 2009 new emission reducing technologies (degassing unit) were installed at the crude oil terminal. Measurements were carried out at the terminal before and after installation show a decrease of 25 % of the NMVOC emission from loading of ships. The reduced emission factors are used for 2010 onwards. The emission factors are listed in Table 3.4.7.

Table 3.4.7 Emission factors for loading of ships onshore and offshore.

	NMVOC, fraction of loaded	Reference
Ships off-shore	0.001	EMEP/EEA, 2013
Ships on-shore, 1990-2009	0.0002	EMEP/EEA, 2013
Ships on-shore, 2010 onwards	0.00015	EMEP/EEA, 2013; Miljøcenter Odense, 2012

### Oil refining

The refineries provide information on consumption of fuel gas and fuel oil. The calorific values are given by the refineries in the reporting for EU ETS since 2006. Before 2006 the calorific values given by the refineries were used when available. When not available standard calorific values from the Danish Energy Agency combined with the conversion factor between fuel gas and fuel oil given by the refinery were used for calculation.

Emissions of SO<sub>2</sub>, NO<sub>x</sub> and VOC are given by the refineries. Only one of the two refineries has made a split between NMVOC and CH<sub>4</sub>. For the other refinery it is assumed that 10 % of the VOC emission is CH<sub>4</sub> and the remaining 90 % is NMVOC (Hjerrild & Rasmussen, 2014).

### Service stations

The NMVOC emission from service stations is calculated by use of different emission factors for the time series as shown in Table 3.4.8.

In 1994 the emission factors for NMVOC from service stations were investigated by Fenhann and Kilde (1994) for 1990 and 1991, individually. The emission factors reported for reloading for 1990 are used for the years 1985-1990, while the emission factor for 1991 is used for 1991 only. In 1995 Stage I was made obligatory, and the emission factor from the 2013 EMEP/EEA Guidebook (EMEP/EEA, 2013) is applied from 1995 and onwards. Linear interpolation is applied for the years 1992-1994.

Fenhann and Kilde (1994) also include NMVOC emission factors for refuelling for the years 1990, 1991, 1992, and 1993. The same value is given for the three years, and this emission factor is applied for the years 1994-1995, when the first legal acts on emission reduction from service stations turned into force. From 2005 the refuelling emission factor is based on the 2013 EMEP/EEA Guidebook (EMEP/EEA, 2013). An abatement rate of 85 % is given in the 2013 EMEP/EEA Guidebook, while 60 % were given in the 2006 EMEP/EEA Guidebook (EMEP/EEA, 2006). The Danish requirement is 85 % abatement under optimal conditions, but 70 % in practice occurrence (Danish Ministry of the Environment, 2011). Based on this, 70 % abatement is applied in the emission calculations. Linear interpolation is used from 1996-2004.

Table 3.4.8 Emission factors used for estimating NMVOC from service stations.

Year	Reloading of tankers, kg NMVOC pr tonnes gasoline	Refuelling of vehicles, kg NMVOC pr tonnes gasoline	Sum of reloading and refuelling, kg NMVOC pr tonnes gasoline	Source
1985-1990	1,28	1,52	2,80	Fenhann & Kilde,1994
1991	0,64	1,52	2,16	Fenhann & Kilde,1994
1992	0,49	1,52	2,01	Interpolation / Fenhann & Kilde,1994
1993	0,35	1,52	1,87	Interpolation / Fenhann & Kilde,1994
1994	0,20	1,52	1,72	Interpolation / Fenhann & Kilde,1994
1995	0,05	1,52	1,57	EMEP/EEA 2013 / Fenhann & Kilde,1994
1996	0,05	1,42	1,47	EMEP/EEA 2013 / interpolation
1997	0,05	1,31	1,37	EMEP/EEA 2013 / interpolation
1998	0,05	1,21	1,26	EMEP/EEA 2013 / interpolation
1999	0,05	1,11	1,16	EMEP/EEA 2013 / interpolation
2000	0,05	1,00	1,06	EMEP/EEA 2013 / interpolation
2001	0,05	0,90	0,95	EMEP/EEA 2013 / interpolation
2002	0,05	0,80	0,85	EMEP/EEA 2013 / interpolation
2003	0,05	0,69	0,75	EMEP/EEA 2013 / interpolation
2004	0,05	0,59	0,64	EMEP/EEA 2013 / interpolation
2005 onwards	0,05	0,49	0,54	EMEP/EEA 2013

### Transmission, storage and distribution of gas

The fugitive emissions from transmission, storage and distribution of natural gas are based on data on gas losses from the companies and on the average annual natural gas composition given by Energinet.dk (Table 3.4.9). For distribution of town gas the emission factor is reduced due to the admixture of 50 % atmospheric air to the natural gas.

Table 3.4.9 Annual gas composition, lower heating value and density for Danish natural gas (Energinet.dk).

		Unit	1990	2000	2005	2010	2011	2012	2013
Methane	CH <sub>4</sub>	molar-%	90.92	86.97	88.97	89.95	89.10	88.84	89,82
Ethane	C <sub>2</sub> H <sub>6</sub>	molar-%	5.08	6.88	6.14	5.71	5.98	6.11	5,65
Propane	C <sub>3</sub> H <sub>8</sub>	molar-%	1.89	3.17	2.50	2.19	2.36	2.44	2,00
i-Butane	i-C <sub>4</sub> H <sub>10</sub>	molar-%	0.36	0.43	0.40	0.37	0.37	0.37	0,32
n-Butane	n-C <sub>4</sub> H <sub>10</sub>	molar-%	0.50	0.61	0.55	0.54	0.55	0.54	0,46
i-Petane	i-C <sub>5</sub> H <sub>12</sub>	molar-%	0.14	0.11	0.11	0.13	0.13	0.13	0,11
n-Petane	n-C <sub>5</sub> H <sub>12</sub>	molar-%	0.10	0.08	0.08	0.08	0.09	0.08	0,08
n-Hexane and heavier hydrocarbons	C <sup>6+</sup>	molar-%	0.09	0.06	0.05	0.06	0.06	0.06	0,05
Nitrogen	N <sub>2</sub>	molar-%	0.31	0.34	0.29	0.31	0.37	0.36	0,55
Carbon dioxide	CO <sub>2</sub>	molar-%	0.60	1.35	0.90	0.66	0.98	1.06	0,96
Lower heating value	H <sub>n</sub>	MJ/m <sup>3</sup> <sub>n</sub>	39.176	40.154	39.671	39.461	39.507	39.548	39.988
Density	pp	kg/m <sup>3</sup> <sub>n</sub>	0.808	0.846	0.825	0.816	0.824	0.827	0.814

### Venting and flaring

#### Venting

Emissions of NMVOC from venting are given in the environmental reports for the gas storage plants (DONG Energy, 2014a; Energinet.dk, 2014).

#### Flaring in refineries

The composition of fuel gas is given for 2008 by one of the two refineries. As the composition for fuel gas is marked different than the composition of natural gas, which has been used in earlier year's calculations, the same fuel gas composition is used in calculations for the other Danish refinery.

The NMVOC emission factor based on these data are applied for both refineries for the entire time series. Emission factors for the remaining pollutants are taken from the 2013 EMEP/EEA Guidebook. The refinery emission factors are listed in Table 3.4.10.

NO<sub>x</sub> emissions are provided by one refinery. Emission factors from the EMEP/EEA Guidebook (2013) are used to calculate emissions of CO and for NO<sub>x</sub> where plant specific data are not available. Emission factors for the remaining pollutants are taken from the 2013 EMEP/EEA Guidebook. The refinery emission factors are listed in Table 3.4.10.

Table 3.4.10 Emission factors for flaring in refineries.

Pollutant	Emission factor	Unit	Source
NO <sub>x</sub>	32.2	g per GJ	Country specific
NMVOC	76.4	g per GJ	Country specific
CO	177	g per GJ	EMEP/EEA, 2013
TSP	0.89	g per GJ	EMEP/EEA, 2013
PM <sub>10</sub>	0.89	g per GJ	EMEP/EEA, 2013
PM <sub>2.5</sub>	0.89	g per GJ	EMEP/EEA, 2013
BC	0.22	g per GJ	EMEP/EEA, 2013
As	0,3	mg per GJ	EMEP/EEA, 2013
Cd	0,7	mg per GJ	EMEP/EEA, 2013
Cr	3	mg per GJ	EMEP/EEA, 2013
Cu	2	mg per GJ	EMEP/EEA, 2013
Hg	0,09	mg per GJ	EMEP/EEA, 2013
Ni	4	mg per GJ	EMEP/EEA, 2013
Pb	2	mg per GJ	EMEP/EEA, 2013
Se	0,42	mg per GJ	EMEP/EEA, 2013
Zn	26	mg per GJ	EMEP/EEA, 2013
Dioxin	0,5	ng I-TEQ per GJ	EMEP/EEA, 2013
Benzo(b)fluoranthene	1,14	µg per GJ	EMEP/EEA, 2013
Benzo(k)fluoranthene	0,63	µg per GJ	EMEP/EEA, 2013
Benzo(a)pyrene	0,67	µg per GJ	EMEP/EEA, 2013
Indeno(1,2,3-c,d)pyrene	0.63	µg per GJ	EMEP/EEA, 2013

#### ***Flaring offshore***

The emission factors for offshore flaring are shown in Table 3.4.11. The NO<sub>x</sub> emission factor is based on the conclusion in a Danish study of NO<sub>x</sub> emissions from offshore flaring carried out by the Danish Environmental Protection Agency (DEPA, 2008). The recommended NO<sub>x</sub> emission factor (31 008 g per GJ or 0.0015 tonnes NO<sub>x</sub> per tonnes gas) corresponds well with the emission factors used to estimate NO<sub>x</sub> emission in other countries with oil production in the North Sea (Netherlands: approximately 0.0014 tonnes NO<sub>x</sub> per tonnes gas and United Kingdom: approximately 0.0013 tonnes NO<sub>x</sub> per tonnes gas). Emission factors for all other pollutants are based on the 2013 EMEP/EEA Guidebook (2013).

Table 3.4.11 Emission factors for offshore flaring.

Pollutant	Emission factor	Unit	Source
SO <sub>2</sub>	0.013	g per Nm <sup>3</sup>	EMEP/EEA, 2013
NO <sub>x</sub>	1.227	g per Nm <sup>3</sup>	DEPA, 2008
NM VOC	1.482	g per Nm <sup>3</sup>	EMEP/EEA, 2013
CO	1.854	g per Nm <sup>3</sup>	EMEP/EEA, 2013
TSP	0.042	g per Nm <sup>3</sup>	EMEP/EEA, 2013
PM <sub>10</sub>	0.042	g per Nm <sup>3</sup>	EMEP/EEA, 2013
PM <sub>2.5</sub>	0.042	g per Nm <sup>3</sup>	EMEP/EEA, 2013
BC	0.008	g per Nm <sup>3</sup>	EMEP/EEA, 2013
As	0.006	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Cd	>0.001	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Cr	>0.001	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Cu	>0.001	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Hg	0.005	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Ni	>0.001	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Pb	0.084	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Se	0.001	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Zn	>0.001	mg per Nm <sup>3</sup>	EMEP/EEA, 2013
Dioxin	0.024	ng I-TEQ per Nm <sup>3</sup>	EMEP/EEA, 2013
Benzo(b)fluoranthene	0.040	µg per Nm <sup>3</sup>	EMEP/EEA, 2013
Benzo(k)fluoranthene	0.040	µg per Nm <sup>3</sup>	EMEP/EEA, 2013
Benzo(a)pyrene	0.026	µg per Nm <sup>3</sup>	EMEP/EEA, 2013
Indeno(1,2,3-c,d)pyrene	0.040	µg per Nm <sup>3</sup>	EMEP/EEA, 2013

### 3.4.5 Emissions

#### Coal storage

The emission from storage of coal is 743Mg TSP in 2013 (297 Mg PM<sub>10</sub> and 30 Mg PM<sub>2.5</sub>). The coal consumption and the related emissions vary from year to year mainly due to the extent of electricity import/export and temperature variations (Table 3.4.12). Note that PM was only included in the inventory from 2000.

Table 3.4.12 PM<sub>10</sub> from storage of solid fuels for selected years of the time series.

	2000	2005	2010	2011	2012	2013
TSP, Mg	962	905	686	920	597	743
PM <sub>10</sub> , Mg	385	362	274	368	239	297
PM <sub>2.5</sub> , Mg	38	36	27	37	24	30

#### Exploration and extraction of oil and gas, loading of ships and oil tanks

NM VOC emissions from exploration and extraction of oil and gas, and from loading of ships are listed in Table 3.4.13, along with emissions from the oil terminal given in the environmental reports from DONG Oil Pipe (2014). A degassing system has been established at the crude oil terminal leading to reduced VOC emissions from storage and handling. The degassing system has been in operation since the summer of 2009 and measurements of VOC emissions were carried out in September 2009 after a period with constant operation. The NM VOC emission factor from the oil terminal has decreased by 38 % from 2008 to 2009 and further by 61 % from 2009 to 2010.

Table 3.4.13 NMVOC (Mg) from onshore activities related to extraction of oil and gas.

NMVOC, Mg	1990	1995	2000	2005	2010	2011	2012	2013
Exploration of oil	0,0029	0,0021	0,0016	0,0028	NO	NO	NO	0,0018
Exploration of gas	4,2867	0,0002	0,083	1,0569	NO	NO	NO	0,0002
Extraction of oil	5,2	8	15,6	16,2	10,5	9,5	8,7	7,5
Extraction of gas	467	575	1030	1049	733	593	511	428
Onshore loading of ships	678	1249	2183	2494	1187	1071	1355	787
Offshore loading of ships	NO	NO	4021	3337	1658	1525	1332	960
Oil tanks	1726	2664	4000	4500	763	638	594	543

### Oil refining

NMVOC emissions from oil refining at the Danish refineries are listed for selected years of the time series in Table 3.4.14. Further, the emissions of SO<sub>2</sub> from oil refining and sulphur recovery in refineries are shown. The emission of SO<sub>2</sub> has shown a pronounced decrease since 1990 because of technical improvements at the refineries. Note that SO<sub>2</sub> from refining and recovery prior to 1994 was summarised and reported as an area source in category 1B2a vi, and that SO<sub>2</sub> from oil refining from 2001 are included in stationary combustion.

Table 3.4.14 Oil Refineries. Emissions of NMVOC and SO<sub>2</sub> from oil refining and SO<sub>2</sub> from sulphur recovery.

	1990 <sup>1</sup>	1995	2000	2005 <sup>2</sup>	2010 <sup>2</sup>	2011 <sup>2</sup>	2012 <sup>2</sup>	2013
NMVOC, Mg	4072	5500	4530	3742	5477	5479	5543	5575
SO <sub>2</sub> , oil refining, Mg		585	178					
SO <sub>2</sub> , sulphur recovery, Mg	3335	2437	803	347	981	1150	881	517

<sup>1</sup> Prior to 1994 SO<sub>2</sub> emissions from oil refining and sulphur recovery are reported as area sources in category 1B2a vi.

<sup>2</sup> From 2001 SO<sub>2</sub> emissions from oil refining are included in stationary combustion.

### Service stations

Emissions from service stations are calculated using the emission factors in Table 3.4.8 and the sales of gasoline given by the Danish Energy statistics. The NMVOC emissions are listed in Table 3.4.15.

Table 3.4.15 Emissions of NMVOC from service stations for selected years of the time series.

	1990	1995	2000	2005	2010	2011	2012	2013
NMVOC, Mg	4 856	2 965	2 616	1 031	851	796	754	725

### Transmission and distribution of gas

The gas transmission company reports emissions of CH<sub>4</sub> for the years 1999 and onwards. Calculations of the CH<sub>4</sub> emissions for transmission are based on registered loss in the transmission grid and the emission from the natural gas consumption in the pressure regulating stations. The distribution companies give data on fugitive losses, and the CH<sub>4</sub> emissions are estimated due to the gas quality given by Energinet.dk. For the years 1991-1998 the CH<sub>4</sub> emissions for transmission are estimated on the basis of registered loss provided by the transmission company and the annual composition of Danish natural gas given by Energinet.dk. Transmission loss is not available for 1990, why the average for 1991-1995 is applied. Calculations of the NMVOC emissions are based on the CH<sub>4</sub> emission and the gas quality measured by Energinet.dk (Table 3.4.9) according to equation 3.4.6.

$$E_{NMVOC} = E_{CH_4} \times (w_{NMVOC} / w_{CH_4}) \quad (\text{eq. 3.4.6})$$

where  $w_{NMVOC}$  is the weight-% NMVOC and  $w_{CH_4}$  is the weight-%  $CH_4$  according to the gas quality of the current year.

As the pipelines in Denmark are relatively new and made of plastic, most emissions are due to construction and maintenance. The decrease of emissions from transmission in 2007 (Table 3.4.16) is caused by the completion of a greater construction work and rerouting of a major pipeline. In preparation for construction work on a new compressor station, there has been laid a number of new line valve stations in 2011. Before this work could be done, larger amounts of natural gas were vented to drain the pipes. Therefore emissions from transmission of natural gas are significantly high in 2011.

The distribution companies provide emissions of  $CH_4$  for the years 1997 and onwards, which are used to calculate NMVOC emissions. For the years 1995-1996 NMVOC emissions are calculated from the registered loss from distribution and the annual composition of Danish natural gas given by Energinet.dk. As distribution losses are not available for the years 1990-1994, the percentage loss for 1995 is used. Emissions from distribution of gas mainly owe to excavations and maintenance of the pipelines, but also difference between the calendar year and the meter reading year might influence the annual variations.

Data on distribution of town gas is limited as more companies are closed. For the active companies, data are available for part of the time series only. Gap filling and extrapolation of distribution and loss amounts to early years are done in collaboration with the individual companies to the extent possible. As the town gas distribution network is significant older the gas losses and thus the emissions are larger than for the natural gas distribution network, even though the distribution rates for natural gas far exceeds the rates for town gas. Emissions from distribution of town gas show large decreases in 2003 and 2005 due to the close down of two distribution companies and following no town gas distribution in those areas.

Emissions of NMVOC from transmission of natural gas (including storage) and distribution of natural gas and town gas are shown in Table 3.4.16.

Table 3.4.16 NMVOC emission from transmission, storage of natural gas and distribution of natural gas and town gas for selected years of the time series.

NMVOC emission, Mg	1990	1995	2000	2005	2010	2011	2012	2013
Transmission	41	135	26	36	6	43	4	4
Distribution, natural gas	12	22	15	17	9	12	18	14
Distribution, town gas	683	700	680	587	53	50	39	30

## Venting and flaring

### Venting

Venting is limited to the gas storage plants and the emissions are of minor importance. Emissions of  $CH_4$  and NMVOC are given in the environmental reports for the gas storage plants (DONG Energy, 2014a; Energinet.dk, 2014a). The time series are shown in Figure 3.4.8.

### Flaring in refineries

Flaring in refineries is a significant fugitive emission source to emissions of  $SO_2$ . In 1990-1993 emissions from petroleum product processing were in-

cluded in emissions from flaring in refineries (NFR category 1B2c). From 1994 the data delivery format was changed, which made it possible to split the emissions into contributions from flaring and processing, respectively. Emissions from processing are from 1994 included in NFR category 1B2a iv.

Emissions for selected years and components are shown in Table 3.4.17. One refinery was shut down in 1996 leading to larger emissions in 1990-1996. Technical improvements of the sulphur recovery system at one of the two Danish refineries lead to a decrease of SO<sub>2</sub> emissions from 1996-1998. The large emissions from 2005 and onwards owe to shut-downs due to maintenance and accidents. Further, construction and initialisation of new facilities and problems related to the ammonium thiosulphate (ATS) plant at the one refinery has led to increased emissions. In 2007 the capacity of the ATS plant was increased followed by commissioning difficulties.

Table 3.4.17 Emissions from flaring in refineries.

Year	1990*	1995	2000	2005	2010	2011	2012	2013
	Mg	Mg	Mg	Mg	Mg	Mg	Mg	Mg
SO <sub>2</sub> <sup>1</sup>	943	203	51	296	326	271	175	166
NO <sub>x</sub>	13	13	11	26	19	18	23	15
NMVOG	31	31	26	32	27	25	32	22
CO	73	73	60	73	62	59	74	52

<sup>1</sup> In 1990-1993 emissions from petroleum product processing were included in flaring in refineries due to the data delivery form. From 1994 emissions from petroleum product processing were given in 1B2a iv.

### Flaring offshore

The time series for the emission of NMVOC from offshore flaring fluctuates due to the fluctuations in the fuel rate (Figure 3.4.12). As shown in Figure 3.4.10 there were marked peaks in the amount of offshore flaring in 1997 and, especially, in 1999. The increase in 1997 was due to the new Dan field and the completion of the Harald field. The increase in 1999 was due to the opening of three new fields; Halfdan, Siri and Syd Arne.




Figure 3.4.12 NMVOC emissions from venting and flaring of gas.

The emissions from offshore flaring are estimated from the same set of emission factors for all years in the time series and the variations reflect only the variations in the flared amounts. Emissions of selected components from flaring in oil and gas extraction including offshore flaring are shown in Table 3.4.18. Flaring in gas storage and gas transmission and distribution only

contribute minor to the emissions from flaring, and are not included in Table 3.4.18.

Table 3.4.18 Emissions from flaring in oil and gas extraction.

Year	1990	1995	2000	2005	2010	2011	2012	2013
	Mg	Mg	Mg	Mg	Mg	Mg	Mg	Mg
SO <sub>2</sub>	1	2	3	2	2	1	1	1
NO <sub>x</sub>	133	186	307	226	146	95	87	106
NM VOC	160	225	371	273	177	115	105	129
CO	201	281	464	342	221	144	132	161

### 3.4.6 Uncertainties and time series consistency

#### Methodology

The applied methodology for uncertainty estimates refers to Pulles & Aardenne (2004). The Danish uncertainty estimates are based on the simple Tier 1 approach described in IPCC Good Practice Guidance (IPCC, 2000).

The uncertainty estimates are based on the calculated emissions for the base year and for the latest inventory year and on the uncertainty rates for both activity data and emission factors. Data is aggregated for the NFR category 1 B - Fugitive Emissions from Fuels. Base year refers to 2000 for particulate matter and to 1990 for the remaining pollutants.

The uncertainty rates are based on the EMEP/EEA emission inventory guidebook (2013), on uncertainty estimates from companies and on estimates and assumptions by DCE. The applied uncertainty levels for activity data and emission factors are given in Table 3.4.19.

Table 3.4.19 Uncertainty levels for activity data and emission factors.

Pollutant	Activity Data	Emission Factor
	Uncertainty level, %	Uncertainty level, %
SO <sub>2</sub>	10	25
NO <sub>x</sub>	8	125
NM VOC	2	125
CO	8	125
TSP	2	50
PM <sub>10</sub>	2	50
PM <sub>2.5</sub>	2	50
BC	2	100
As	8	500
Cd	8	500
Cr	8	500
Cu	8	500
Hg	8	500
Ni	8	500
Pb	8	500
Se	8	500
Zn	8	500
Benzo(b)	8	500
Benzo(k)	8	500
Benzo(a)	8	500
Indeno	8	500



## Results

The uncertainty model estimates uncertainties for both the emission level and the trend. The uncertainty on the emission level for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC and CO is 27 %, 125 %, 125 % and 125 %, respectively.

For PM the uncertainty is 50 %, for BC the uncertainty is 100 % and for HM and PAHs the uncertainty is 500 %. The individual uncertainty estimates for the fugitive emission inventory are shown in Table 3.4.20.

Table 3.4.20 Estimated emission uncertainty and trend uncertainty for fugitive emissions. The trend refers to the years 1990-2013 for all pollutants except PM where the trend refers to 2000-2013.

Pollutant	Emission uncertainty	Trend uncertainty
	%	%
SO <sub>2</sub>	27	2
NO <sub>x</sub>	125	9
NMVOC	125	2
CO	125	8
TSP	50	2
PM <sub>10</sub>	50	2
PM <sub>2.5</sub>	50	2
BC	100	2
As	500	8
Cd	500	8
Cr	500	8
Cu	500	8
Hg	500	8
Ni	500	8
Pb	500	8
Se	500	8
Zn	500	8
Benzo(b)	500	8
Benzo(k)	500	8
Benzo(a)	500	8
Indeno	500	8

### 3.4.7 Source specific QA/QC and verification

A list of QA/QC tasks are performed directly in relation to the fugitive emission part of the Danish emission inventories. The following procedures are carried out to ensure the data quality:

- The emission from the large point sources (refineries, gas treatment and gas storage plants) is compared with the emission reported the previous year.
- Annual environmental reports are kept for subsequent control of plant-specific emission data.
- Checks of data transfer are incorporated in the fugitive emission models, e.g. sum checks.
- Verification of activity data from external data when data are available through more data sources (offshore fuel and flaring rates).
- Data sources are incorporated in the fugitive emission models
- A manual log table in the emission databases is applied to collect information about recalculations.

- Comparison with the inventory of the previous year. Any major changes are verified.
- Total emission, when aggregated to reporting tables, is compared with totals based on SNAP source categories (control of data transfer).
- Checking of time series in the NFR and SNAP source categories. Significant dips and jumps are controlled and explained.

The QC work will continue in future years.

#### Data deliveries

Table 3.4.21 lists the external data deliveries used for the inventory of fugitive emissions. Further the table holds information on the contacts at the data delivery companies.

Table 3.4.21 List of external data sources.

Category	Data description	Activity data, emission factors or emissions	Reference	Contact(s)	Data agreement/ Comment
Exploration of oil and gas	Dataset for extraction of oil and gas, including amounts and composition.	Activity data	The Danish Energy Agency	Jan H. Andersen	Data agreement
Production of oil and gas	Gas and oil production. Dataset for production of oil, gas and number of platforms. Amounts of offshore loading of ships	Activity data	The Danish Energy Agency	Jan H. Andersen	Not necessary due to obligation by law
Offshore flaring	Flaring offshore in oil and gas extraction (ETS data)	Activity data	The Danish Energy Agency	Dorte Maimann	Data agreement
Service stations	Data on gasoline sales from the Danish energy statistics.	Activity data	The Danish Energy Agency	Jane Rusberg	Data agreement
Gas transmission	Natural gas from the transmission company, sales and losses (meter differences)	Activity data	Energinet.dk	Christian Friberg B. Nielsen	Not necessary due to obligation by law
Onshore activities	Amounts of oil transport in pipeline and onshore loading to ships. Emissions from storage of raw oil in the terminal.	Activity data and emission data	DONG Olierør A/S	Stine B. Bergmann	No formal data agreement.
Gas distribution	Natural gas from the distribution company, sales and losses (meter differences)	Activity data	Naturgas Fyn, HMN Aalborg Forsyning	Hanne Mochau, Søren K. Andersen, Andreas Bech Jensen	No formal data agreement.
Air emissions from refinery	Fuel consumption and emission data.	Activity data and emission data	Statoil A/S, A/S Danish Shell	Anette Holst, Lis Rønnow Rasmussen	No formal data agreement.
Storage and treatment of gas	Environmental reports from plants defined as large point sources (Lille Torup, Stenlille, Nybro)	Activity data	Various plants		Not necessary due to obligation by law
Emission factors	Emission factors origin from a large number of sources	Emission factors	See chapter regarding emission factors		

### 3.4.8 Source specific recalculations

The following recalculations regarding fugitive emissions from fuels have been applied for the time series:

#### Exploration of oil and gas

Exploration has been included as a new source in the emission inventory as activity data has now been made available by the Danish Energy Agency. Emissions only occur in years with exploration and appraisal wells (E/A wells) (1990-2000, 2002, 2005, 2009 and 2013). The largest E/A productions occurred in 1990 and 2002, contributing 2.4 % and 3.7 % to the fugitive NO<sub>x</sub> emission, respectively, while the corresponding shares for SO<sub>2</sub> and NMVOC are ≤ 0.1 %.

#### Refineries

In cooperation with one refinery, the SO<sub>2</sub> emissions have been examined. This work has led to a reallocation of part of the SO<sub>2</sub> emissions in the years 2005-2012. SO<sub>2</sub> emissions from the ammoniumthiosulfat (ATS) unit have been reallocated to the flaring category for the years 2008-2012, as the refinery has informed that H<sub>2</sub>S from the ATS unit is being led to the flare. In previous inventory these emissions were allocated to the category refining/storage.

For 2012 the SO<sub>2</sub> emission has been updated according to supplementing information from one refinery, leading to a decrease of 8.1 ktonnes SO<sub>2</sub>. Also a small correction of the fuel gas consumption has been applied according to an update in the annual environmental report.

The methodology for estimating NMVOC from one refinery has been changed, resulting in an increase of the NMVOC emissions for the years 1994-2012. The refinery report annual VOC emissions based on results from measurement campaigns, the last in carried out in 2006. In previous inventories, the split between NMVOC and CH<sub>4</sub> given by the refinery has been used. This methodology has been changed, as the CH<sub>4</sub> share was much higher than for the other Danish refinery, and also much higher than corresponding shares found in a literature study. The CH<sub>4</sub> share has therefore been changed from 56 % to 10 % as given by the other Danish refinery and supported by shares of 10-20 % found for Swedish refineries. The largest increase of the NMVOC emissions are estimated for the years 2006-2012 (1611 tonnes per year), corresponding 10 % - 16 % of the total fugitive NMVOC emissions (16 % in 2011 and 2012).

#### Service stations

The activity data has been updated for 2011-2012 according to the latest energy statistics published by the Danish Energy Agency. Also an error for the emission factors has been corrected for the years 1992-2012. The recalculation has changed the NMVOC emission by -0.711 ktonnes (2005) to +0.738 ktonnes (1992), which is insignificant compared to the total fugitive NMVOC emission (< 0.01%).

#### Natural gas transmission and distribution

Activity data has been updated for one town gas distribution company for the year 2012. The change of 0.27 tonnes NMVOC has insignificant impact on the total fugitive emissions (< 0.01%).

### **Venting**

Activity data has been corrected for one gas storage plant for 2012 according to the annual environmental report. The change is insignificant for the total fugitive emissions (< 0.01%).

### **Flaring in refineries**

The NO<sub>x</sub> emission factor has been updated for the years 1994-1996 according to the 2013 EMEP/EEA Guidebook. The changes of the emissions are largest in 1994 with an increase of 2 ktonnes NO<sub>x</sub>, corresponding 0.4 % of the total fugitive emissions.

### **Flaring in gas storage and treatment plants**

NO<sub>x</sub> emissions are updated according to the environmental report for the gas treatment plant for 2012. The decrease of the NO<sub>x</sub> emission is due to installation of new low-NO<sub>x</sub> burner. NO<sub>x</sub> has been updated from 45 tonnes to 0.007 tonnes. The decrease of the NO<sub>x</sub> emission account for 0.02 % of the total fugitive NO<sub>x</sub> emission.

### **Flaring in gas transmission and distribution**

Flaring in gas transmission and distribution has been included as a new source in the emission inventory, only occurring in the years 2011-2013. The gas transmission company inform that they have started using a mobile flare in large construction works, and also one distribution company is flaring gas. The change of 0.4 tonnes NMVOC has insignificant impact on the total fugitive emissions (< 0.01%).

## **3.4.9 Source specific planned improvements**

The following future improvements are suggested.

Emissions from storage of fuels in tank facilities: The recent edition of the Danish emission inventory holds emissions from extraction of fuels, combustion of fuels and from service stations. To make the inventory complete emissions from storage of fuels in tank facilities should be included in the future if data is available. Work is going on to locate greater tank facilities in Denmark and collect the available data. In cases where no emission estimates or measurements are available a set of emission factors have to be set up.

## **3.4.10 References**

A/S Dansk Shell, 2014: Annual environmental report 2012.

Andersen, 2014: Personal communication, -mail correspondence, Danish Energy Agency.

Danish Energy Agency, 2014a: Oil and Gas Production in Denmark 2013. Available at:

[http://www.ens.dk/sites/ens.dk/files/energistyrelsen/Nyheder/2014/oil\\_and\\_gas\\_production\\_in\\_denmark\\_2013\\_uk.pdf](http://www.ens.dk/sites/ens.dk/files/energistyrelsen/Nyheder/2014/oil_and_gas_production_in_denmark_2013_uk.pdf) (29-01-2015).

Danish Energy Agency, 2014b: The Danish energy statistics (Energistatistik) (in Danish). Available at:

<http://www.ens.dk/info/tal-kort/statistik-noglestal/arlig-energistatistik> (29-01-2015).

DEPA, 2008: Emissionsfaktorer for NO<sub>x</sub>-emissioner fra flaring fra platforme i Nordsøen (in Danish). Not published.

Danish Ministry of the Environment, 2011: BEK nr 1158 af 07/12/2011. Legal Act (in Danish). Available at:  
<https://www.retsinformation.dk/Forms/R0710.aspx?id=139278&exp=1>  
(13-01-2015)

DONG Energy, 2014a: Annual environmental reports from DONG Energy for Stenlille gas storage plants (in Danish). Available at:  
<http://www.dongenergy.com/da/ansvarlighed/resultater/gr%C3%B8nne-regnskaber> (29-01-2015).

DONG Energy, 2014b: Annual environmental reports from DONG Energy for Nybro gas treatment plant (in Danish). Available at:  
<http://www.dongenergy.com/da/ansvarlighed/resultater/gr%C3%B8nne-regnskaber> (29-01-2015).

DONG Oil Pipe A/S, 2014: Dong Energy Raw Oil Terminal, Fredericia – Self-regulation Report 2013 (in Danish).

EMEP/EEA, 2013: Air emission inventory guidebook, prepared by the UNECE/EMEP Task Force on Emission Inventories and Projections, 2013 update. Available on the internet at:  
<http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>  
(17-01-2014).

Energinet.dk, 2014: Annual report from Energinet.dk for Lille Torup gas storage plant (in Danish). Available on the internet at:  
<http://gaslager.energinet.dk/EN/about/Pages/The-Company.aspx>  
(29-01-2015).

Fenhann, J. & Kilde, N.A. 1994: Inventory of Emissions to the Air from Danish Sources 1972-1992. Risø National Laboratory, Roskilde, Denmark.

Hjerrild & Rasmussen, 2014: Fugitive VOC from refineries.

IPCC, 2000: Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/gp/english/> (17-01-2014).

IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/>

Jensen, 2008: Town gas distribution loss.

Karll, B. 2002: Personal communication, e-mail 2002, Danish Gas Technology Centre.

Karll, B. 2003: Personal communication, e-mail 2003, Danish Gas Technology Centre.

Karll, B. 2004: Personal communication, e-mail 2004, Danish Gas Technology Centre.

Karll, B. 2005: Personal communication, e-mail 2005, Danish Gas Technology Centre.

Miljøcenter Odense, 2012: Revurdering af miljøgodkendelse for A/S Dansk Shell, Havneterminalen (in Danish) Available at: [http://mst.dk/media/mst/Attachments/Revurdering\\_Miljoegodkendelse\\_KLHOU\\_21mar12.pdf](http://mst.dk/media/mst/Attachments/Revurdering_Miljoegodkendelse_KLHOU_21mar12.pdf) (29-01-2015).

Oertenblad, M. 2006: Personal communication, e-mail 2006, Danish Gas Technology Centre.

Oertenblad, M. 2007: Personal communication, e-mail 2007, Danish Gas Technology Centre.

Pulles, T. & Aardenne, J.v. 2004: Good Practice for CLRTAP Emission Inventories, 24. June 2004. Available at: <http://www.eea.europa.eu> (29-01-2015).

Statoil A/S, 2014: Annual environmental report 2013 (in Danish).

Visschedijk, A., Pacyna, J., Pulles, T., Zandveld, P. & van der Gon, H.D. 2004: Coordinated European Particulate Matter Emission Inventory Program (CEPMEIP).

## 4 Industrial processes (NFR sector 2)

### 4.1 Overview of the sector

The chapter on *Industrial processes* (NFR sector 2) is outlined as follows:

- Mineral Products (NFR 2A)
- Chemical Industry (NFR 2B)
- Metal Production (NFR 2C)
- Non-energy products from fuels and solvent use (NFR 2D)
- Other Product Use (NFR 2G)
- Other Production (NFR 2H)
- Other production, consumption, storage, transportation or handling of bulk products (NFR 2L)

The industrial processes included in the Danish inventory are those in large companies, e.g. cement factories, as well as a number of smaller companies e.g. iron foundries.

Table 4.1.1 presents an overview of sources and groups of pollutants included in the present reporting. Explanations to the abbreviations are given below the table. In addition to the indicated groups of pollutants some groups do not include all relevant pollutants or the time series are not complete. For some processes, it is not possible to separate emissions from the fuels and the emissions stemming from the raw materials. This is especially the case for processes with contact, e.g. cement and lime production. Detailed information on this subject can be found in the following table.

Table 4.1 .1 Survey of IPPU sector with SNAP-code and NFR-code included in the Danish inventory.

Industrial sector	SNAP	NFR	SO <sub>2</sub> /NO <sub>x</sub> / NH <sub>3</sub>	NMVOC/ CO	PMs	HMs	POPs
Cement production	030311	2A1	IE	IE	IE	IE	IE
Lime production	030312	2A2	IE	IE	x	-	x
Container glass production	030315	2A3	-	-	x	x	-
Glass wool production	030316	2A3	x	x	x	-	-
Production of yellow bricks	040691	2A6	x	-	-	-	x
Production of expanded clay products	040692	2A6	x	-	-	-	x
Mineral wool production	030318	2A6	x	x	x	-	x
Sulphuric acid production	040401	2B10a	x	-	-	-	-
Nitric acid production	040402	2B2	x	-	x	-	-
Catalyst production	040416	2B10a	x	-	x	-	-
Production of chemical ingredients	040500	2B10a	-	x	-	-	-
Pesticide production	040525	2B10a	-	x	-	-	-
Production of tar products	040527	2B10a	x	x	-	x	-
Electric arc furnace steel production	040207	2C1	IE	IE	x	x	x
Rolling mills steel production	040208	2C1	IE	IE	x	x	-
Grey iron foundries	030303	2C1	IE	IE	x	x	x
Allied metal manufacturing	040306	2C2	IE	IE	-	x	-
Secondary aluminium production	030310	2C3	IE	IE	x	x	x
Secondary lead production	030307	2C5	IE	IE	x	x	x
Domestic solvent use incl. fungicides	060408/ 060411	2D3a	-	x	-	-	-
Road paving with asphalt	040611	2D3b	-	x	-	-	-
Asphalt roofing	040610	2D3c	-	x	-	-	-
Coating applications	060100	2D3d	-	x	-	-	-
Degreasing	040200	2D3e	-	x	-	-	-
Dry cleaning	040202	2D3f	-	x	-	-	-
Chemical products	060300	2D3g	-	x	-	-	-
Printing	060403	2D3h	-	x	-	-	-
Other solvent use	060400	2D3i	-	x	-	-	-
Other product use: Fireworks	060601	2G	x	x	x	x	-
Other product use: Tobacco	060602	2G	x	x	x	x	x
Other product use: Charcoal	060605	2G	x	x	x	x	x
Other product use: Paraffin wax use	060606	2G	-	x	x	-	x
Bread	040605	2H2	-	x	-	-	-
Wine	040606	2H2	-	x	-	-	-
Beer	040607	2H2	-	x	-	-	-
Spirits	040608	2H2	-	x	-	-	-
Sugar production	040625	2H2	-	x	-	-	-
Meat curing	040627	2H2	-	x	-	-	-
Margarine and solid cooking fats	040698	2H2	-	x	-	-	-
Coffee roasting	040699	2H2	-	x	-	-	-
Other (slaughterhouse waste)	040617	2L	x	-	-	-	-

x Included in the present inventory.

- Not included/not relevant.

IE Included elsewhere.

Table 4.1.2 presents an overview of the most significant source categories for 2013. Many changes have occurred over the time series; some factories have closed and others have opened, Table 4.1.2 is therefore only representable for 2013.



Table 4.1.2 Overview of 2013 emissions from Industrial processes and product use (IPPU)

	Total emission from IPPU		Fraction of national total, %	Largest contributor in IPPU	Emission from largest contributor		Fraction of IPPU, %
SO <sub>2</sub>	1.98	Gg	14.5	2A6 Other mineral products	1.75	Gg	88.2
NO <sub>x</sub>	0.08	Gg	0.1	2G Other product use	0.06	Gg	74.1
NMVOC	32.24	Gg	28.2	2D3i Other solvent use	21.30	Gg	66.1
CO	4.00	Gg	1.2	2G Other product use	3.74	Gg	93.4
NH <sub>3</sub>	0.42	Gg	0.6	2A6 Other mineral products	0.18	Gg	43.2
TSP	0.68	Gg	0.8	2G Other product use	0.37	Gg	55.1
HMs	6.55	Mg	5.5	Cu from 2G Other product use	1.98	Mg	30.2
POPs	0.10	Mg	1.4	PAHs from 2G Other product use	0.10	Mg	99.8

## 4.2 Mineral products

### 4.2.1 Source category description

The sub-sector *Mineral products* (NFR 2A) cover the following processes relevant for the Danish inventories:

- 2A1 Cement production (SNAP 030311).
- 2A2 Lime production (SNAP 030312).
- 2A3 Glass production (SNAP 030315, 030316)
- 2A6 Other mineral products (SNAP 030318, 040691, 040692)

The time series for emission of acidifying substances, NMVOC, particulate matter, heavy metals, and POPs from *Mineral products* (NFR 2A) is available in the NFR tables. Table 4.2.1 presents an overview of emissions from 2013.

Table 4.2.1 Overview of 2013 emissions from Mineral products.

	Total emission from Mineral industries		Fraction of IPPU; %	Largest contributor in Mineral industries	Emission from largest contributor		Fraction of Mineral industries, %
SO <sub>2</sub>	1.75	Gg	88.2	2A6 Other mineral products	1.75	Gg	100.0
NMVOC	0.04	Gg	0.1	2A3 Glass production	0.04	Gg	100.0
CO	0.01	Gg	0.3	2A6 Other mineral products	0.01	Gg	87.3
NH <sub>3</sub>	0.30	Gg	71.4	2A6 Other mineral products	0.18	Gg	60.6
TSP	0.14	Gg	19.9	2A6 Other mineral products	0.06	Gg	45.9
HMs	0.05	Mg	0.8	Pb from 2A3 Glass production	0.02	Mg	44.2
POPs	10.6	kg	4.8	PCB from 2A2 Lime production	10.0	kg	94.9

### 4.2.2 Cement production

It has not been possible to separate emissions from fuel consumption and emissions from process activities. Process emissions from the production of cement are therefore included in the energy section.

### 4.2.3 Lime production

The production of limestone and lime/burned lime/quicklime is located at a few localities: Faxe Kalk (Lhoist group) situated in Faxe, Scandinavian Calcium Oxide ApS situated in Støvring, dankalk A/S situated in Løgstør with limestone quarries/limeworks in Aggersund, Mjels, Poulstrup and Batum. The following SNAP-codes are covered:

- 03 03 12 Lime production

The following pollutants are relevant for the lime production process:

- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>
- BC
- HCB
- PCDD/F
- PCB

Emissions associated with the fuel use are estimated and reported in the energy sector.

### Methodology

#### Activity data

The activity data regarding production of lime is obtained from Statistics Denmark (2014). The data are presented in Table 4.2.2.

Table 4.2.2 Production of burnt lime, Mg (Statistics Denmark, 2014).

	1990	1991	1992	1993	1994	1995	1996	1997
Burnt lime	127978	86222	104526	106587	112480	100789	95028	102587
	1998	1999	2000	2001	2002	2003	2004	2005
Burnt lime	88922	95177	92002	96486	122641	87549	77844	71239
	2006	2007	2008	2009	2010	2011	2012	2013
Burnt lime	78652	75504	74981	46202	50397	59430	69136	66805

Slaking of lime does not emit any pollutants. All burnt lime that is later slaked, is included in statistics shown in the table above. Adding the production of slaked lime to the activity data, would therefore result in a double counting.

#### Emission factors

The emission factors used to calculate the emissions from lime production are shown in Table 4.2.3 along with their respective sources.

Table 4.2.3 Emission factors for production of lime

Pollutant	Unit	Value	Source
TSP	kg/Mg	0.5	CEPMEIP
PM <sub>10</sub>	kg/Mg	0.20	CEPMEIP
PM <sub>2.5</sub>	kg/Mg	0.04	CEPMEIP
BC	g/Mg	0.18	EMEP/EEA Guidebook (2013)
HCB	mg/Mg	0.01	Nielsen et al. (2013)
PCDD/F	µg/Mg	0.02	Henriksen et al. (2006)
PCB	mg/Mg	0.15	Nielsen et al. (2013)

#### Emission trends

The emission trends for particles and POPs for lime production are shown in Table 4.2.4 and in Figure 4.2.1. Emission data for the different particle size distributions and individual POPs are available in the NFR tables.

Table 4.2.4 Emission of particles and POPs.

	Unit	1990	1991	1992	1993	1994	1995	1996	1997
TSP		-	-	-	-	-	-	-	-
POPs	kg	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
		1998	1999	2000	2001	2002	2003	2004	2005
TSP	Gg	-	-	0.05	0.05	0.07	0.05	0.05	0.04
POPs	kg	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01
		2006	2007	2008	2009	2010	2011	2012	2013
TSP	Gg	0.05	0.04	0.04	0.03	0.03	0.02	0.03	0.03
POPs	kg	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01




Figure 4.2.1 Emission trends for particles and POPs from lime production.

There is a peak in emissions in 2002 due to a corresponding peak for the activity data. The activity data are based on the official statistics from Statistics Denmark and there is no immediate explanation for this peak. There are very few producers in Denmark and therefore it will not be possible to obtain more detailed data from Statistics Denmark.

#### 4.2.4 Glass production

Glass production covers production of:

- Flat glass
- Container glass
- Glass wool

The production of flat glass (SNAP 03 03 14 Flat glass) is concentrated at few European producers and none of these have plants in Denmark. The processes in Denmark are limited to mounting of sealed glazing units. The mounting process is not considered to contribute to emission of pollutants to air in Denmark.

The production of container glass for packaging is concentrated at one company: Ardagh Glass Holmegaard A/S (previously Rexam Glass Holmegaard A/S) and for art industrial glass products: Holmegaard A/S both situated in Fensmark, Næstved. Saint-Gobain Isover situated in Vamdrup produces glass wool. The following SNAP-codes are covered:

- 03 03 15 Container glass
- 03 03 16 Glass wool

The following pollutants are relevant for the glass production process:

- NMVOC
- CO
- NH<sub>3</sub>
- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>
- Heavy metals: As, Cd, Cr, Ni, Pb, Se, Zn

### Methodology

#### Activity data

Activity data for the production of container glass and glass wool are presented in Table 4.2.5 and Figure 4.2.2.

Table 4.2.5 Production of container glass and glass wool, Gg.

	1990	1991	1992	1993	1994	1995	1996	1997
Container glass	164	159	145	140.5	150.2	140	140	140
Glass wool	36	36	36	36	36	36	36	35
	1998	1999	2000	2001	2002	2003	2004	2005
Container glass	187	198	180	179	161	145	139	126
Glass wool	34	39	40	37	35	37	41	37
	2006	2007	2008	2009	2010	2011	2012	2013
Container glass	48	52	130	79	86	88	90	68
Glass wool	43	41	41	33	25	30	27	28




Figure 4.2.2 Activity data for container glass and glass wool production.

The annual produced amount of container glass is estimated based on the consumption of raw materials. Data on raw materials are gathered from environmental reports (1990-2005) and EU-ETS data (2006-2013).

The produced amount of glass wool is available in the company's environmental reports (1997-2013). Production data back to 1990 are estimated as the constant average of 1997-1999.

The drastic decrease in the calculated production of container glass in 2006 and 2007 is a result of a similar decrease in the consumption of limestone reported to the EU-ETS database. It will be investigated if the temporary decrease is due to EU-ETS teething troubles or something else.

Both the container glass and glass wool production displays a significant decrease from 2008 to 2010 that can be explained by the financial crises.

### Emission factors

Yearly measurements of the emissions from production of container glass provide emissions of TSP (2000-2013), lead (1997-2013), selenium (1997-2013) and zinc (1997-2001) (Ardagh Glass Holmegaard, 2014). PM<sub>10</sub> and PM<sub>2.5</sub> are estimated from the distribution between TSP, PM<sub>10</sub> and PM<sub>2.5</sub> (1/0.9/0.8) from the EMEP/EEA Guidebook (2013). Emissions of arsenic, cadmium, chromium and nickel are estimated from standard emission factors, where direct emissions are not available for lead, selenium and zinc; these are also calculated using emission factors. All used emission factors are shown in Table 4.2.6. From 2006, measured particle emissions from the singular Danish container glass producer decrease 90 % due to installation of abatement equipment; all calculated heavy metal emissions are therefore also lowered with 90 % from 2006.

The emission of NH<sub>3</sub> and TSP from the production of glass wool has been measured yearly since 1996 and are available in the company's environmental reports (Saint-Gobain Isover, 2014). PM<sub>10</sub> and PM<sub>2.5</sub> are estimated from the distribution between TSP, PM<sub>10</sub> and PM<sub>2.5</sub> (1/0.9/0.8) from the EMEP/EEA Guidebook (2013). NMVOC and CO have also been measured for 2007-2013 and 1996-1997 respectively. For the years where no measured emission data are available, emissions are calculated using implied emission factors (IEF) based on the available measurements. All used emission factors are shown in Table 4.2.6. Since it has not been possible to separate process emissions from the emissions from fuel combustion, the measured/calculated emissions from glass wool production presented here account for the entire production.

Table 4.2.6 Emission factors for production of glass production.

Pollutant	Container glass production			Glass wool production		
	Unit	Value	Source	Unit	Value	Source
NMVOC		-		kg/Mg	1.4	IEF (2007-2009)
CO		-		kg/Mg	0.1	IEF (1996-1997)
NH <sub>3</sub>		-		kg/Mg	7.4	IEF(1996-1998)
TSP	kg/Mg	0.28	Guidebook (2013)	kg/Mg	2.0	IEF (2000-2013)
PM <sub>10</sub>	kg/Mg	0.25	Guidebook (2013)	kg/Mg	1.8	IEF (2000-2013)
PM <sub>2.5</sub>	kg/Mg	0.22	Guidebook (2013)	kg/Mg	1.6	IEF (2000-2013)
As	g/Mg	0.29	Guidebook (2013)		-	
Cd	g/Mg	0.12	Guidebook (2013)		-	
Cr	g/Mg	0.37	Guidebook (2013)		-	
Ni	g/Mg	0.24	Guidebook (2013)		-	
Pb	g/Mg	2.90	Guidebook (2013)		-	
Se	g/Mg	1.50	Guidebook (2013)		-	
Zn	g/Mg	0.23	Guidebook (2013)		-	

### Emission trends

The only pollutants to which both container glass and glass wool productions contribute are particles. Therefore, only particle emissions are presented in the table below. Table 4.2.7 shows the individual emissions from the two sources.

Table 4.2.7 Emission of particles (TSP) from glass production

	2000	2001	2002	2003	2004	2005	2006
Container glass	26.0	25.0	21.0	26.0	23.0	7.0	0.9
Glass wool	111.0	119.0	114.0	102.0	99.0	85.0	82.0
	2007	2008	2009	2010	2011	2012	2013
Container glass	1.0	1.2	1.2	1.7	1.8	4.5	1.6
Glass wool	52.0	54.0	33.0	26.0	42.0	47.0	38.0

Emissions of the remaining pollutants can be found in the NFR tables, where NMVOC, CO and NH<sub>3</sub> emissions stem only from glass wool production and heavy metal emissions only from container glass production.

#### 4.2.5 Other mineral products

The sub-sector "Other" covers production of bricks and tiles (aggregates or bricks/blocks for construction), expanded clay products for different purposes (aggregates as absorbent for chemicals, cat litter, and for other miscellaneous purposes) and mineral wool from the company Rockwool situated at three localities in Denmark: Hedehusene<sup>1</sup>, Vamdrup and Øster Doense produces stone wool. The following SNAP-codes are covered:

- 04 06 91 Production of yellow bricks
- 04 06 92 Expanded clay products
- 03 03 18 Mineral wool

The following pollutants are covered:

- SO<sub>2</sub>
- NMVOC
- CO
- NH<sub>3</sub>
- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>

#### Methodology

The production of yellow bricks is found all over the country, where clay is available. Producers of expanded clay products are located in the northern part of Jutland. The SO<sub>2</sub> emission from the production of bricks and expanded clay products is related to sulphur content in the raw material. No other process emissions are calculated from these source categories.

Mineral wool is produced from mineral fibres and a binder. The raw materials are melted in a cupola fired by coke and natural gas. The consumption of raw material as well as amount of produced mineral wool is confidential. Information on emissions from some years has in combination with yearly raw material consumption been used to extrapolate the emissions to other years. The data have been extracted from company reports (Rockwool, 2014) and reports to PRTR. Measured emissions of CO and NH<sub>3</sub> are available for the years 2001, 2004 and 2007-2013 and emissions of particulate matter are available for 1995-2013. For PCDD/F, the inventory is based on measured emissions for 2004.

<sup>1</sup> The melting of minerals (cupola) has been closed down in 2002.

### Activity data

National statistics on bricks, tiles and expanded clay contain a broad range of different products, most of them in units of numbers (no.). The consumption of limestone is therefore used as alternative activity data for these source categories.

Data on the produced amount of stone wool is confidential; however the consumption of carbonates at the three Danish Rockwool factories is available from the annual environmental reports. The different raw materials such as lime, waste, bottom ash etc. are added up to the activity data of CaCO<sub>3</sub> equivalents.

The chosen activity data for “Other mineral products” are shown in Table 4.2.8 and Figure 4.2.3.

Table 4.2.8 Production of “Other mineral products”, Mg CaCO<sub>3</sub> equivalents.

	1990	1991	1992	1993	1994	1995	1996	1997
Brickworks	56844	56873	59240	54344	76054	70768	73683	81834
Expanded clay	37054	30030	31599	32201	42863	38073	41111	45426
Stone wool	17856	17856	17856	17856	17856	17963	17238	16500
	1998	1999	2000	2001	2002	2003	2004	2005
Brickworks	82580	79066	80929	68669	66762	66723	71290	79592
Expanded clay	36233	36790	35313	25944	26768	23655	31477	34724
Stone wool	18632	18948	17258	15588	15910	15022	18072	18033
	2006	2007	2008	2009	2010	2011	2012	2013
Brickworks	78989	86356	61784	35814	35075	45979	39743	36710
Expanded clay	47503	61100	36551	14731	13714	15075	13388	23821
Stone wool	15490	19347	22579	16486	17050	16752	15012	13755




Figure 4.2.3 Consumption of CaCO<sub>3</sub> equivalents in the production of ceramics and stone wool.

Both the brickworks and expanded clay production displays a significant decrease from 2007 to 2009 that can be explained by the financial crises

### Emission factors

For production of bricks and expanded clay products the emission factors for SO<sub>2</sub> are determined from the individual companies reporting of SO<sub>2</sub> emission (environmental reports) for the years 2009-2011 and actual activity

for the corresponding years. These years were selected as the most complete data sets were available for these years. The SO<sub>2</sub> emissions have been adjusted for fuel related emissions to derive the process emissions. The PCDD/F emission factors shown in Table 4.2.9 are calculated from 0.018 µg per Mg product (Henriksen et al., 2006), using the total carbonate consumption (environmental reports), national production statistics (Statistics Denmark) and assumption of 2 kg per brick.

Stone wool emission factors for CO, NH<sub>3</sub> and TSP are average values measured and reported in the annual environmental reports for each Rockwool factory for the years 2001, 2004 and 2007-2013 (TSP for 1995-2013). PM<sub>10</sub> and PM<sub>2.5</sub> are estimated from the distribution between TSP, PM<sub>10</sub> and PM<sub>2.5</sub> (1/0.9/0.7). PCDD/F is known from Henriksen et al. (2006).

Table 4.2.9 Emission factors for Other mineral products, units are per Mg CaCO<sub>3</sub> equivalent.

Pollutant	Brickworks		Expanded clay		Mineral wool		Source
	Value	Unit	Value	Unit	Value	Unit	
SO <sub>2</sub>	6.5	kg	63.3	kg			Environmental reports*
CO					641	kg	Environmental reports**
NH <sub>3</sub>					16.0	kg	Environmental reports**
TSP					5.5	kg	Environmental reports**
PM <sub>10</sub>					5.0	kg	Environmental reports**
PM <sub>2.5</sub>					3.9	kg	Environmental reports**
PCDD/F	0.19	µg	0.17	µg	3.2	kg	Henriksen et al. (2006)*

\* Some recalculations were necessary to derive the desired units.

\*\* Calculated average from all measurements at all plants. For CO, only years before installation of abatement equipment are chosen.

Since 2007, emissions of CO, NH<sub>3</sub> and TSP have been measured yearly for mineral wool production, and the emission factors from Table 4.2.9 are therefore not used for these years. The measurements show a strong decrease in CO emissions from the two mineral wool factories in 2009 and 2010 respectively due to installation of abatement equipment.

### Emission trends

The only pollutants to which more than one source category contributes are SO<sub>2</sub> and PCDD/F. Therefore, only these two emissions are presented in the figures below. Figure 4.2.4 and Figure 4.2.5 show the emissions of SO<sub>2</sub> and PCDD/F respectively, emissions are presented individual for the three sources in "Other mineral products".




Figure 4.2.4 Emissions of SO<sub>2</sub> from ceramics.






Figure 4.2.5 Emissions of PCDD/F from ceramics and mineral wool.

Emissions of the remaining pollutants can be found in the NFR tables, where CO, NH<sub>3</sub> and particle emissions stem only from stone wool production.

#### 4.2.6 Source specific recalculations and improvements

##### Lime production

The activity data for lime no longer includes slacked lime. Personal communication with the industry made it clear that the inclusion of slacked lime resulted in a double counting because statistic on burned lime also includes lime that is later slacked.

An emission factor for black carbon has been added for lime production.

A factor 1000 error for the PCDD/F emission factor has been corrected for the years 1991-1999 and 2001-2003, resulting in increased emissions for these years.

##### Glass production

The activity data for container glass production has been altered for 1997-2012. Emissions were previously calculated based on fuel consumptions but this has now been changed to total production (mass product). This change necessitates a change and reassessment of the heavy metal emission factors. Where direct measured emissions are not available, Guidebook (2013) emission factors are now used to calculate emissions of heavy metals; As, Cd, Cr and Ni are new in this submission.

NMVOC from glass wool was not reported prior to 2010 causing a jump in the implied emission factor for NMVOC. In addition to the direct emissions of NMVOC known from the environmental reports, data on NMVOC has now been estimated for 1990-2006 using the average implied emission factor for 2007-2009.

Lesser chances were made for CO and NH<sub>3</sub> emissions from glass wool production for 1990-2012 and 1990-1995 respectively. Calculated emissions of these two pollutants were reassessed resulting in a small decrease of CO and increase of NH<sub>3</sub>. Particle size distribution for PM<sub>2.5</sub> was changed from 70 % of TSP to 80 % of TSP in accordance with the Guidebook (2013).

### Other mineral products

All activity data for the three source categories in “Other mineral products” have been changed to mass of CaCO<sub>3</sub> equivalent consumption instead of mass product produced. This was done because data on the mass produced is impossible to obtain, and uncertainties on the estimated data were unreasonably high. The consumption of raw materials, such as lime, on the other hand are available from 2006 and are easily estimated back to 1990 using surrogate data. All emission factors are changed to match the new activity data.

A factor 1000 error in the bricks and expanded clay production emission factors for PCDD/F has been corrected for the years 1991-1999 and 2001-2003, resulting in increased emissions for these years. For the same categories, a calculation error has resulted in lesser corrections for SO<sub>2</sub>.

### 4.2.7 Source specific planned improvements

The choice of CEPMEIP as source of the emission factors for particulate matter for lime production will be re-evaluated and a change to the latest edition of the EMEP/EEA guidebook will be considered.

The time series for production of yellow bricks and expanded clay products will be extended to include 1980-1989.

The SO<sub>2</sub> emission factor for bricks and expanded clay products will be improved based on additional data for coming years.

Emissions of black carbon will be added for container glass and glass wool production.

## 4.3 Chemical industry

### 4.3.1 Source category description

The sub-sector *Chemical industry* (NFR 2B) covers the following processes:

- 2B2 Nitric acid/fertiliser production (SNAP 040402/040407)
- 2B10a Other chemical industry
  - Sulphuric acid production (SNAP 040401)
  - Catalyst/fertiliser production (SNAP 040416/040407)
  - Production of chemical ingredients (SNAP 040500)
  - Pesticide production (SNAP 040525)
  - Production of tar products (SNAP 040527)

The time series for emission of acidifying substances, NMVOC, particulate matter, heavy metals, and POPs from *Mineral products* (NFR 2A) is available in the NFR tables. Table 4.2.1 presents an overview of emissions from 2013.

Table 4.3.1 Overview of 2013 emissions from Chemical industry.

	Total emission from Chemical industries	Fraction of IPPU, %	Largest contributor in Chemical industries	Emission from largest contributor	Fraction of Chemical industries, %
SO <sub>2</sub>	0.18 Gg	9.0	2B10a Other chemical industry	0.18 Gg	100.0
NO <sub>x</sub>	0.02 Gg	25.9	2B10a Other chemical industry	0.02 Gg	100.0
NMVOC	0.02 Gg	0.05	2B10a Other chemical industry	0.02 Gg	100.0
NH <sub>3</sub>	0.02 Gg	5.0	2B10a Other chemical industry	0.02 Gg	100.0
TSP	0.01 Gg	1.5	2B10a Other chemical industry	0.01 Gg	100.0

### 4.3.1 Nitric and sulphuric acid production

The production of sulphuric acid, nitric acid as well as NPK fertilisers has been concentrated at one company; Kemira GrowHow A/S (Kemira GrowHow, 2004). The production of sulphuric acid and nitric acid/fertiliser ceased in 1996/7 and in the middle of 2004, respectively. The following SNAP codes are covered:

- 04 04 01 Sulphuric acid
- 04 04 02 Nitric acid

The following pollutants are relevant for the nitric and sulphuric acid production process:

- SO<sub>2</sub>
- NO<sub>x</sub>
- NH<sub>3</sub>
- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>

#### Methodology

Information on emissions is obtained from environmental reports, contact to the company as well as information from the county. Information on emissions of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is available for 1990; 1994-2004, 1990; 1994-1997 and 1990-2004 respectively, TSP is available for 2000-2004. Implied emission factors are calculated for the years where measurements are available; these implied emission factors are then used to calculate emissions for the remaining years.

#### Activity data

The activity data regarding production of nitric and sulphuric acids is obtained through personal communication with Kemira (Kemira GrowHow, 2004 and 2005). The data are presented in Table 4.3.2.

Table 4.3.2 Production of nitric and sulphuric acid, Mg.

	1980	1981	1982	1983	1984	1985	1986	1987	1988
Nitric acid	350	350	350	350	350	350	315	357	383
Sulphuric acid	188	188	188	188	188	188	97	126	184
	1989	1990	1991	1992	1993	1994	1995	1996	1997
Nitric acid	402	450	412	364	343	348	390	360	366
Sulphuric acid	215	148	65	58	63	80	102	55	2
	1998	1999	2000	2001	2002	2003	2004		
Nitric acid	348	410	433	382	334	386	229		
Sulphuric acid	NO	NO	NO	NO	NO	NO	NO		

#### Emission factors

The calculated implied emission factors for SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> are presented in Table 4.3.3.

Table 4.3.3 Implied emission factors (IEF) for production of nitric and sulphuric acid.

Process	Pollutant	IEF	Unit
Nitric acid	NO <sub>x</sub>	0.95-1.79	kg/Mg
Nitric acid	NH <sub>3</sub>	0.03-0.26	kg/Mg
Sulphuric acid	SO <sub>2</sub>	1.40-2.69	kg/Mg

PM<sub>10</sub> and PM<sub>2.5</sub> are estimated from the distribution between TSP, PM<sub>10</sub> and PM<sub>2.5</sub> (1/0.8/0.6) from CEPMEIP (Values for 'Production of nitrogen fertiliser').

### Emission trends

The time series for SO<sub>2</sub> follows the amount of sulphuric acid produced, i.e. the fluctuation follows the activity until the activity ceased in 1997. The same is the case for NO<sub>x</sub> from production of nitric acid; see Figure 4.3.1.




Figure 4.3.1 Emissions from nitric and sulphuric acid production.

### 4.3.2 Catalyst production

Production of a wide range of catalysts and potassium nitrate (fertiliser) is concentrated at one company: Haldor Topsøe A/S situated in Frederikssund. The products are catalysts for many purposes (for hydro-processing, ammonia, DeNO<sub>x</sub>, methanol, hydrogen and synthesis gas, sulphuric acid, formaldehyde, and combustion catalysts) and potassium nitrate (fertiliser). The following SNAP code is covered:

- 04 04 16 Other: catalysts

The following pollutants are relevant for the catalyst production process:

- NO<sub>x</sub>
- NH<sub>3</sub>
- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>

### Methodology

The emissions of NO<sub>x</sub>, NH<sub>3</sub> and TSP from production of catalysts and fertilisers are measured yearly from 1996 to 2013 (TSP from 2000 to 2013) (Haldor Topsøe, 2014). The emissions from 1985-1995 were extrapolated.

The process-related NO<sub>x</sub> emission has been estimated as 80 % of the total NO<sub>x</sub> emission; Haldor Topsøe reports this assumption in their environmental report. The plant is equipped with DeNO<sub>x</sub> flue gas cleaning systems and depending on the efficiency of the cleaning system emission of NH<sub>3</sub> will occur.

### Activity data

The activity data regarding production of catalysts and fertiliser is obtained through environmental reports from Haldor Topsøe. The data are presented in Table 4.3.4.

Table 4.3.4 Production of catalysts and potassium nitrate, Gg (Haldor Topsøe, 2014).

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
Catalysts <sup>1</sup>						17.0	17.0	17.0	17.0	17.0
Potassium nitrate <sup>1</sup>						18.4	18.4	18.4	18.4	18.4
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Catalysts <sup>1</sup>	17.0	17.0	17.0	17.0	17.0	17.0	17.0	16.9	14.4	17.0
Potassium nitrate <sup>1</sup>	18.4	18.4	18.4	18.4	18.4	18.4	18.4	18.8	15.6	18.1
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Catalysts	17.2	19.5	19.3	15.3	22.0	23.2	20.3	20.7	28.1	22.5
Potassium nitrate	19.2	20.4	21.7	19.6	27.1	23.3	24.9	27.0	31.4	22.1
	2010	2011	2012	2013						
Catalysts	19.2	22.3	22.9	23.0						
Potassium nitrate	24.8	25.3	32.9	33.0						

<sup>1</sup> Production 1985-1996 assumed to be the average of 1997-2001.

### Emission factors

The calculated implied emission factors for NO<sub>x</sub>, NH<sub>3</sub> and particles are presented in Table 4.3.5.

Table 4.3.5 Implied emission factors for production of catalysts and potassium nitrate, Mg/Gg.

NO <sub>x</sub>	NH <sub>3</sub>	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>
0.30-1.76	0.26-3.70	0.11-0.59	0.09-0.48	0.06-0.36

PM<sub>10</sub> and PM<sub>2.5</sub> are estimated from the distribution between TSP, PM<sub>10</sub> and PM<sub>2.5</sub> (1/0.8/0.6) from CEPMEIP (Values for 'Production of nitrogen fertiliser').

### Emission trends

The emission of NH<sub>3</sub> shows an increasing trend throughout the 00's; from 14 Mg in 2000 to 165 Mg in 2009; in the same period the IEF fluctuates around the average 0.54 Mg per Gg but shows no trend. For the remaining time series, the NH<sub>3</sub> emission only varies between 9-21 Mg with the exception of 2010 where 123 Mg were emitted.

The emission of NO<sub>x</sub> decreases from the end of the 90's to the beginning of the 00's, in spite of the increasing production.

Emissions of NO<sub>x</sub>, NH<sub>3</sub> and TSP are shown in Figure 4.3.2.




Figure 4.3.2 Emissions from catalyst and fertiliser production.

### 4.3.3 Production of chemical ingredients

The production of chemical ingredients takes place in a number of different companies. One of the major companies is Danisco Grindsted located in Grindsted (Danisco Grindsted, 2014). The following SNAP code is covered:

- 04 05 00

The following pollutants are relevant for the production process of chemical ingredients:

- NMVOC

#### Methodology

The following description of the production of chemical ingredients is based on the environmental report from the company (Danisco Grindsted, 2014).

The raw materials are primarily natural or nature identical raw materials/substances: vegetable oils, animal fatty acids, glycerine, other organic substances, mineral acidic and alkaline compounds, solvents etc. The products are emulsifiers, stabilisers, flavours, enzymes, antioxidants, pharmaceuticals, and preservatives.

#### Activity data

Due to confidentiality no activity data are available.

#### Emission factors

Due to confidentiality no emission factors are available.

#### Emission trends

The emission of NMVOC from production of chemical ingredients has been measured from 1996 to 2013 (Danisco Grindsted, 2014). The emission has decreased from 100 to 12 Mg NMVOC in this period. However, no explanation can be given on these conditions, as information on activity is not available. The NMVOC emissions are presented in Table 4.3.6.

Table 4.3.6 Emissions from the production of chemical ingredients, Mg (Danisco Grindsted, 2014).

	1990	1991	1992	1993	1994	1995	1996	1997
NMVOG	100	100	100	100	100	100	100	93
	1998	1999	2000	2001	2002	2003	2004	2005
NMVOG	103	62	40	18	18	15	16	14
	2006	2007	2008	2009	2010	2011	2012	2013
NMVOG	15	17	15	12	12	12	12	12

#### 4.3.4 Pesticide production

The production of pesticides in Denmark is concentrated at one company: Cheminova A/S situated in Harboøre. The following SNAP code is covered:

- 04 05 25 Pesticide production

The following pollutants are relevant for the pesticide production process:

- SO<sub>2</sub>
- NMVOG

#### Methodology

The air emissions from Cheminova are measured from a number of sources; e.g. exhaust from process plant, sulphur recovery plant and biological sewage treatment plant. The environmental reports do only include some of the emissions and they do only present aggregated data.

Emissions of SO<sub>2</sub> and NMVOG are measured yearly for 1990-2013 and 1990-2000 respectively.

#### Activity data

Activity data for the production of pesticides are presented in Table 4.3.7.

Table 4.3.7 Production of pesticides, Mg (Cheminova, 2014).

	1990	1991	1992	1993	1994	1995	1996	1997	1998
Pesticides	37 671	39 631	29 764	38 988	41 913	45 320	55 800	56 500	52 985
	1999	2000	2001	2002	2003	2004	2005	2006	2007
Pesticides	64 264	60 284	60 376	55 464	52 849	65 310	53 504	52 575	49 796
	2008	2009	2010	2011	2012	2013			
Pesticides	49 747	37 484	31 000	31 000	31 000	30 000			

Activity data for 1990-1995 are estimated with the production in million DKK as surrogate data.

#### Emission factors

The implied emission factors for pesticide production are presented in Table 4.3.8.

Table 4.3.8 Implied emission factors for pesticide production, Claus process.

	Substance	Interval, kg/Mg	Average, kg/Mg
Pesticides	SO <sub>2</sub>	0.1-15.5	5.3
	NMVOG	0.5-10.4	2.2

### Emission trends

The emission of NMVOC from production of pesticides is reduced significantly from 1990 to 1993 (Cheminova, 2014). The decrease can be explained by introduction of flue gas cleaning equipment rather than any decrease in activity. The emission of SO<sub>2</sub> is from the sulphur regeneration plant (Claus plant).

The emission of NMVOC from production of pesticides is measured yearly from 1990 to 2000 (Cheminova, 2014) and estimated for 2001 to 2013. The implied emission factor based on 2000 data is used for estimation of 2001 to 2013 emissions.

Emissions are presented in Figure 4.3.3.




Figure 4.3.3 Emissions of SO<sub>2</sub> and NMVOC from pesticide production.

### 4.3.5 Production of tar products

One Danish factory situated in Nyborg produces tar products. The following SNAP code is covered:

- 04 05 27 Production of tar products

The following pollutants are relevant for the production process of tar products:

- SO<sub>2</sub>
- NMVOC
- Hg

#### Methodology

##### Activity data

No activity data are available.

##### Emission factors

The emissions are based on yearly measured process emissions reported in the environmental reports (Koppers, 2014). The emissions for the years 1985 – 2004 are assumed to be the same as for 2005.

##### Emission trends

Emissions are presented in Table 3.3.9.



Table 3.3.9 Emissions from production of tar products (Koppers, 2014).

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
SO <sub>2</sub>	210	210	210	210	210	210	210	210	210	210
NMVOG	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Hg	NE	NE	NE	NE	NE	4.9	4.9	4.9	4.9	4.9
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
SO <sub>2</sub>	210	210	210	210	210	210	210	210	210	210
NMVOG	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Hg	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9
	2005	2006	2007	2008	2009	2010	2011	2012	2013	
SO <sub>2</sub>	212	122	61	95	93	105	166	203	174	
NMVOG	0.9	1.2	1.0	0.5	2.0	1.3	1.1	1.2	0.6	
Hg	4.9	4.9	4.9	12.1	0.5	1.5	1.4	10.0	0.0	

NE: Not estimated

#### 4.3.6 Source specific recalculations and improvements

For the production of nitric and sulphuric acid, activity data and emission factors are now completed for the time series back to 1980.

The available data for production of catalysts and tar products was reassessed, leading to minor changes for 1990-1996 and 2007-2012.

#### 4.3.7 Source specific planned improvements

For catalyst production, it will be attempted to verify the assumptions on the split between combustion and process emissions for CO<sub>2</sub> and NO<sub>x</sub>.

For pesticides, the time series for SO<sub>2</sub> and NMVOG will be completed back to 1980. In addition to this, the exiting data will be reassessed as the interval of the implied emission factors seems very large.

For tar products, the emission of SO<sub>2</sub> will be extrapolated back to 1980 to comply with the requirement to have a time-series back to the base year. If possible emissions of NMVOG and Hg will also be extrapolated. And it will be evaluated whether the assumption to keep emissions constant back in time at the 2005 level is appropriate.

### 4.4 Metal production

#### 4.4.1 Source category description

The processes within metal industry in Denmark in relation to emission of other pollutants are:

- Steel production (SNAP 040207/040208)
- Iron production (SNAP 030303)
- Red bronze production (SNAP 040306)
- Secondary aluminium production (SNAP 030310)
- Secondary lead production (SNAP 030307)

The time series for emission of particulate matter, heavy metals, and POPs from *Metal production* (NFR 2C) is available in the NFR tables. Table 4.4.1 presents an overview of emissions from 2013.

Table 4.4.1 Overview of 2013 emissions from Metal production.

	Total emission from Metal industries	Fraction of IPPU; %	Largest contributor in Metal industries	Emission from largest contributor	Fraction of Metal industries, %
TSP	0.16 Gg	19.90	2C1 Iron and steel production	0.16 Gg	98.69
HMs	2.86 Mg	43.67	Zn from 2C7c Other metal production	0.63 Mg	22.16
POPs	0.21 kg	0.09	HCBs from 2C3 Aluminium production	0.14 kg	65.78

In the NFR tables, steel production and iron production is summed into one category called "Iron and steel production". This NFR sector 2C1 comprises three activities: An electric arc furnace (EAF) (until 2001/2002 and in 2005), rolling mills (from 2003) and grey iron foundries (whole time series). The most interesting activity from an emission perspective is the EAF. After the closing of the EAF, the site has since 2003 been used for rolling steel slabs imported from steelworks in other countries. This change in production results in large changes in activity data and emissions reported for the year 2002. In 2005, the EAF was shortly reopened, which explains the higher activity level this year.

Regarding the steelworks that use iron and steel scrap as raw material, the emissions to a large degree depend on the quality of the scrap. This fact may result in large annual variations for one or more of the heavy metals. This may also be the case for iron foundries, as they also use scrap as raw material, but they have not been subject to the same requirements to analyse emissions of heavy metals to air.

#### 4.4.2 Steel production

The production of semi-manufactured steel products (e.g. steel sheets/plates and bars) is concentrated at one company: Det Danske Stålvalseværk A/S situated in Frederiksværk. The following SNAP codes are covered:

- 04 02 07 Electric furnace steel plant
- 04 02 08 Rolling mill

The following pollutants are relevant for the steel production processes:

- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>
- Heavy metals: As, Cd, Cr, Cu, Hg, Ni, Pb, Zn
- HCB
- PCDD/F
- PCB

#### Methodology

The steelwork was closed down in January 2002 and then partly re-opened again in November 2002. The production of steel sheets/plates was reopened by DanSteel in 2003, the production of steel bars was reopened by DanScan Metal in March 2004, and the electro steelwork was reopened by DanScan Steel in January 2005. The production at DanScan Metal and Steel ceased in the last part of 2005 and in June 2006 DanScan Metal was taken over by Duferco; the future for the electro steelwork (DanScan Steel) is still uncertain and the plant has not been in operation since 2005. The timeline is presented in Activity data

Statistical data on activities are available in environmental reports from the single Danish plant (Stålvalseværket) supplemented with other literature; see Table 4.4.2.

Table 4.4.2 Overall mass flow for Danish steel production, Gg (Stålvalseværket, 2002; DanSteel, 2014).

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<b>Det danske stålvalseværk</b>											
Raw material	Iron and steel scrap	-	6301	557	-	673	657	664	735	737	691
Intermediate product	Steel slabs etc.	-	-	599	-	730	654	744	794	800	727
Product	Steel sheets	444	444	444	451	459	478	484	571	514	571
	Steel bars	170	170	170	217	264	239	235	245	238	226
	Products, total	6142	6142	614	6683	722	717	720	816	752	798
<b>DanSteel</b>											
Raw material	Steel slabs										
Product	Steel sheets										
		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Det danske stålvalseværk</b>											
Raw material	Iron and steel scrap	731	680					-			
Intermediate product	Steel slabs etc.	803	746					-			
Product	Steel sheets	380	469					-			
	Steel bars	251	256					-			
	Products, total	631	725				2504				
<b>DanSteel</b>											
Raw material	Steel slabs				553	600	515	561	635	590	254
Product	Steel sheets				469	506	433	468	520	484	211
		2010	2011	2012	2013						
<b>Det danske stålvalseværk</b>											
Raw material	Iron and steel scrap										
Intermediate product	Steel slabs etc.										
Product	Steel sheets										
	Steel bars										
	Products, total										
<b>DanSteel</b>											
Raw material	Steel slabs	457	490	338	460						
Product	Steel sheets	381	390	275	379						

<sup>1</sup>Jensen & Markussen (1993).

<sup>2</sup>Extrapolation.

<sup>3</sup>Intrapolation.

<sup>4</sup>Assumed

The mass balances / flow sheets presented in the annual environmental reports do not for all years tell about the changes in the stock and therefore the balance cannot be checked off.

### Emission factors

The applied emission factors are presented in Table 4.4.3. Regarding the electric arc furnace the emissions for all other pollutants than TSP have been estimated by use of emission factor from literature.

Table 4.4.3 Emission factors for steel production (Environmental reports).

	Unit	Electric Arc Furnace	Rolling Mill
TSP	g/Mg	61-147	6.44
PM10	g/Mg	80 % of TSP1	6.12
PM2.5	g/Mg	70 % of TSP1	3.86
As	g/Mg	0.0151	-
Cd	g/Mg	0.01-0.082	0.006
Cr	g/Mg	0.11	-
Cu	g/Mg	0.021	-
Hg	g/Mg	0.1-0.42	0.023
Ni	g/Mg	0.4-1.42	0.087

Pb	g/Mg	1.0-5.02	0.087
Zn	g/Mg	3.6-19.02	0.377
HCB	mg/Mg	3.23	-
PCDD/F	mg/Mg	0.81	-
PCB	mg/Mg	2.53	-

<sup>1</sup> EMEP/EEA Guidebook, Tier 2 no abatement

<sup>2</sup> Illerup et al. (1999)

<sup>3</sup> Nielsen et al. (2013)

### Emission trends

Emissions from the electro steelwork and rolling mill are presented in Table 4.4.4.

Table 4.4.4 Emissions from the electro steelwork and rolling mill.

	Unit	1990	1995	2000	2005	2010	2011	2012	2013
TSP	Mg			41.2	2.8	2.5	2.5	1.8	2.4
PM10	Mg			32.9	2.7	2.3	2.4	1.7	2.3
PM2.5	Mg			23.0	1.7	1.5	1.5	1.1	1.5
As	kg	9.2	10.8	9.5					
Cd	kg	38.9	21.5	16.4	2.5	2.2	2.3	1.6	2.2
Cr	kg	61.4	71.7	63.1					
Cu	kg	12.3	14.3	12.6					
Hg	kg	245.5	143.4	63.1	10.1	8.8	9.0	6.4	8.8
Ni	kg	757.4	430.2	252.4	37.7	33.1	33.8	23.9	32.9
Pb	kg	2966.9	1720.4	669.0	37.7	33.1	33.8	23.9	32.9
Zn	kg	11491.8	6546.7	3084.9	163.3	143.6	146.8	103.5	142.7
HCB	kg	2.0	2.3	2.0	2.14	2.31	2.29	2.12	2.61
PCDD/F	g	0.5	0.6	0.5					
PCB	kg	1.5	1.8	1.6	1.67	1.81	1.79	1.66	2.04

Due to the change in production process in the beginning of the 00's, the emissions (and even more so the implied emission factors) change drastically from 2001 to 2002 and from 2002 to 2003. Please refer to Figure 4.4.1 and Table 4.4.2.




Figure 4.4.1 Timeline for production at the Danish steelwork.

### Activity data

Statistical data on activities are available in environmental reports from the single Danish plant (Stålvalseværket) supplemented with other literature; see Table 4.4.2.

Table 4.4.2 Overall mass flow for Danish steel production, Gg (Stålvalseværket, 2002; DanSteel, 2014).

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<b>Det danske stålvalseværk</b>											
Raw material	Iron and steel scrap	-	630 <sup>1</sup>	557	-	673	657	664	735	737	691
Intermediate product	Steel slabs etc.	-	-	599	-	730	654	744	794	800	727
Product	Steel sheets	444	444	444	451	459	478	484	571	514	571
	Steel bars	170	170	170	217	264	239	235	245	238	226
	Products, total	614 <sup>2</sup>	614 <sup>2</sup>	614	668 <sup>3</sup>	722	717	720	816	752	798
<b>DanSteel</b>											
Raw material	Steel slabs										
Product	Steel sheets										
		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Det danske stålvalseværk</b>											
Raw material	Iron and steel scrap	731	680					-			
Intermediate product	Steel slabs etc.	803	746					-			
Product	Steel sheets	380	469					-			
	Steel bars	251	256					-			
	Products, total	631	725					250 <sup>4</sup>			
<b>DanSteel</b>											
Raw material	Steel slabs				553	600	515	561	635	590	254
Product	Steel sheets				469	506	433	468	520	484	211
		2010	2011	2012	2013						
<b>Det danske stålvalseværk</b>											
Raw material	Iron and steel scrap										
Intermediate product	Steel slabs etc.										
Product	Steel sheets										
	Steel bars										
	Products, total										
<b>DanSteel</b>											
Raw material	Steel slabs	457	490	338	460						
Product	Steel sheets	381	390	275	379						

<sup>1</sup>Jensen & Markussen (1993).

<sup>2</sup>Extrapolation.

<sup>3</sup>Intrapolation.

<sup>4</sup>Assumed

The mass balances / flow sheets presented in the annual environmental reports do not for all years tell about the changes in the stock and therefore the balance cannot be checked off.

### Emission factors

The applied emission factors are presented in Table 4.4.3. Regarding the electric arc furnace the emissions for all other pollutants than TSP have been estimated by use of emission factor from literature.

Table 4.4.3 Emission factors for steel production (Environmental reports).

	Unit	Electric Arc Furnace	Rolling Mill
TSP	g/Mg	61-147	6.44
PM <sub>10</sub>	g/Mg	80 % of TSP <sup>1</sup>	6.12
PM <sub>2.5</sub>	g/Mg	70 % of TSP <sup>1</sup>	3.86
As	g/Mg	0.015 <sup>1</sup>	-
Cd	g/Mg	0.01-0.08 <sup>2</sup>	0.006
Cr	g/Mg	0.1 <sup>1</sup>	-
Cu	g/Mg	0.02 <sup>1</sup>	-
Hg	g/Mg	0.1-0.4 <sup>2</sup>	0.023

Ni	g/Mg	0.4-1.4 <sup>2</sup>	0.087
Pb	g/Mg	1.0-5.0 <sup>2</sup>	0.087
Zn	g/Mg	3.6-19.0 <sup>2</sup>	0.377
HCB	mg/Mg	3.2 <sup>3</sup>	-
PCDD/F	mg/Mg	0.8 <sup>1</sup>	-
PCB	mg/Mg	2.5 <sup>3</sup>	-

<sup>1</sup> EMEP/EEA Guidebook, Tier 2 no abatement

<sup>2</sup> Illerup et al. (1999)

<sup>3</sup> Nielsen et al. (2013)

### Emission trends

Emissions from the electro steelwork and rolling mill are presented in Table 4.4.4.

Table 4.4.4 Emissions from the electro steelwork and rolling mill.

	Unit	1990	1995	2000	2005	2010	2011	2012	2013
TSP	Mg			41.2	2.8	2.5	2.5	1.8	2.4
PM <sub>10</sub>	Mg			32.9	2.7	2.3	2.4	1.7	2.3
PM <sub>2.5</sub>	Mg			23.0	1.7	1.5	1.5	1.1	1.5
As	kg	9.2	10.8	9.5					
Cd	kg	38.9	21.5	16.4	2.5	2.2	2.3	1.6	2.2
Cr	kg	61.4	71.7	63.1					
Cu	kg	12.3	14.3	12.6					
Hg	kg	245.5	143.4	63.1	10.1	8.8	9.0	6.4	8.8
Ni	kg	757.4	430.2	252.4	37.7	33.1	33.8	23.9	32.9
Pb	kg	2966.9	1720.4	669.0	37.7	33.1	33.8	23.9	32.9
Zn	kg	11491.8	6546.7	3084.9	163.3	143.6	146.8	103.5	142.7
HCB	kg	2.0	2.3	2.0	2.14	2.31	2.29	2.12	2.61
PCDD/F	g	0.5	0.6	0.5					
PCB	kg	1.5	1.8	1.6	1.67	1.81	1.79	1.66	2.04

Due to the change in production process in the beginning of the 00's, the emissions (and even more so the implied emission factors) change drastically from 2001 to 2002 and from 2002 to 2003. Please refer to Figure 4.4.1 and Table 4.4.2.

### 4.4.3 Iron production

Multiple grey iron foundries exist in Denmark, producing a wide range of products like e.g. cast iron pipes, central heating boilers and flywheels. The following SNAP codes are covered:

- 03 03 03 Grey iron foundries

The following pollutants are relevant for the iron production process:

- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>
- Heavy metals: As, Cd, Cr, Ni, Pb, Se, Zn
- HCB
- PCB

### Methodology

The emission of heavy metals from iron foundries is based on standard emission factors and yearly production statistics from Statistics Denmark (2013). The emission of TSP and distribution between TSP, PM<sub>10</sub> and PM<sub>2.5</sub>

(1/0.3/0.045) is obtained from CEPMEIP (2003). Emission factors for HCB and PCB are obtained from Nielsen et al. (2013).

#### Activity data

Statistical data on production in grey iron foundries are available from Statistics Denmark (2014) since 1998; activity data prior to this year are estimated. The activity data are presented in Table 4.4.5.

Table 4.4.5 Activity data, iron foundries, Gg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Grey iron foundries	103.0	100.4	97.8	95.3	92.7	90.2	87.6	85.1	85.8	86.0
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Grey iron foundries	101.7	94.5	103.0	104.5	113.0	99.9	81.5	72.8	71.4	41.4
	2010	2011	2012	2013						
Grey iron foundries	72.9	83.6	76.4	77.4						

#### Emission factors

The applied emission factors are presented in Table 4.4.6.

Table 4.4.6 Emission factors for grey iron foundries.

	Unit	Grey iron foundries
TSP	g/Mg	2000
PM <sub>10</sub>	g/Mg	600
PM <sub>2.5</sub>	g/Mg	90
As	g/Mg	0.3
Cd	g/Mg	0.14
Cr	g/Mg	1.1
Ni	g/Mg	1.3
Pb	g/Mg	7.2
Se	g/Mg	5
Zn	g/Mg	5
HCB	mg/Mg	0.04
PCB	mg/Mg	0.5

#### Emission trends

Emissions from grey iron foundries are presented in Table 4.4.7.

Table 4.4.7 Emissions from grey iron foundries.

	Unit	1990	1995	2000	2005	2010	2011	2012	2013
TSP	Mg			203.4	199.8	145.7	167.1	152.9	154.7
PM <sub>10</sub>	Mg			61.0	59.9	43.7	50.1	45.9	46.4
PM <sub>2.5</sub>	Mg			9.2	9.0	6.6	7.5	6.9	7.0
As	kg	30.9	27.1	30.5	30.0	21.9	25.1	22.9	23.2
Cd	kg	14.4	12.6	14.2	14.0	10.2	11.7	10.7	10.8
Cr	kg	113.3	99.2	111.9	109.9	80.2	91.9	84.1	85.1
Ni	kg	133.9	117.3	132.2	129.9	94.7	108.6	99.4	100.6
Pb	kg	741.6	649.4	732.2	719.4	524.6	601.7	550.3	557.1
Se	kg	515.0	451.0	508.5	499.6	364.3	417.8	382.1	386.9
Zn	kg	515.0	451.0	508.5	499.6	364.3	417.8	382.1	386.9
HCB	g	4.1	3.6	4.1	4.0	2.9	3.3	3.1	3.1
PCB	g	51.5	45.1	50.8	50.0	36.4	41.8	38.2	38.7

#### 4.4.1 Red bronze production

The following SNAP code is covered:

- 04 03 06 Allied metal manufacturing (Red bronze production)

The following pollutants are relevant for the red bronze production process:

- Heavy metals: Cd, Cu, Pb, Zn

#### Methodology

##### Activity data

Activity data are available for the years 1990 and 1995-1997. Data for 1991-1994 was interpolated and 1998-2013 are kept constant on the same level as 1997.

Table 4.4.8 Activity data for red bronze production.

	Unit	1990	1995	1996	1997
Red bronze production	Mg	3895	4350	4400	4532

#### Emission factors

The applied emission factors are presented in Table 4.4.9 and are all referenced to Illerup et al. (1999).

Table 4.4.9 Emission factors for red bronze production.

Cd	g/Mg	1
Cu	g/Mg	10
Pb	g/Mg	15
Zn	g/Mg	140

#### Emission trends

Emissions trends for Cd, Cu, Pb, and Zn from red bronze production are presented in Table 4.4.10.

Table 4.4.10 Emissions from red bronze production, kg.

	1990	1991	1992	1993	1994	1995	1996	1997-2013
Cd	3.9	4.0	4.1	4.2	4.3	4.4	4.4	4.5
Cu	39.0	39.9	40.8	41.7	42.6	43.5	44.0	45.3
Pb	58.4	59.8	61.2	62.5	63.9	65.3	66.0	68.0
Zn	545.3	558.0	570.8	583.5	596.3	609.0	616.0	634.5



#### 4.4.1 Secondary aluminium production

Two active Danish production sites have been identified for secondary aluminium; “Stena Aluminium” and “Jydsk Aluminium Industri”. The following SNAP codes are covered:

- 03 03 10 Secondary aluminium production

The following pollutants are relevant for the secondary aluminium production:

- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>
- Heavy metals: Cd, Pb
- HCB
- PCDD/F
- PCB

#### Methodology

Secondary aluminium industries were identified from a list of companies with the relevant environmental approvals acquired from the Danish Environmental Agency. The largest producer, called Stena Aluminium, stood for approximately 90 % of the total Danish production until the factory was closed in the end of 2008.

#### Activity data

The activity data are presented in Table 4.4.11.

Table 4.4.11 Activity data for secondary aluminium production (Stena Aluminium, 2008 and Jydsk Aluminiums Industri, 2014), Gg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Stena Aluminium <sup>1</sup>	30.23	30.23	30.23	30.23	30.23	30.23	25.06	31.79	32.98	28.48
Jydsk Aluminium <sup>2, 3, 4</sup>	1.26	1.26	1.26	1.26	1.74	2.22	2.70	3.00	3.30	3.61
Total	31.49	31.49	31.49	31.49	31.97	32.45	27.76	34.79	36.28	32.08
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Stena Aluminium	32.87	33.38	32.13	26.42	19.56	23.43	31.32	35.12	36.22	
Jydsk Aluminium <sup>2</sup>	3.91	4.21	4.51	4.81	5.11	5.42	5.72	6.02	6.07	3.80
Total	36.78	37.59	36.64	31.23	24.67	28.85	37.03	41.14	42.29	3.80
	2010	2011	2012	2013						
Stena Aluminium										
Jydsk Aluminium <sup>2</sup>	5.20	6.26	6.49	6.84						
Total	5.20	6.26	6.49	6.84						

<sup>1</sup>1990-1995: Calculated average of 1996-2000.

<sup>2</sup>1993, 1996, 2007-2013: Estimated based on information from the environmental reports.

<sup>3</sup>1990-1992: Estimated based on 1993.

<sup>4</sup>1994-1995, 1997-2006: Interpolated.

#### Emission factors

Emission factors for the production of secondary aluminium are presented in Table 4.4.12.

Table 4.4.12 Emission factors for secondary aluminium production.

Pollutant	Unit	Value	Source
TSP	kg/Mg	0.120	Calculated based on environmental reports from Stena, average for 1998-2000
PM <sub>10</sub>	kg/Mg	0.084	Particle distribution from EMEP/EEA Guidebook (2013)
PM <sub>2.5</sub>	kg/Mg	0.033	Particle distribution from EMEP/EEA Guidebook (2013)
Cd	g/Mg	0.032	Calculated based on environmental reports from Stena, average for 1998-2000
Pb	g/Mg	0.146	Calculated based on environmental reports from Stena, average for 1998-2000
HCB	g/Mg	0.020	Nielsen et al. (2013)
PCDD/F	mg/Mg	0.035	EMEP/EEA Guidebook (2013)
PCB	mg/Mg	3.40	Nielsen et al. (2013)

#### Emission trends

Emissions from secondary aluminium production are available in the NFR tables.

#### 4.4.2 Secondary lead production

One Danish company producing secondary lead has been identified; Hals Metal. The following SNAP codes are covered:

- 03 03 07 Secondary lead production

The following pollutants are relevant for the secondary lead production:

- TSP
- PM<sub>10</sub>
- PM<sub>2.5</sub>
- Heavy metals: As, Cd, Pb, Zn
- HCB
- PCDD/F
- PCB

#### Methodology

Only one Danish company, called Hals metal, has been identified as producing secondary lead from scrap metal. In addition to Hals metal, old lead tiles from castles, churches etc. are melted and recast on site during preservation of the many historical buildings in Denmark.

#### Activity data

Activity data from Hals metal is provided by the company. A clause affected in 2002 meant that Hals metal could no longer burn cables containing lead. The processing of cables was therefore stopped and the company's activity changed to melting. This transition resulted in a low activity in 2003.

The activity of recasting lead tiles is not easily found because it is spread out on many craftsmen and poorly regulated. However, an estimate by Lassen et al. (2004) stated that 200-300 Mg lead tiles were recast in 2000. Since the building stock worthy of preservation is constant, it is considered reasonable to also let the activity of recasting of lead tiles be constant.

Activity data for secondary lead is shown in Table 4.4.13.

Table 4.4.13 Activity data for secondary lead production (Hals metal, 2014 and Lassen et al., 2004), Mg.

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Hals metal	540	540	540	750	750	750	540	540	540	540
Lead tiles	250	250	250	250	250	250	250	250	250	250
Total	790	790	790	1000	1000	1000	790	790	790	790
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Hals metal	540	1080	419	64	520	691	500	670	582	780
Lead tiles	250	250	250	250	250	250	250	250	250	250
Total	790	1330	669	314	770	941	750	920	832	1030
	2010	2011	2012	2013						
Hals metal	635	938	412	533						
Lead tiles	250	250	250	250						
Total	885	1188	662	783						

#### Emission factors

Emission factors are presented in Table 4.4.14.

Table 4.4.14 Emission factors for secondary lead production.

Pollutant	Value	Unit	Reference
TSP	1.63	kg/Mg	EMEP/EEA Guidebook (2013)
PM <sub>10</sub>	1.30	kg/Mg	EMEP/EEA Guidebook (2013)
PM <sub>2.5</sub>	0.65	kg/Mg	EMEP/EEA Guidebook (2013)
As	3.5	g/Mg	EMEP/EEA Guidebook (2013)
Cd	1.1	g/Mg	EMEP/EEA Guidebook (2013)
Pb	426	g/Mg	EMEP/EEA Guidebook (2013)
Zn	2.6	g/Mg	EMEP/EEA Guidebook (2013)
HCB	0.3	mg/Mg	Nielsen et al. (2013)
PCDD/F	8.0	µg/Mg	EMEP/EEA Guidebook (2013)
PCB	7.3	mg/Mg	Nielsen et al. (2013)

#### Emission trends

Emissions from secondary lead production are available in the NFR tables.

#### 4.4.3 Source specific recalculations and improvements

For the electro steelwork, emission factors for As, Cr, Cu and PCDD/F have been added to this year's inventory. Emission factors for Cd, Hg, Ni, Pb and Zn were reassessed and are now calculated based on the measurements provided by the factory's environmental reports, the change in emission factors have led to a strong increase in emissions. The emission factor for TSP was lowered for 2001 down to the measured level of 2000.

The source of secondary aluminium production has been reassessed for this year's inventory. The work on this source category has resulted in all new activity data (4-26 % increase) and a completion of the time series; meaning that 1990-1994 and 2009-2013 are new years for this source. For the emission factors, particles are now calculated based on measurements from the industry, this change has caused a decrease. The PCDD/F emission factor now refers to the EMEP/EEA Guidebook (2013); this change has caused an increase in PCDD/F emissions. Cd, Pb, HCB and PCB from aluminium production are new in this year's inventory.

The source of secondary lead production has also been reassessed. The work on this source category has resulted in all new activity data (from 5 Gg in all years to 0.3-1.2 Gg in the new time series). Emission factors for As, Zn and PCDD/F are new in this year's inventory while Cd, Pb and particle emission factors now refer to the EMEP/EEA Guidebook (2013) resulting in large in-

creases in emissions. The emission factor for Cu was removed due to the lack of a reference.

Secondary zinc production (SNAP 030308) does not occur in Denmark and never has. The reporting of emissions in this category was the result of a mistake that has now been corrected.

#### **4.4.4 Source specific planned improvements**

All emission factors for iron foundries will be reviewed and clear references provided. Activity data from Statistics Denmark will be used for iron foundries for the whole time series.

It will be investigated whether activity data for red bronze production (Allied metal manufacturing) are available from Statistics Denmark.

An emission factor for black carbon will be added for secondary aluminium production.

### **4.5 Non-energy products from fuels and solvent use**

#### **4.5.1 NMVOCs used as solvents**

##### **Description of source category**

Non-energy products from fuels and solvent use (NFR 2D) includes the following categories:

- Domestic solvent use including fungicides (NFR 2D3a, SNAP 060408 & 060411)
- Road paving with asphalt (NFR 2D3b, SNAP 040611)
- Asphalt roofing (NFR 2D3c, SNAP 040610)
- Coating applications (NFR 2D3d, SNAP 0601)
- Degreasing (NFR 2D3e, SNAP 0602)
- Dry cleaning (NFR 2D3f, SNAP 060202)
- Chemical products (NFR 2D3g, SNAP 0603)
- Printing (NFR 2D3h, SNAP 060403)
- Other solvent use (NFR 2D3i, SNAP 0604)

Methodologies, activity data, emission factors and pollutants for Road paving with asphalt (NFR 2D3b) and Asphalt roofing (NFR 2D3c) are described in their respective sections below. For all other categories in NFR 2D the methods for compiling activity data and emission factors, values of activity data and emission factors and calculated emissions are presented in this section.

Solvent use categories are aggregated according to the following four categories, which corresponds to the grouping in IPCC (2006):

- Coating applications (NFR 2D3d)
- Degreasing (NFR 2D3e) and Dry cleaning (NFR 2D3f)
- Chemical products (NFR 2D3g)
- Domestic solvent use including fungicides (NFR 2D3a), Printing (NFR 2D3h) and Other solvent use (NFR 2D3i)

Only NMVOCs used as solvents are relevant for these categories. Solvents are chemical compounds that are used on a global scale in industrial processes and as constituents in final products to dissolve e.g. paint, cosmetics,

adhesives, ink, rubber, plastic, pesticides, aerosols or are used for cleaning purposes, i.e. degreasing. NMVOCs are main components in solvents - and solvent use in industries and households is typically the dominant source of anthropogenic NMVOC emissions (UNFCCC, 2008; Pärt, 2005; Karjalainen, 2005). In industrial processes where solvents are produced or used, NMVOC emissions to air and as liquid can be recaptured and either used or destroyed. Solvent containing products are used indoor and outdoor and the majority of solvent sooner or later evaporate. A small fraction of the solvent ends up in waste or as emissions to water and may finally also contribute to air pollution by evaporation from these compartments. Emission inventories for solvents are based on model estimates, as direct and continuous emissions are only measured from a limited number of sources.

### **Methodology**

Until 2002 the Danish solvent emission inventory was based on questionnaires, which were sent to selected industries and sectors requiring information on solvent use. In 2003 it was decided to implement a method that is more complete, accurate and transparent with respect to including the total amount of used solvent, attributing emissions to industrial sectors and households and establishing a reliable model that is readily updated on a yearly basis.

The method is mainly based on the detailed approach and methodology described in EMEP/EEA (2013) and IPCC (2006), and emissions are calculated for industrial sectors, households for the stated NFR sectors, as well as for individual pollutants.

Emission modelling of solvents can basically be done in two ways: 1) By estimating the amount of (pure) solvents consumed, or 2) By estimating the amount of solvent containing products consumed, taking account of their solvent content (EMEP/EEA, 2013).

In 1) all relevant solvents must be estimated, or at least those together representing more than 90 % of the total pollutant emission, and in 2) all relevant source categories must be inventoried or at least those together contributing more than 90 % of the total pollutant emission. A simple approach is to use a per capita emission for each category, whereas a detailed approach is to get all relevant consumption data (EMEP/EEA, 2013; IPCC, 2006).

The detailed method 1) is used in the Danish emission inventory for solvent use, thus representing a chemicals approach, where each pollutant is estimated separately. The sum of emissions of all estimated pollutants used as solvents equals the pollutant emission from solvent use.

### **Pollutant list**

The definitions of solvents and (NM)VOC that are used in the Danish emission inventory are as defined in the solvent directive (Directive 1999/13/EC) of the EU legislation: "Organic solvent shall mean any VOC which is used alone or in combination with other agents, and without undergoing a chemical change, to dissolve raw materials, products or waste materials, or is used as a cleaning agent to dissolve contaminants, or as a dissolver, or as a dispersion medium, or as a viscosity adjuster, or as a surface tension adjuster, or a plasticiser, or as a preservative". VOCs are defined as follows: "Volatile organic compound shall mean any organic compound having at 293.15 K a

vapour pressure of 0.01 kPa or more, or having a corresponding volatility under the particular condition of use”.

This implies that some NMVOCs, e.g. ethylene glycol, that have vapour pressures just around 0.01 kPa at 20 °C, may only be defined as VOCs at use conditions with higher temperature. However, use conditions at elevated temperatures are typically found in industrial processes. Here the capture of solvent fumes is often efficient, thus resulting in small emissions (communication with industries).

The Danish list of NMVOCs comprises approx. 30 pollutants or pollutant groups representing more than 95 % of the total emission from solvent use, cf. Table 4.5.4.

#### **Activity data**

For each pollutant or product a mass balance is formulated:

$$\text{Consumption} = (\text{production} + \text{import}) - (\text{export} + \text{destruction/disposal} + \text{hold-up}) \quad (\text{Eq. 1})$$

Data on production, import and export amounts of solvents and solvent containing products are collected from Statistics Denmark (2014), which contains detailed statistical information on the Danish society. Manufacturing and trading industries are committed to reporting production and trade figures to the Danish Customs & Tax Authorities in accordance with the Combined Nomenclature. Import and export figures are available on a monthly basis from 1990 to present and contain trade information from 272 countries world-wide. Production figures are reported quarterly as “industrial commodity statistics by commodity group and unit” from 1990 to present. Prior to 1990 the figures are assumed constant on a 1990 level.

Destruction and disposal of solvents lower the pollutant emissions. In principle this amount must be estimated for each pollutant in all industrial activities and for all uses of pollutant containing products. At present the solvent inventory only considers destruction and disposal for a limited number of pollutants. For some pollutants it is inherent in the emission factor, and for others the reduction is specifically calculated from information obtained from the industry or literature.

Hold-up is the difference in the amount in stock in the beginning and at the end of the year of the inventory. No information on solvents in stock has been obtained from industries. Furthermore, the inventory spans over several years so there will be an offset in the use and production, import and export balance over time.

In some industries the solvents are consumed in the process, e.g. in the graphics and plastic industry, whereas in the production of paints and lacquers the solvents are still present in the final product. These products can either be exported or used in the country. In order not to double count consumption amounts of pollutants it is important to keep track of total solvent use, solvents not used in products and use of solvent containing products. Furthermore some pollutants may be represented as individual pollutants and also in chemical groups, e.g. “o-xylene”, “mixture of xylenes” and “xylene”. Some pollutants are better inventoried as a group rather than individual pollutants, due to missing information on use or emission for the indi-

vidual pollutants. The Danish inventory considers single pollutants, with a few exceptions.

Activity data for pollutants are thus primarily calculated from Equation 1 with input from Statistics Denmark (2014). When Statistics Denmark holds no information on production, import and export or when more reliable information is available from industries, scientific reports or expert judgements the data can be adjusted or even replaced. The used amounts of products (activity data) in Table 4.5.1 are derived from used amounts of pollutants by assessing the amount of pollutants that is comprised within products belonging to each of the categories.

Table 4.5.1 Activity data (AD) in Gg per year. Complete time series can be seen in [Annex 3C-1](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Coating applications (NFR 2D3d)	83.2	83.2	92.1	105	75.2	45.8	43.8	43.3	47.3
Degreasing (NFR 2D3e) and Dry cleaning (NFR 2D3f)	1.41	1.41	1.53	0.59	0.37	0.25	0.22	0.055	0.097
Chemical products (NFR 2D3g)	406	406	504	567	740	641	640	516	518
Domestic solvent use - including fungicides (NFR 2D3a), Printing (NFR 2D3h) and Other solvent use (NFR 2D3i)	197	197	247	230	204	170	169	169	180

#### Emission factors

For each pollutant the emission is calculated by multiplying the consumption with the fraction emitted (emission factor), according to:

$$\text{Emission} = \text{consumption} * \text{emission factor}$$

The present Danish method uses emission factors that represent specific industrial activities, such as processing of polystyrene, dry cleaning etc. or that represent use categories, such as paints and detergents. Some pollutants have been assigned emission factors according to their water solubility. Higher hydrophobicity yields higher emission factors, since a lower amount ends in waste water, e.g. ethanol (hydrophilic) and turpentine (hydrophobic).

Emission factors for solvents are categorised in four groups in ascending order: (1) Lowest emission factors in the chemical industry, e.g. lacquer and paint manufacturing, due to emission reducing abatement techniques and destruction of solvent containing waste, (2) Other processes in industry, e.g. graphic industry, have higher emission factors, (3) Non-industrial use, e.g. auto repair and construction, have even higher emission factors, (4) Diffuse use of solvent containing products, e.g. painting, where practically all the pollutant present in the products will be released during or after use.

For a given pollutant the consumed amount can thus be attributed with two or more emission factors; one emission factor representing the emissions occurring at a production or processing plant and one emission factor representing the emissions during use of a solvent containing product. If the chemical is used in more processes and/or is present in several products more emission factors are assigned to the respective chemical amounts.

Emission factors can be defined from surveys of specific industrial activities or as aggregated factors from industrial branches or sectors. Furthermore, emission factors may be characteristic for the use pattern of certain products.

The emission factors used in the Danish inventory also rely on the work done in a joint Nordic project (Fauser et al., 2009).

In Table 4.5.2 the emission factors are listed. They are based on the values in the Guidebook (EMEP/EEA, 2013) and adjusted on a country specific basis according to the assessment described above. See more details in the QA/QC section.

Table 4.5.2 Emission factors in Gg NMVOC per Gg AD. Complete time series can be seen in [Annex 3C-2](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Coating applications (NFR 2D3d)	0.0614	0.0614	0.0641	0.0608	0.0575	0.0600	0.0656	0.0660	0.0652
Degreasing (NFR 2D3e) and Dry cleaning (NFR 2D3f)	5E-05	5E-05	5E-05	5E-05	5E-05	5E-05	5E-05	5E-05	5E-05
Chemical products (NFR 2D3g)	0.0200	0.0200	0.0185	0.0123	0.00844	0.00787	0.00753	0.00944	0.00889
Domestic solvent use including fungicides (NFR 2D3a), Printing (NFR 2D3h) and Other solvent use (NFR 2D3i)	0.125	0.125	0.121	0.121	0.102	0.115	0.114	0.113	0.118

#### Source allocation

The Danish Working Environment Authority (WEA) is administrating the registrations of chemicals and products to the Danish product register. All manufacturers and importers of products for occupational and commercial use are obliged to register. The following products are comprised in the registration agreement:

- Chemicals and materials that are classified as dangerous according to the regulations set up by the Danish Environmental Protection Agency (EPA).
- Chemicals and materials that are listed with a limit value on the WEA "limit value list".
- Materials, containing 1 % or more of a chemical, which is listed on the WEA "limit value list".
- Materials, containing 1 % or more of a chemical, which are classified as hazardous to humans or the environment according to the EPA rules on classification.

There are the following important exceptions for products, which do not need to be registered:

- Products exclusively for private use.
- Pharmaceuticals ready for use.
- Cosmetic products.

The Danish product register does therefore not comprise a complete account of used pollutants. Source allocations of exceptions from the duty of declaration are done based on information from trade organisations, industries, scientific reports and information from the internet.

The data base Substances in Preparations in the Nordic Countries (SPIN) ([www.spin2000.net](http://www.spin2000.net)) holds information on use of various pollutants in product and activities, i.e. Use Categories Nordic (UCN), and on use in industrial categories, i.e. according to the standard nomenclature for economic activities (NACE) system ([http://ec.europa.eu/comm/competition/mergers/cases/index/nace\\_all.html](http://ec.europa.eu/comm/competition/mergers/cases/index/nace_all.html)). The use amount from DK Statistics is first distributed in SNAP catego-



ries according to UCN data, and second according to NACE industrial use in NFR categories.

### Emission trends

Table 4.5.3 and Figure 4.5.1 show the emissions of NMVOC from 1985 to 2013, where the used amounts of single pollutants have been assigned to specific products and NFR sectors. From 1985 to 1990 the emission level is set constantly equal to the 1990 emission level, due to missing reliable data. A general increase is seen for all sectors from 1990 to 1996 followed by a decrease from 1997 to 2006 and stagnation in the period 2007 to 2013.

In Table 4.5.4 the emission for 2013 is split into individual pollutants. The most abundantly used solvents are ethanol, turpentine, or white spirit defined as a mixture of stoddard solvent and solvent naphtha and propylalcohol. Ethanol is used as solvent in the chemical industry and as windscreen washing agent. Turpentine is used as thinner for paints, lacquers and adhesives. Propylalcohol is used in cleaning agents in the manufacture of electrical equipment, flux agents for soldering, as solvent and thinner and as windscreen washing agent. Household emissions are dominated by propane and butane, which are used as aerosols in spray cans, primarily in cosmetics. For some pollutants the emission factors are precise but for others they are rough estimates. The division of emission factors into four categories implies that high emission factors are applicable for use of solvent containing products and lower emission factors are applicable for use in industrial processes.

Table 4.5.3 Emissions in Gg NMVOC per year. Complete time series can be seen in [Annex 3C-3](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Coating applications (NFR 2D3d)	5.11	5.11	5.90	6.40	4.32	2.75	2.87	2.86	3.09
Degreasing (NFR 2D3e) and Dry cleaning (NFR 2D3f)	7E-05	7E-05	8E-05	3E-05	2E-05	1E-05	1E-05	3E-06	5E-06
Chemical products (NFR 2D3g)	8.14	8.14	9.32	6.96	6.25	5.05	4.81	4.87	4.60
Domestic solvent use including fungicides (NFR 2D3a), Printing (NFR 2D3h) and Other solvent use (NFR 2D3i)	24.7	24.7	30.0	27.8	20.8	19.4	19.2	19.1	21.3
<b>Total NMVOC</b>	<b>38.0</b>	<b>38.0</b>	<b>45.2</b>	<b>41.1</b>	<b>31.4</b>	<b>27.2</b>	<b>26.9</b>	<b>26.9</b>	<b>29.0</b>




Figure 4.5.1 NMVOC emissions in Gg NMVOC per year. The methodological approach for finding emissions in the period 1985 – 2013 is described in the text. Figures can be seen in Table 4.5.3 and in [Annex 3C-3](#).

Table 4.5.4 2013 NMVOC emissions of single pollutants or pollutant groups.

Pollutant	CAS no	Emissions 2013 (t)
ethanol	64-17-5	9740
turpentine (white spirit: stoddard solvent and solvent naphtha)	64742-88-7	7213
	8052-41-3	
cyanates	79-10-7	2324
propyl alcohol	67-63-0	2301
pentane	109-66-0	1631
propylene glycol	57-55-6	1085
methanol	67-56-1	1047
acetone	67-64-1	700
propane	74-98-6	654
butane	106-97-8	654
butanone	78-93-3	283
xylene	1330-20-7	264
	95-47-6	
	108-38-3	
	106-42-3	
glycol ethers	110-80-5	199
	107-98-2	
	108-65-6	
	34590-94-8	
	112-34-5	
	and others	
ethylene glycol	107-21-1	142
toluene	108-88-3	142
styrene	100-42-5	125
phenol	108-95-2	119
cyclohexanones	108-94-1	113
butanols	78-92-2	107
	2517-43-3	
	and others	
formaldehyde	50-00-0	76.2
ethyl acetate	141-78-6	34.1
1-butanol	71-36-3	16.1
butyl acetate	123-86-4	13.7
acyclic aldehydes	78-84-2	3.2
	111-30-8	
	and others	
tetrachloroethylene	127-18-4	0.80
Total 2011		28988

## 4.5.2 Road paving with asphalt

### Description of source category

Road paving with asphalt is an activity that can be found all over the country and especially in relation to establishing new traffic facilities. The raw materials for construction of transport facilities are prepared on one of the plants located near the locality of application to limit the transport distance. The asphalt concrete is mixed and brought to the locality of application on a truck.

Transport facilities are constructed by a number of different layers:

- a load bearing layer (e.g. course gravel)
- an adhesive layer (liquefied asphalt e.g. "cutback" asphalt or asphalt emulsion)
- a wearing coarse (e.g. hot mix asphalt concrete)

Different qualities of "cutback" asphalt (e.g. asphalt dissolved in organic solvents/petroleum distillates) and asphalt emulsion contains different

kinds and amounts of solvent. Cutback asphalt contains 25-45%v/v solvent e.g. heavy residual oil, kerosene-type solvent, naphtha or gasoline solvent. Approximately 500.000 litre solvent evaporates annually from the use of "cutback" asphalt (Asfaltindustrien, 2003). This amount of solvent, which is added to the asphalt, is comprised in the solvent categories above with an emission factor of approximately unity. This means that NMVOC emissions from "cutback" asphalt in Road paving NFR 2D3b only include emissions from the asphalt fraction as quantified in Table 4.5.5.

#### Methodology

Emissions are calculated from Eq. 1 for the pollutants NMVOC and CO.

#### Activity data

The use amounts of asphalt for road paving have been compiled from production, import and export statistics of asphalt products in Statistics Denmark (2014).

Table 4.5.5 Activity data for asphalt in road paving in Gg per year. Complete time series can be seen in [Annex 3C-4](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Road paving with asphalt (NFR 2D3b)	2 370	2 370	3 144	2 933	3 879	3 005	3 896	3 233	3 339

#### Emission factors

Default emission factors are derived from EMEP/EEA (2013) and US EPA (2004).

Table 4.5.6 Emission factors for NMVOC and CO from road paving with asphalt.

		Road paving with asphalt (incl. cutback)	
NMVOC	g per tonnes		16
CO	g per tonnes		75

#### Emission trends

Table 4.5.7 NMVOC and CO emissions in Gg per year from road paving with asphalt. Complete time series can be seen in [Annex 3C-5](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
NMVOC	0.0379	0.0379	0.0503	0.0469	0.0621	0.0481	0.0623	0.0517	0.0534
CO	0.178	0.178	0.236	0.220	0.291	0.225	0.292	0.242	0.250

### 4.5.3 Asphalt roofing

#### Methodology

Emissions are calculated from Eq. 1 for the pollutants NMVOC and CO.

#### Activity data

The use amounts of asphalt for roofing has been compiled from production, import and export statistics of asphalt products in Statistics Denmark (2014).

Table 4.5.8 Activity data for asphalt roofing in Gg per year. Complete time series can be seen in [Annex 3C-6](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Asphalt roofing (NFR 2D3c)	120	120	123	204	187	105	134	131	125

### Emission factors

Default emission factors are derived from EMEP/EEA (2013) and US EPA (2004).

Table 4.5.9 Emission factors for NMVOC and CO from asphalt roofing.

Asphalt roofing		
NMVOC	g per tonnes	80
CO	g per tonnes	9.5

### Emission trends

Table 4.5.10 NMVOC and CO emissions in Gg per year from asphalt roofing. Complete time series can be seen in [Annex 3C-7](#).

	1985	1990	1995	2000	2005	2010	2011	2012	2013
NMVOC	0.0096	0.0096	0.0099	0.0163	0.0150	0.0084	0.0108	0.0105	0.0100
CO	0.00114	0.00114	0.00117	0.00194	0.00178	0.00100	0.00128	0.00124	0.00119

### 4.5.4 Source specific recalculations and improvements

The amount of solvent, which is added to the asphalt in “cutback”, is comprised in Other solvent use (NFR 2D3i), above with an emission factor of approximately unity. This amount was previously included in Road paving with asphalt (NFR 2D3b) as “cutback”. In the improved inventory NMVOC emissions from “cutback” asphalt in Road paving (NFR 2D3b) only include emissions from the asphalt fraction.

A change in allocation of amounts from Statistics Denmark (2014) has caused an increase in activity data for roofing, e.g. 2012: from 75.5 Gg to 131 Gg, and a relatively small increase for road paving, e.g. for 2012: from 3223 Gg to 3233 Gg.

Emission factors for NMVOC and CO have been updated with values from EMEP/EEA (2013) and US EPA (2004) for road paving and asphalt roofing.

### 4.5.5 Source specific planned improvements

Emissions of TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, BC, PAHs and possibly heavy metals will be included for road paving and asphalt roofing, with emissions factors from EMEP/EEA (2013) and US EPA (2004).

## 4.6 Other product use

### Methodology

The following sources and pollutants are included:

- Fireworks (SNAP 060601) (SO<sub>2</sub>, CO, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, As, Cd, Cr, Cu, Hg, Ni, Pb, Zn)
- Tobacco (SNAP 060602) (SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, PCDD/F, Benzo[b]fluoranthene, Benzo[k]fluoranthene, Benzo[a]pyrene, Indeno[1,2,3-cd]pyrene)
- Charcoal (SNAP 060605) (SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, As, Cd, Cr, Cu, Hg, Ni, Pb, Se, Zn, HCB, PCDD/F, Benzo[b]fluoranthene, Benzo[k]fluoranthene, Benzo[a]pyrene, Indeno[1,2,3-cd]pyrene)
- Paraffin wax use (SNAP 060606) (CO, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, PCDD/F, Benzo[k]fluoranthene, Benzo[a]pyrene, Indeno[1,2,3-cd]pyrene)

Method 2) in EMEP/EEA (2013) is used for calculating pollutant emissions from fireworks, tobacco, charcoal for barbeques and paraffin wax use (candles).

#### **Activity data**

Activity data are derived from Eq. 1 with import, export and production data from Statistics DK (2014). See [Annex 3C-8](#).

#### **Emission factors**

The emission factor for fireworks for Pb was changed in 2000 and Hg and Pb, along with any compounds derived here from, were forbidden in 2003 and 2007, respectively. Emissions are therefore noted as not occurring for these years and forward. Full time series for emission factors can be found in [Annex 3C-9 to 3C-12](#).

#### **Emission trends**

See [Annex 3C-13 to 3C-16](#).

### **4.6.1 Source specific recalculations and improvements**

No recalculations.

### **4.6.2 Source specific planned improvements**

No planned recalculations.

## **4.7 Other production**

### **4.7.1 Source category description**

The sub-sector *Other production* (NFR 2H) cover the following processes relevant for the Danish inventories: 2H2 Food and beverages industry

### **4.7.2 Food and beverages industry**

The following SNAP-codes are covered:

- 04 06 05 Bread
- 04 06 06 Wine
- 04 06 07 Beer
- 04 06 08 Spirits
- 04 06 25 Sugar production
- 04 06 27 Meat, fish etc. frying/curing
- 04 06 98 Margarine and solid cooking fats
- 04 06 99 Coffee roasting

The pollutant relevant for the lime production process is NMVOC.

#### **Methodology**

The emission of NMVOC from production of foods and alcoholic beverages is estimated from production statistics (Statistics Denmark, 2014), standard emission factors from the EMEP/EEA Guidebook (2013) and a country specific emission factor for sugar refining.

#### **Activity data**

The production statistics for the relevant processes have been aggregated based on data from Statistics Denmark and presented in Table 4.7.1. Data be-

fore 1985 are not available and therefore the activity level for 1980-1984 has been assumed to be equal to the average of 1985-1989.

Table 4.7.1 Production of foods and beverages (Statistics Denmark, 2014).

	Unit	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
Bread (rye and wheat)	Gg	181	181	181	181	181	193	178	179	176	178
Biscuits, cakes and other bakery products	Gg	105	105	105	105	105	119	99	101	101	103
Red wine	million l	12	12	12	12	12	12	20	12	8	8
White wine	million l	NO	NO	NO	NO	NO	NO	NO	NO	0.4	0.6
Beer	million l	886	886	886	886	886	836	879	875	916	922
Malt whisky	million l	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2
Other spirits	million l	33	33	33	33	33	39	33	31	30	31
Sugar production	Gg	506	506	506	506	506	533	532	522	468	476
Poultry curing	Gg	7	7	7	7	7	4	5	5	9	12
Fish and shellfish curing	Gg	37	37	37	37	37	35	32	36	38	42
Other meat curing	Gg	480	480	480	480	480	531	503	474	455	439
Margarine and solid cooking fats	Gg	247	247	247	247	247	233	241	260	254	248
Coffee roasting	Gg	53	53	53	53	53	53	53	53	51	54
<i>Continued</i>		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Bread (rye and wheat)	Gg	190	205	196	208	218	231	213	240	247	234
Biscuits, cakes and other bakery products	Gg	99	123	129	136	138	148	164	163	150	131
Red wine	million l	10	14	14	12	6	5	7	11	10	10
White wine	million l	3.2	6.6	6.7	5.9	0.1	0.5	0.6	3.5	6.2	6.7
Beer	million l	930	967	978	944	941	990	959	918	804	821
Malt whisky	million l	0.02	0.01	0.01	0.01	0.01	0.003	NO	NO	NO	NO
Other spirits	million l	33	33	36	37	30	27	29	29	28	23
Sugar production	Gg	506	510	471	484	496	444	432	487	557	535
Poultry curing	Gg	11	14	16	15	13	14	15	16	18	19
Fish and shellfish curing	Gg	52	48	52	46	40	31	40	36	33	35
Other meat curing	Gg	448	487	457	477	506	464	434	443	440	400
Margarine and solid cooking fats	Gg	233	237	238	221	201	199	193	217	227	205
Coffee roasting	Gg	52	51	56	55	56	49	55	53	55	61
<i>Continued</i>		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Bread (rye and wheat)	Gg	244	271	252	247	256	257	270	256	254	267
Biscuits, cakes and other bakery products	Gg	138	146	151	156	165	157	149	136	124	115
Red wine	million l	5	4	3	2	2	1	1	0	1	3
White wine	million l	0.3	2.4	2.5	2.2	2.6	3.1	2.1	1.7	1.1	1.2
Beer	million l	745	723	820	835	855	868	817	766	647	604
Malt whisky	million l	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Other spirits	million l	24	25	25	24	24	26	25	20	25	12
Sugar production	Gg	443	563	508	512	453	503	458	357	466	395
Poultry curing	Gg	24	30	32	33	33	35	38	39	40	50
Fish and shellfish curing	Gg	44	46	44	43	45	41	45	39	84	76
Other meat curing	Gg	393	390	385	400	357	361	342	318	310	307
Margarine and solid cooking fats	Gg	196	178	225	209	211	200	181	191	191	175
Coffee roasting	Gg	56	59	57	51	55	37	35	34	35	35
<i>Continued</i>		2010	2011	2012	2013						
Bread (rye and wheat)	Gg	245	207	233	227						
Biscuits, cakes and other bakery products	Gg	118	114	109	115						
Red wine	million l	4	3	2	4						
White wine	million l	4.6	3.6	5.5	8.2						
Beer	million l	634	659	608	617						
Malt whisky	million l	NO	NO	NO	NO						
Other spirits	million l	17	21	16	15						
Sugar production	Gg	262	218	262	493						
Poultry curing	Gg	54	57	63	65						
Fish and shellfish curing	Gg	73	63	62	67						
Other meat curing	Gg	303	307	249	241						
Margarine and solid cooking fats	Gg	180	191	182	146						
Coffee roasting	Gg	37	23	19	17						

NO: not occurring

### Emission factors

The emission factors used to calculate the NMVOC emissions from food and beverage production are shown in Table 4.7.2. Regarding refining of sugar, the default emission factor has been revised based on company specific measurements obtained from Nielsen (2011). TOC has been measured in order to solve odour issues. The emission of TOC has been used as indicator for NMVOC assuming a conversion factor at: 0.6 kg C/kg NMVOC.

Table 4.7.2 Emission factors for NMVOC for production of alcoholic beverages.

Production	Unit	Value	Reference
Bread (rye and wheat)	kg/Mg bread	4.5	EMEP/EEA Guidebook (2013)
Biscuits, cakes and other bakery products	kg/Mg product	1	EMEP/EEA Guidebook (2013)
Red wine	kg/m <sup>3</sup> wine	0.8	EMEP/EEA Guidebook (2013)
White wine	kg/m <sup>3</sup> wine	0.35	EMEP/EEA Guidebook (2013)
Beer	kg/m <sup>3</sup> beer	0.35	EMEP/EEA Guidebook (2013)
Malt whisky	kg/m <sup>3</sup> alcohol	150	EMEP/EEA Guidebook (2013)
Other spirits	kg/m <sup>3</sup> alcohol	4	EMEP/EEA Guidebook (2013)
Sugar production	kg/Mg sugar	0.2	Nielsen (2011)
Meat, fish and poultry	kg/Mg product	0.3	EMEP/EEA Guidebook (2013)
Margarine and solid cooking fats	kg/Mg product	10	EMEP/EEA Guidebook (2013)
Coffee roasting	kg/Mg beans	0.55	EMEP/EEA Guidebook (2013)

### Emission trends

The emission trend for emission of NMVOC from production of food and beverage is presented in Figure 4.7.1.




Figure 4.7.1 NMVOC emissions from the production of food and beverages, Mg.

The emission of NMVOC from production of food and beverage follows the activity as the same emission factors have been used for the entire period.

### 4.7.3 Source specific recalculations and improvements

New source categories that were added in this year's inventory include:

- Palmin (added to Margarine and solid cooking fats)

- Caffeine free coffee (added to coffee roasting)
- Wine (separated in red and white).

1980-1989 data was reassessed for all categories. As a result of the assessment 1980-1984 data were added to this year's inventory and 1985-1989 are now actual statistical data and not estimates.

An error was corrected for beer production resulting in an increase in activity data for 1999 of 18m litres.

The activity data for meat curing has been drastically reduced. Previously, the dataset included the entire meat production, but this has now been corrected to only meat that is fried, smoked or otherwise treated in ways that results in NMVOC emissions. The overall reduction for meat curing (including poultry and fish) is 82-89 %.

It was investigated whether it was possible to separate the production of spirits into categories depending on whether the spirit is matured or not. As a result, the inventory now separates between malt whisky and other spirits.

It was investigated whether the bread production could be subdivided to allow for the application of more disaggregated emission factors. This was however not possible with the available statistical data.

#### **4.7.4 Source specific planned improvements**

Other activities not currently included, such as flour production, grain drying and fish meal processing will be investigated further.

### **4.8 Other production, consumption, storage, transportation or handling of bulk products**

#### **4.8.1 Source category description**

The sub-sector *Other production, consumption, storage, transportation or handling of bulk products* (NFR 2L) covers the following SNAP-code: 04 06 17 Slaughterhouse waste.

#### **4.8.2 Slaughterhouse waste**

One company treats slaughterhouse waste: Daka with five departments located in Løsning, Randers, Lunderskov, Ortved, and Nyker. The pollutant relevant

#### **Methodology**

The raw materials for the processes are by-products from the slaughterhouse, animals dead from accident or disease, and blood. The outputs from the processes are protein and fat products as well as animal fat, meat and bone meal.

The emissions from the processes are related to the consumption of energy and emissions of e.g. NH<sub>3</sub> and odour. The last-mentioned emissions are related to storage of the raw materials as well as to the drying process.

The emission of NH<sub>3</sub> from treatment of slaughterhouse waste has been calculated from an average emission factor based on measurements from Danish plants (Daka, 2014).



### Activity data

The activity data for treatment of slaughterhouse waste are compiled from different sources. Due to changes in the company structure environmental reports are only available for some of the years (Daka, 2014). These environmental reports in combination with environmental reports for one of the merging companies are used to identify the corresponding data in the statistical information from Statistics Denmark (2014). The activity data are presented in Table 4.8.1.

Table 4.8.1 Activity data for treatment of slaughterhouse waste, Gg.

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
Meat/bone meal <sup>1</sup>	146.6	146.6	146.6	146.6	146.6	134.4	142.0	166.0	143.6	147.1
Animal fat <sup>1</sup>	20.1	20.1	20.1	20.1	20.1	11.1	14.3	15.0	41.6	18.6
Blood meal <sup>2</sup>	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0
Total	177.7	177.7	177.7	177.7	177.7	156.5	167.3	192.0	196.2	176.7
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Meat/bone meal <sup>1</sup>	128.8	186.1	181.0	209.2	197.8	197.0	191.3	166.6	177.7	201.5
Animal fat <sup>1</sup>	72.1	81.9	85.9	75.8	63.9	54.2	57.7	72.6	72.8	89.2
Blood meal <sup>2</sup>	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.4	11.5
Total	211.9	279.0	277.9	296.0	272.7	262.2	260.0	250.2	261.8	302.2
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Meat/bone meal <sup>1</sup>	198.6	154.0	209.1	227.8	241.6	177.4	161.7	142.6	140.5	116.4
Animal fat <sup>1</sup>	73.4	56.4	108.5	93.8	98.6	90.2	75.6	82.6	84.7	70.9
Blood meal <sup>2</sup>	11.4	9.7	8.9	9.5	10.6	10.2	8.9	10.6	10.0	7.5
Total	283.4	220.1	326.6	331.0	350.7	277.9	246.2	235.8	235.2	194.8
	2010	2011	2012	2013						
Meat/bone meal <sup>1</sup>	104.6	96.3	73.7	78.5						
Animal fat <sup>1</sup>	75.3	77.7	76.2	48.2						
Blood meal <sup>2</sup>	7.5	7.5	7.5	7.5						
Total	187.4	181.5	157.4	134.3						

<sup>1</sup> Statistics Denmark (2014).

<sup>2</sup> Based on environmental reports from Daka (2014) for the years 1998 – 2009 and assumed for the other years.

### Emission factors

The emission of NH<sub>3</sub> from treatment of slaughterhouse waste has been calculated from an average emission factor based on measurements from Danish plants (Daka, 2014). Measurements of NH<sub>3</sub> during the years 2002/3 from three locations (Lunderskov, Løsning and Randers) with different product mix have been included in the determination of an emission factor.

Weighted emission factors covering all the products within the sector have been estimated for 2002 and 2003 as well as a weighted emission factor covering 1990-2001. The estimated emission factors are presented in Table 4.8.2.

Table 4.8.2 Emission factors for treatment of slaughterhouse waste.

	EF	1990-2001	2002	2003-2012
NH <sub>3</sub>	g/Mg	120	151	475

### Emission trends

Emissions from the treatment of slaughterhouse waste are available in the NFR tables.

### 4.8.3 Source specific recalculations and improvements

The activity data for slaughter house waste was reassessed with the following results: The dataset was extended back to 1980, some increases in activity for the beginning of the 90's and a few changes for other years (2001, 2002, 2006 and 2012).

### 4.8.4 Source specific planned improvements

There are no planned improvements for this sector.

## 4.9 QA/QC and verification

Please refer to the Danish National Inventory Report reported to the UN-FCCC (Nielsen et al., 2014).

## 4.10 Uncertainty estimates

Table 4.10.1 Tier 1 uncertainties for Industrial Processes (NFR 2).

Pollutant	Uncertainty total	Trend 1990-2013	Uncertainty trend
	emission	(2000-2013)	
	%	%	%-age points
SO <sub>2</sub>	17,95	-45	2,7
NO <sub>x</sub>	40,81	-91	5,2
NMVOG	16,52	-32	8,8
CO	94,68	-72	24,9
NH <sub>3</sub>	539,80	-36	104,0
TSP	111,34	-48	26,2
PM <sub>10</sub>	127,74	-51	30,3
PM <sub>2.5</sub>	147,35	-50	34,5
BC	50,04	-27	2,1
Arsenic	642,06	-61	190,5
Cadmium	604,54	-71	80,5
Chromium	534,86	-38	162,5
Copper	149,06	22	162,8
Mercury	901,08	-96	3,1
Nickel	499,57	-72	119,3
Lead	629,58	-85	40,8
Selenium	933,15	-46	204,1
Zinc	336,22	-81	143,5
Dioxins/Furans	544,82	-99	5,1
benzo(b)flouranthene	150,75	92	40,6
benzo(k)flouranthene	150,75	89	40,1
benzo(a)pyrene	150,75	86	39,4
indeno(1,2,3-c,d)pyrene	150,75	90	40,2
HCB	963,65	-95	56,0
PCB	585,76	-95	23,5

Important uncertainty issues related to the mass-balance approach used for Non-energy products from fuels and solvent use (NFR 2D) are:

(i) Identification of pollutants that qualify as NMVOGs. Although a tentative list of 650 pollutants from NAEI (2000) has been used, it is possible that relevant pollutants are not included, e.g. pollutants that are not listed with their name in Statistics Denmark (2014) but as a product.

(ii) Collection of data for quantifying production, import and export of single pollutants and products where the pollutants are comprised. For some

pollutants no data are available in Statistics Denmark (2014). This can be due to confidentiality or that the amount of pollutants must be derived from products wherein they are comprised. For other pollutants the amount is the sum of the single pollutants *and* product(s) where they are included. The data available in Statistics Denmark (2014) is obtained from Danish Customs & Tax Authorities and they have not been verified in this assessment.

(iii) Distribution of pollutants on products, activities, sectors and households. The present approach is based on amounts of single pollutants. To differentiate the amounts into industrial sectors it is necessary to identify and quantify the associated products and activities and assign these to the industrial sectors and households. No direct link is available between the amounts of pollutants and products or activities. From the Nordic SPIN database it is possible to make a relative quantification of products and activities used in industry, and combined with estimates and expert judgement these products and activities are differentiated into sectors. The contribution from households is also based on estimates. If the household contribution is set too low, the emission from industrial sectors will be too high and vice versa. This is due to the fact that the total amount of pollutant is constant. A change in distribution of pollutants between industrial sectors and households will, however, affect the total emissions, as different emission factors are applied in industry and households, respectively.

A number of activities are assigned as "other", i.e. activities that cannot be related to the comprised source categories. This assignment is based on expert judgement but it is possible that the assigned amount of pollutants may more correctly be included in other sectors. More detailed information from the industrial sectors is continuously being implemented.

(iv) Rough estimates and assumed emission factors are used for some pollutants. For some pollutants more reliable information has been obtained from the literature and from communication with industrial sectors. In some cases it is more appropriate to define emission factors for sector specific activities rather than for the individual pollutants.

A quantitative measure of the uncertainty has not been assessed. Single values have been used for emission factors and activity distribution ratios etc.

The main issues leading to uncertainties for Other Product Use (NFR 2G) are:

(i) Collection of data for quantifying production, import and export of products. Some data, like private import (cross-border shopping) of fireworks, are not available in Statistics Denmark (2014). Other lacking data like the composition of mineral containing charcoal for barbequing are unobtainable due to confidentiality.

(ii) Reliable emission factors are not easy to obtain for other product use categories. Some chosen emission factors apply to countries that are not directly comparable to Denmark, and hereby is introduced an increased uncertainty. This is the case with e.g. some heavy metals from charcoal for barbequing.

## 4.11 References

Asfaltindustrien, 2003: Asfalt 3/03, Asfalt vore veje. Direktion, Ib Frandsen.

Ardagh Glass Holmegaard, 2014: Grønt regnskab, CO<sub>2</sub>-opgørelse og afrapportering 2013. Ardagh Glass Holmegaard A/S, CVR nr. 18445042 (In Danish) Partly confidential. (Environmental reports).

CEPMEIP: The Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance (CEPMEIP). Database. Available at: <http://www.air.sk/tno/cepmeip/> (14/01-2015).

Cheminova, 2014: Grønt regnskab 2013 for Cheminova A/S. (Environmental reports).

Daka, 2014: Grønt regnskab 2013. (Environmental reports).

Danisco Grindsted, 2014: Grønt regnskab 2013. (Environmental reports).

DanSteel, 2014: NLMK DanSteel A/S 2014, Grønt Regnskab 2013. (Environmental reports).

Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations, Brüssel, 1999.

EMEP/EEA Guidebook 2013: EMEP/EEA air pollutant emission inventory guidebook 2013. Published by the EEA with the CLRTAP Task Force on Emission Inventories and Projections responsible for the technical content of the chapters. Technical report No 12/2013. Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (23/01-2015).

Fausser, P., Saarinen, K., Harðardóttir, K., Kittilsen, M.O., Holmengen, N. & Skårman, T., 2009: Improvement of Nordic Emission Models for Solvent Use in Selected Sectors. TemaNord 2009:556. Nordic Council Of Ministers, Copenhagen.

Haldor Topsøe, 2014: Miljøredegørelse for katalysatorfabrikken 2013 (16. regnskabsår) (Environmental reports).

Hals metal, 2014: Personal communication with Helle Bjerrum Holm September 2014.

Illerup, 1999: Illerup, J.B., Geertinger A.M., Hoffmann, L. & Christiansen, K., 1999: Emissionsfaktorer for tungmetaller 1990-1996. Faglig rapport fra DMU, nr. 301. Miljø- og Energiministeriet, Danmarks Miljøundersøgelse. Available at: [http://www2.dmu.dk/1\\_viden/2\\_publicationer/3\\_fagrapporter/rapporter/fr301.pdf](http://www2.dmu.dk/1_viden/2_publicationer/3_fagrapporter/rapporter/fr301.pdf) (in Danish) (14/01-2015).

IPCC, 2006: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Available at: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/> (23/01-2015).

Jensen & Markussen, 1993: Jensen, A., & Markussen, J. 1993. Forbrug af og forurening med cadmium (Miljøprojekt nr. 213). København: Miljøstyrelsen.

- Jydsk Aluminiums Industri, 2014: Grønt Regnskab 2012 A/S Jydsk Aluminium Industri. (Environmental reports).
- Karjalainen, T. 2005: Commission research in Action: tackling the hormone disrupting chemicals issue, EUR report 21941.
- Kemira GrowHow, 2004: Miljø & arbejdsmiljø. Grønt regnskab 2003. (Environmental reports).
- Kemira GrowHow, 2005: Personal communication with Gert Jacobsen, September 2005.
- Koppers, 2014: Miljøreddegørelse 2013. Koppers European Operations, Carbon Materials and Chemicals, Nyborg Works, Denmark, 2014. (Environmental reports).
- Lassen, C, Christensen, C.L. & Skårup, S. 2004: Cowi A/S, Massestrømsanalyse for bly 2000 - revideret udgave, Miljøprojekt Nr. 917 2004. (Danish). Available at:  
[http://www.unep.org/chemicalsandwaste/Portals/9/Lead\\_Cadmium/docs/submissions/Massestromsanalyse\\_for\\_bly\\_2000\\_revised\\_Nr\\_917.pdf](http://www.unep.org/chemicalsandwaste/Portals/9/Lead_Cadmium/docs/submissions/Massestromsanalyse_for_bly_2000_revised_Nr_917.pdf) (29/01-2015)
- NAEI, 2000: British National Atmospheric Inventory performed by the National Environmental Technology Centre, June 2000. Available at:  
[http://www.aeat.co.uk/netcen/airqual/naei/annreport/annrep99/app1\\_2\\_8.html](http://www.aeat.co.uk/netcen/airqual/naei/annreport/annrep99/app1_2_8.html) (05/05-2006).
- Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Fauser, P., Mikkelsen, M.H., Albrektsen, R., Hjelgaard, K., Hoffmann, L., Thomsen, M., Bruun, H.G., 2013: Danish emission inventory for hexachlorobenzene and polychlorinated biphenyls. Aarhus University, DCE – Danish Centre for Environment and Energy, 65 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 103. Available at:  
<http://www.dce2.au.dk/pub/SR103.pdf> (13/01-2015).
- Nielsen, O.-K., Plejdrup, M.S., Winther, M., Nielsen, M., Gyldenkerne, S., Mikkelsen, M.H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H.G., Johannsen, V.K., Nord-Larsen, T., Vesterdal, L., Møller, I.S., Caspersen, O.H., Rasmussen, E., Petersen, S.B., Baunbæk, L. & Hansen, M.G. 2014: Denmark's National Inventory Report 2014. Emission Inventories 1990-2012 - Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol. Aarhus University, DCE – Danish Centre for Environment and Energy, 1214pp. Scientific Report from DCE – Danish Centre for Environment and Energy. Available at:  
<http://www.dce2.au.dk/pub/SR101.pdf> (23/01-2015).
- Nielsen, 2011: Personal communication Vibeke Vestergaard Nielsen, Danish EPA, 9 September 2011.
- Pärt, P. (main author) 2005: Environment and health, EEA Report No 10/2005, Copenhagen.
- Rockwool, 2014: Miljøreddegørelse 2013 for fabrikkerne i Hedehusene, Vamdrup og Øster Doense.

Saint-Gobain Isover, 2014: Redegørelse for arbejdsmiljø, miljø, sundhed og socialt engagement 2013. (Environmental reports).

SPIN on the Internet. Substances in Preparations in Nordic Countries. Available at: <http://www.spin2000.net/spin.html>.

Statistics Denmark, 2014: Statistics Denmark, StatBank Denmark (Danish/English). Available at: <http://www.dst.dk/> (15/01-2015).

Stena Aluminium, 2008: Miljøredogørelse/Grønt Regnskab for Stena Aluminium A/S 2007/2008. (Environmental reports).

Stålvalseværket, 2002: Grønt regnskab og miljøredogørelse 2001. Det Danske Stålvalseværk A/S. (Environmental reports).

UNFCCC, 2008: Available at: [http://unfccc.int/national\\_reports/annex\\_i\\_ghg\\_inventories/national\\_inventories\\_submissions/items/3929.php](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/3929.php) (29/01-2015).

US EPA, 2004: Technology Transfer Network Clearinghouse for Inventories & Emissions Factors. AP 42, Fifth Edition, Volume I, Chapter 11: Mineral Products Industry. Available at: <http://www.epa.gov/ttnchie1/ap42/ch11/> (29/01-2015).

## 5 Agriculture (NFR sector 3)

### 5.1 Overview of the sector

The emission from agricultural activities covers ammonia (NH<sub>3</sub>) emission from manure management, agricultural soils, field burning and agricultural other. Emission of particulate matter (PM) from manure management, field operations and field burning. Nitrogen oxides (NO<sub>x</sub>) and non-methane volatile organic compounds (NMVOC) emission from manure management, agricultural soils and field burning. Emission of hexachlorbenzen (HCB) from use of pesticides. Furthermore emission of carbon monoxide (CO), sulphur dioxide (SO<sub>2</sub>), black carbon (BC), heavy metals (As, Cd, Cr, Cu, Hg, Ni, Pb, Se and Zn), dioxins and furanes (PCDD/F) and polycyclic aromatic hydrocarbons (PAH – benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene and indeno(1,2,3-cd)pyrene), HCB and perchlorbenzen (PCB) from field burning.

Buffalos, camels, lamas, mules and donkeys are not farmed in Denmark therefore no emission estimates from these animal categories.

Table 5.1 shows the agricultural contribution of emissions to the national total in 2013. The main part of the NH<sub>3</sub> emission (95 %) is related to the agricultural sector, while the agricultural contribution of TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are 74 %, 35 % and 9 %, respectively. Due to implementation of NMVOC emission from manure management, the agricultural share now accounts for 33 % of the total. This year the inventory also includes the NO<sub>x</sub> emission from application of inorganic fertilizers and animal manure, which result in an agricultural part on 12 % of the total. The agricultural part of the total emissions of SO<sub>x</sub> is lower than 1 %.

Table 5.1 Emission 2013, Agricultural share of the Danish emission.

	NH <sub>3</sub>	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>	NMVOC	SO <sub>x</sub>	NO <sub>x</sub>
National total, Gg	74	89	32	21	114	14	124
Agricultural total, Gg	71	66	11	2	38	<1	14
Agricultural part, %	95	74	35	9	33	<1	12

#### 5.1.1 Ammonia

The majority of the Danish NH<sub>3</sub> emission, corresponding to 95 %, originates from the agricultural sector. The remaining 5 % originates from stationary combustion, traffic and industrial processes. Figure 5.1 shows the distribution of sources of NH<sub>3</sub> emission from the agricultural sector 2013. The main part of the agricultural emission is directly related to the livestock production by 51 % from manure management, 28 % from manure applied to soils and 3% from grazing animals. Emissions from use of inorganic fertilizer and cultivated crops contribute with 10 % and 7 %, respectively. Emissions from NH<sub>3</sub>-treated straw, field burning of agricultural residues and sewage sludge used as fertilizer amount to less than 1 %.




Figure 5.1 NH<sub>3</sub> emissions from the agricultural sector, 2013.

The NH<sub>3</sub> emission from the agricultural sector has decreased between 1985 and 2013 from 125.97 Gg NH<sub>3</sub> to 70.53 Gg NH<sub>3</sub>, corresponding to a 44 % reduction (Table 5.2). This significant drop in NH<sub>3</sub> emissions should be read in a conjunction of a very active national environmental policy designed to reduce the loss of nitrogen to the aquatic environment. A string of measures have been introduced by action plans, for example the NPO (Nitrogen, phosphor, organic matter) Action Plan (1986), Action Plans for the Aquatic Environment (1987, 1998, 2004), the Action Plan for Sustainable Agriculture (1991), the Ammonia Action Plan (2001) and latest the action plan the Agreement on Green Growth (2009 and 2010). Based on these action plans have legislative changes and actions led to an optimization of manure as a resource. Requirements to capacity of slurry storage and requirements to handling of manure during spreading has led to a decrease in animal nitrogen excretion, improvement in use of nitrogen in manure and a fall in the use of inorganic fertilizer. A Danish environmental approval act for livestock holdings was acted in January 2007 and according to the act, farmers are required to apply for an environmental approval if the farmer wants to change or expand the livestock production facilities. In order to get environmental approval farmers has to fulfil requirements concerning Best Available Technique (BAT) and specific environmental requirements as for example emission of ammonia. All of these action plans have helped to reduce the overall NH<sub>3</sub> emission significantly and the new Danish environmental approval act for livestock will contribute to a further reduction in emissions in future.

Table 5.2 Total NH<sub>3</sub> emissions from the agricultural sector 1985 to 2013, Gg NH<sub>3</sub>.

		1985	1990	1995	2000	2005	2010	2011	2012	2013
3B	Manure Management, total	53.33	50.95	46.77	46.85	47.16	40.02	39.97	39.29	36.31
	<i>Cattle</i>	13.81	12.62	11.54	11.84	10.47	10.72	11.06	11.20	11.23
	<i>Swine</i>	31.55	28.86	26.09	24.99	24.71	17.91	17.81	16.94	16.23
	<i>Other animals</i>	7.98	9.47	9.14	10.02	11.97	11.39	11.11	11.15	8.86
3Da1	Inorganic N-fertilizer	15.66	16.87	15.27	9.90	7.81	7.14	7.36	7.06	7.29
3Da2a	Manure applied to soil	39.75	36.72	29.67	25.61	20.89	20.70	19.78	19.55	19.53
3Da2b	Sewage sludge applied to soil	0.05	0.07	0.11	0.08	0.05	0.06	0.06	0.06	0.06
3Da3	Urine and dung deposite by grazing animals	3.12	2.91	3.02	2.92	2.21	1.87	1.81	1.84	1.86
3De	Cultivated crops	5.97	5.92	5.28	5.21	5.34	5.41	5.42	5.40	5.37
3F	Field burning of Agricultural Residue	1.53	0.08	0.09	0.11	0.13	0.09	0.09	0.10	0.11
3I	NH <sub>3</sub> treated straw	6.54	10.19	6.63	2.47	0.26	0.24	0.24	NO	NO
3.	Agricultural Sector - total	125.97	123.71	106.83	93.15	83.84	75.52	74.72	73.30	70.53



The management of manure has to be considered as the most important emission source. Most of the emission originates from the production of swine and cattle, which contributed, respectively with 45 % and 31 %.

It is noteworthy that the overall emission from swine has decreased by 49 % from 1985 to 2013 despite a considerable increase in swine production from 14.7 million produced fattening pigs in 1985 to 20.2 million in 2013. The most important reason for this is the improvement in feed efficiency. In 1985, the nitrogen excretion for a fattening pig was estimated to 5.09 kg N (Poulsen & Kristensen, 1998). In 2013, that figures were considerably lower at 2.85 kg N per fattening pig produced (Poulsen, 2014). Due to the large contribution from the swine production, the lower level of N-excretion has a significant influence on total agricultural emissions.

Since 1985, changes in practice of manure application to the fields have taken place, which has reduced the emission from manure applied to soils. From the beginning of the 1990s slurry has increasingly been spread using trailing hoses. From the late 1990s the practice of slurry injection or mechanical incorporation into the soil has increased. This development is a consequence of a ban on broad spreading but it is also a consequence of the general requirement to improve the utilisation of nitrogen in the manure - e.g. requirements that a larger part of the nitrogen in manure has to be included in the farmer's nitrogen accounting. This has forced farmers to consider the manure as a fertilizer resource instead of a waste product.

### **5.1.2 Particulate matter**

In the NFR, the emission of particulate matter (PM) is reported for the years 2000 to 2013. The emission from the agricultural sector includes the emission of dust from animal housing systems, field operations and field burning of agricultural residues.

TSP (total suspended particulate) emission from the agricultural sector contributes with 73 % to the national TSP emission in 2013 and the emission shares for PM<sub>10</sub> and PM<sub>2.5</sub> are 34 % and 8 % respectively. The majority of the TSP emission originates from the field operations 85 % while the emission from animal housings contributes with 15 % and field burning of agricultural residues, contributes with less than 1 % to the agricultural emission in 2013.

The PM emission from agricultural activities, given in TSP, is decreased 7 % during the period from 2000 to 2013 (Figure 5.2) mainly to decrease in the emission from field operations.




Figure 5.2 Emission of particulate matter (TSP) from the agricultural sector 2000 to 2013.

## 5.2 References – sources of information

DCE – the Danish Centre for Environment and Energy, Aarhus University, which is responsible for the emission inventory, has established data agreements with the institutes and organisations to assure that the necessary data are available for timely completion of the emission inventory. The main part of the emission is related to livestock production and most of the data are based on Danish standards.

Activity data, emissions factors (EF) and additional values are collected, evaluated and discussed in cooperation with Statistics Denmark, DCA - Danish Centre for Food and Agriculture, Aarhus University, the Danish Agricultural Advisory Service, Danish Environmental Protection Agency and the Danish AgriFish Agency. It means that both the data and the methods used are evaluated continuously according to latest knowledge and information. Table 5.3 shows the source of data input from the different institutes.

Table 5.3 List of institutes involved in the emission inventory.

References	Abbreviation	Data / information
Statistics Denmark - Agricultural Statistics ( <a href="http://www.dst.dk/en.aspx">http://www.dst.dk/en.aspx</a> )	DSt	-livestock production -milk yield -slaughtering data -land use -crop production -crop yield
Danish Centre for Food and Agriculture, Aarhus University	DCA	-N-excretion -feeding situation -NH <sub>3</sub> emissions factor -PM emissions factor
The Danish Agricultural Advisory Service ( <a href="http://www.vfl.dk">http://www.vfl.dk</a> )	DAAS	-housing type (until 2004) -grazing situation -manure application time and methods -field burning of agricultural residue
Danish Environmental Protection Agency ( <a href="http://www.mst.dk">http://www.mst.dk</a> )	EPA	-sewage sludge used as fertilizer (until 2004)
The Danish AgriFish Agency ( <a href="http://naturehverv.fvm.dk">http://naturehverv.fvm.dk</a> )	DAFA	-inorganic fertilizer -number of animals from CHR -housing type (from 2005) -sewage sludge used as fertilizer(from 2005)

### 5.2.1 Methods

The emission calculation is based on the methodologies provided in the EMEP/EEA air pollutant emission inventory guidebook (EMEP/EEA, 2013).

The emissions from agricultural activities include Animal Husbandry and Manure Management, Crop Production and Agricultural Soils, Field Burning of Agricultural Residue, Use of Pesticides and Agriculture Other. In general the field burning of agricultural residue has been prohibited since 1989. However, burning of straw may take place in connection with fields continuously cultivating seed grass or in cases where weather conditions result in surplus of straw in form of wet or broken bales.

The emissions from the agricultural sector are calculated in a comprehensive agricultural model complex called IDA (Integrated Database model for Agricultural emissions). The model complex is designed in a relational database system (MS Access). Input data are stored in tables in one database called IDA\_Backend and the calculations are carried out as queries in another linked database called IDA. The model, as shown in Figure 5.3, is implemented and it is used to calculate emissions of NH<sub>3</sub>, PM, NO<sub>x</sub>, CO, NMVOC, SO<sub>2</sub>, heavy metals, dioxin, PAH, HCB, PCB and greenhouse gases (N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub>). Thus, there is direct link between the NH<sub>3</sub> emission and the emission estimation of N<sub>2</sub>O.

DCA, Danish Centre for Food and Agriculture, Aarhus University delivers Danish standards relating to feeding consumption, manure type in different housing types, nitrogen content in manure, etc. Previously, the standards were updated and published every third or fourth year – the last one is Poulsen et al. from 2001. From year 2001, DCE receives updated data annually directly from DCA in the form of spread sheets. These standards have been described and published in English in Poulsen & Kristensen (1998). From 2004 the standards are uploaded every year at <http://anis.au.dk/forskning/sektioner/husdyrernaering-og-miljoe/normtal/>.

## IDA - Integrated Database model for Agricultural emissions




Figure 5.3 Overview of the data process for calculation of agricultural emissions.

IDA includes 39 different livestock categories, divided on weight class and age. Each of these subcategories is subdivided according to housing system and manure type, which results in 266 different combinations of subcategories and housing type (Table 5.4). The emissions are calculated from each of these subcategories and then aggregated in accordance with the livestock categories given in the NFR. It is important to point out that changes in the emission and the implied emission factor over the years are not only a result of changes in the number of animals, but also depend on changes in the allocation of subcategories, changes in feed consumption, changes in housing type and changed practices with regard to the handling of livestock manure in relation to storage and application.

Table 5.4 Livestock categories and subcategories.

NFR	Animal categories	Includes	No. of sub-categories in IDA, animal type/housing system/manure type
3B 1a	Dairy Cattle <sup>1</sup>	Dairy Cattle	35
3B 1b	Non-dairy Cattle <sup>1</sup>	Calves (<½ year), heifers, bulls, suckling cattle	129
3B 2	Sheep	Sheep and lambs	2
3B 3	Swine	Sows, weaners, fattening pigs	37
3B 4d	Goats	Including kids (meet, dairy and mohair)	3
3B 4e	Horses	<300 kg, 300 - 499 kg, 500 - 700 kg, >700 kg	4
3B 4gl-gIV	Poultry	Hens, pullet, broilers, turkey, geese, ducks, ostrich, pheasant	47
3B 4h	Other	Fur bearing animals, deer	9

<sup>1)</sup> For all cattle categories, large breed and jersey cattle are distinguished from each other.

### 5.3 Manure management

For the sector manure management is the emissions of NH<sub>3</sub>, PM, NMVOC and NO<sub>x</sub> estimated.

#### 5.3.1 Activity data

##### Animals

Table 5.5 shows the development in livestock production from 1985 to 2013 based on the Agricultural Statistics (Statistics Denmark). The number of animal corresponds to average annual production (AAP), which means the number of animals that are present on average within the year (EMEP/EEA, 2013). For many animal categories the number given in the annual Agricultural Statistics can be used directly. However, for weaners, fattening pigs, bulls and poultry the number is based on slaughter data also collected from the Agricultural Statistics. This is because the production cycle for these animals is under one year and the normative figures are based on produced animals. See Annex 3D Table 3D-1 for number of animals allocated on subcategories.

Only farms larger than five hectares are included in the annual census. Especially horses, goats and sheep are placed on small farms, which mean that the number of animals given in the Agricultural Statistics is not representative. Therefore, the number of sheep and goats is based on the Central Husbandry Register (CHR) which is the central register of farms and animals managed by the Ministry of Food, Agriculture and Fisheries. The number of deer and ostriches is also based on CHR because these are not included in the Agricultural Statistics published by Statistics Denmark. The number of horses is based on data from The Danish Agricultural Advisory Service. The number of pheasants is based on expert judgement from Department of Bioscience, Aarhus University and the Danish pheasant breeding association.

Since 1985, the production of swine, poultry and fur has increased significantly. This is contrary to the production of cattle, which has decreased as a result of increasing milk yields. Buffalos, camels, lamas, mules and donkeys are not farmed in Denmark.

Table 5.5 Livestock production 1985 to 2013 given in AAP, 1000 head - NFR category 3B.

NFR	Animal category	1985	1990	1995	2000	2005	2010	2011	2012	2013
3B 1a	Dairy Cattle	896	753	702	636	564	568	565	587	582
3B 1b	Non-dairy cattle	1 721	1 486	1 388	1 232	1 006	1 003	1 003	1 020	1 032
3B 2	Sheep*	99	230	202	279	316	278	234	226	221
3B 3	Swine	9 089	9 497	11 084	11 922	13 534	13 173	12 932	12 331	12 076
3B 4d	Goats*	8	7	7	8	11	16	13	13	13
3B 4e	Horses*	140	135	143	150	175	165	155	155	150
3B 4gl	Laying hens	5 577	5 696	6 088	4 935	5 168	5 248	5 679	5 597	5 766
3B 4gll	Broilers	8 490	9 802	12 585	16 047	11 905	12 836	12 528	12 576	13 215
3B 4glll	Turkeys	308	238	456	456	516	494	400	460	289
3B 4glV	Other poultry	1 822	1 600	1 563	1 374	1 509	1 510	1 963	1 444	1 263
3B 4h	Other									
3B 4h	Fur bearing animals	1 906	2 264	1 850	2 199	2 552	2 699	2 757	2 948	3 143
3B 4h	Deer	9	10	10	10	10	10	8	7	8

\*Includes animals on small farms (less than 5 ha), which are not included in the Agricultural Statistics published by Statistics Denmark.

See Annex 3D Table 3D-1 for number of animals allocated on subcategories.

### N-excretion

The normative figures for both total nitrogen excretion and the content of TAN are provided by DCA, Aarhus University.

The emission of NH<sub>3</sub> from manure management is calculated on the basis on nitrogen excreted from livestock. Most of the N excreted that is readily degradable and broken down to NH<sub>4</sub>-N is found in the urine. The relationship between NH<sub>4</sub>-N and total N will not remain constant over time due to changes in feed composition and feed use efficiency. In order to be able to implement the effect of NH<sub>3</sub> reducing measures as improvements in feed intake and composition in the emission inventory, it is necessary to calculate the emission based on the Total Ammoniacal Nitrogen (TAN) content. Since 2007, DCA has established Danish standards based on TAN for liquid manure, which is incorporated in the inventory. The emission for solid manure and deep litter is based on the total N excreted because DCA's estimate of TAN follows urine-N.

In Annex 3D Table 3D.2 is given the average N-excretion based on Total-N for each NFR livestock category from 1985 to 2006 (Table 3D.2a) and N-excretion based on TAN for 2007-2013 (Table 3D.2b). These values include N excretion from grazing animals. Notice that each livestock category is an aggregated average of different subcategories (see Table 5.4).

### Housing system

A systematic registration of the housing of husbandry for all farms does not exist from 1985 to 2004 and the housing type distribution is therefore based on estimates from the Danish Agricultural Advisory Services (Rasmussen, 2006) and Lundgaard (2006). From 2005 the distribution of housing system is based on information from the Danish AgriFish Agency, which is based on information from the farmers.

The structural development in the agricultural sector has an influence on the changes in housing type distribution. The trend in housing system for dairy cattle goes from older tied-up housings, which is replaced by bigger housings with loose-holding. In 1985 85 % of the dairy cattle were kept in tied-up housings and in 2013 the share is reduced to 8 %. In loose-holding systems the cattle have more space and more straw bedding and this will

in general increase the NH<sub>3</sub> emission per animal compared to the tied-up housings. In Annex 3D Table 3D.3 the distribution of housing type for all animals for 1985-2013 is listed.

### 5.3.2 NH<sub>3</sub>

#### Description

The main part of the NH<sub>3</sub> emission (51 %) is related to manure management. Figure 5.4 shows the emission from manure management (NFR category 3.B) distributed according to the different livestock categories. The main part of the emission is related to cattle and swine production, corresponding to 76 %. The fall in the emission from swine 1985 to 2013 is mainly due to an active environmental policy in combination with improvements within the genetic development. The emission has increased slightly from "other", which is mainly due to an increase in number of produced mink.




Figure 5.4 NH<sub>3</sub> emission from manure management 1985 to 2013.

#### Methodological issues

NH<sub>3</sub> emission from manure management covers emission from housings and storage and is based on N excreted and emission factors given in the normative figures (Poulsen et al, 2001, and Poulsen 2014).

#### Activity data

See Chapter 5.2.1

#### Emission factor

Emission factors - Housing

The emission from housings is thus determined by a number of different conditions that depends on housing type and the different kinds of manure disposal systems placed in these housings. Danish Centre for Food and Agriculture, Aarhus University has carried out a number of emission surveys and estimated emission coefficients for each type of housings (Poulsen et al., 2001 and Poulsen, 2014). In Table 5.6 is shown the emission factors for the most important animal categories; dairy cattle and fattening pigs in different housing systems. For the slurry and liquid manure is giv-

en TAN emission factors (TAN ex animal) and for solid and deep litter manure is given N ex animal.

Table 5.6 NH<sub>3</sub> emission factors in different housing systems 2013 – dairy cattle and fattening pigs.

Manure system	Manure type	NH <sub>3</sub> emission	
		Pct. NH <sub>3</sub> -N of N ex Animal	Pct. NH <sub>3</sub> -N of TAN ex Animal
<b>Dairy cattle</b>			
Tied-up	Solid manure	6.0	
	+ Liquid		10.0
Tied-up	Slurry		6.0
Loose-holding with beds, slatted floor	Slurry		16.0
Loose-holding with beds, slatted floor, scrapes	Slurry		12.0
Loose-holding with beds, solid floor	Slurry		20.0
Loose-holding with beds, drained floor	Slurry		8.0
Deep litter (all)	Deep litter	6.0	
Deep litter, slatted floor	Deep litter	6.0	
	+ Slurry		16.0
Deep litter, slatted floor, scrapes	Deep litter	6.0	
	+ Slurry		12.0
Deep litter, solid floor, scrapes	Deep litter	6.0	
	+ Slurry		20.0
<b>Fattening pigs</b>			
Full slatted floor	Slurry		24.0
Partly slatted floor (50-75% solid floor)	Slurry		13.0
Partly slatted floor (25-49% solid floor)	Slurry		17.0
Solid floor	Solid manure	15.0	
	+ Liquid		27.0
Deep litter	Deep litter	15.0	
Partly slatted floor and partly deep litter	Deep litter	15.0	
	+ Slurry		18.0

#### Emission factors - Storage

Livestock manure is collected either as solid manure or as slurry depending on housing type. In Table 5.7 are shown the emission factors used for storage. It is assumed that the part of solid manure taken directly from the housing into the field is 65 % from cattle, 25 % from pigs, 50 % from sows, 15 % from poultry and 5 % from hens (Poulsen, 2008). The remaining part of the solid manure is deposited in stock piles in the field before field application.

By law all slurry tanks have to be covered by a fixed cover or a full floating cover in order to reduce NH<sub>3</sub> emission. However, it can be difficult to establish a natural full floating cover every day all year especially for tank with pig slurry. In 2013 it is assumed that 5 % of the tanks with swine slurry and 2 % of tanks with cattle slurry are incompletely covered (Annex 3D Table 3D-4).



Table 5.7 NH<sub>3</sub> emission factors for storage 2013.

Animal category	Liquid manure		Slurry	Solid manure	Deep litter	
	Loss of NH <sub>3</sub> -N in %					
	of TAN ex housing	of TAN ex housing	of N ex housing	of N ex housing	of N ex housing	
Cattle	2.2		3.5	4.0	1.05	
Swine		Fattening pigs	2.2	2.9	19.0	9.75
		Sows		2.9	19.0	6.50
Poultry		Hens and pullet		2.0 <sup>a</sup>	7.5	4.75
		Broilers, geese and ducks			7.5	6.80
		Turkeys			7.5	8.00
Fur bearing animals			3.1	11.5		
Sheep/goats						3.0
Horses						3.0

<sup>a</sup> Loss of NH<sub>3</sub>-N in % of N ex housing.

### Implied emission factor

Table 5.8 shows the implied emission factors for each NFR livestock category from 1985 to 2013. The implied emission factors express the average emission of NH<sub>3</sub> from housing and storage per AAP (annual average population) per year. The implied emission factors are changing from year to year depending on a combination of several factors, such as:

- change in number of animals or change in the share of different subcategories,
- change in feed intake and N-excretion,
- change in housing type,

It should be mentioned that the emission from urine and dung deposited by grazing animals is included in the emission from agricultural soils (NFR – 3Da3).

For most of the animal categories are the implied emission factor decreased from 1985 to 2013, which is mainly the result of measures in relation to the environmental Action Plans. Strict requirements to obtain improvements in utilisation of nitrogen in manure have resulted in reduction of N-excretion and especially for fattening pigs. For dairy cattle the implied emission factor has increased and this is due to increase in feed intake and milk production per cow.

Table 5.8 Implied emission factor, manure management 1985 to 2013, kg NH<sub>3</sub> per AAP per year.

NFR	Animal category	1985	1990	1995	2000	2005	2010	2011	2012	2013
3B 1a	Dairy cattle	9.64	10.50	10.37	11.49	13.50	12.86	13.32	13.07	13.06
3B 1b	Non-dairy cattle	3.00	3.18	3.06	3.68	2.84	3.40	3.52	3.46	3.52
3B 2	Sheep	0.51	0.52	0.53	0.44	0.44	0.40	0.40	0.40	0.40
3B 3	Swine	3.47	3.04	2.35	2.10	1.83	1.36	1.38	1.37	1.34
3B 4d	Goats	1.39	1.40	1.44	1.12	1.05	0.98	0.98	0.99	0.99
3B 4e	Horses	5.44	5.34	4.80	4.84	4.84	4.34	4.34	4.34	4.34
3B 4gl	Laying hens	0.15	0.20	0.25	0.27	0.34	0.27	0.27	0.24	0.24
3B 4gII	Broilers	0.15	0.20	0.18	0.17	0.21	0.15	0.14	0.13	0.07
3B 4gIII	Turkeys	0.49	0.51	0.65	0.63	0.63	0.52	0.52	0.52	0.52
3B 4glV	Other poultry	0.10	0.10	0.14	0.10	0.08	0.03	0.02	0.03	0.02
3B 4h	Other	2.46	2.28	2.15	2.13	2.44	2.55	2.47	2.39	1.80

## Emissions

The NH<sub>3</sub> emission from manure management is estimated to 36.31 Gg NH<sub>3</sub> in 2013 (Table 5.9). From 1985 to 2013, the emission is reduced by 32 %. As mentioned in Chapter 6.1.1 this development is mainly due to implementation of a number of action plans to reduce nitrogen losses from the agricultural production.

In 2013, cattle production contributes with 31 % of the total emission from manure management. The swine production contributes in 2013 with 45 % of the total emission from manure management. The number of cattle has decreased as a result of a growth in milk yield and the milk quota which limits the production. The production of fattening pigs has increased by more than 50 % compared with 1985. However, despite this development the emission from swine is still decreasing. This is due to a breeding of pigs with focus on a biological development and improvement in fodder efficiency. Thus the N-excretion for fattening pigs has decreased from 5.09 kg per pig per year in 1985 to 2.85 in 2013.

Table 5.9 Emission of NH<sub>3</sub> from manure management 1985 to 2013, Gg NH<sub>3</sub>.

NFR	Animal category	1985	1990	1995	2000	2005	2010	2011	2012	2013
3B 1a	Dairy cattle	8.64	7.90	7.29	7.30	7.62	7.31	7.53	7.67	7.60
3B 1b	Non-dairy cattle	5.17	4.72	4.25	4.54	2.85	3.41	3.53	3.53	3.63
3B 2	Sheep	0.05	0.12	0.11	0.12	0.14	0.11	0.09	0.09	0.09
3B 3	Swine	31.55	28.86	26.09	24.99	24.71	17.91	17.81	16.94	16.23
3B 4d	Goats	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01
3B 4e	Horses	0.76	0.72	0.68	0.73	0.85	0.72	0.67	0.67	0.65
3B 4gI	Laying hens	0.86	1.17	1.51	1.35	1.76	1.44	1.53	1.34	1.37
3B 4gII	Broilers	1.24	1.99	2.31	2.68	2.52	1.89	1.71	1.68	0.87
3B 4gIII	Turkeys	0.15	0.12	0.29	0.29	0.33	0.26	0.21	0.24	0.15
3B 4gIV	Other poultry	0.18	0.16	0.22	0.14	0.12	0.04	0.04	0.04	0.03
3B 4h	Other	4.72	5.18	4.01	4.70	6.25	6.92	6.84	7.07	5.69
3B	Total	53.33	50.95	46.77	46.85	47.16	40.02	39.97	39.29	36.31

Figure 5.5 shows the percentage distribution of the NH<sub>3</sub> emission from housing, storage and application of manure. The main part of the reduction in NH<sub>3</sub> emission has taken place in connection with the application of manure in fields, due to changes in manure application practice, see Chapter 5.4.2. There has been a reduction in emissions associated with storage of manure, which is a result of improvement in coverage of slurry tanks. As a consequence of this development, the percentage of emission from housing is increased from 38 % in 1985 to 54 % in 2013.

The possibilities for NH<sub>3</sub> reduction will likely be focused on measures in housings by various technological solutions. Some ammonia reducing technology is already implemented in housing e.g. air cleaning systems and slurry acidification, but the reduced effect of these technologies is not taken into account in the Danish inventory. Improvement in documentation is needed. Current, there is no survey which covers the number of farmers or number of housing using the ammonia reducing technology.




Figure 5.5 The percentage distribution of the NH<sub>3</sub> emission in manure management 1985-2013.

### 5.3.3 PM

#### Description

Investigations have shown that farmers, as well as livestock, are subject to an increased risk of developing lung and respiratory related diseases due to the particulate emissions (Hartung and Seedorf, 1999). This is because the particles are able to carry bacteria, viruses and other organic compounds.

In 2013 the PM emission from housings, given as TSP, is estimated to 10.23 Gg, which correspond to 14 % of the emission of TSP from the agricultural sector. Of the 10.23 Gg TSP, 69% relates to swine production. The emission from cattle and poultry contributes with 14 % and 16 %, respectively and the remainder animals contribute with 1 %.

#### Methodological issues

The estimation of PM emission is based on the EMEP/EEA guidebook (2013) where the scientific data mainly are based on an investigation of PM emission in North European housings (Takai et al., 1998).

The PM emission includes primary particles in the form of dust from housings. The inventory includes PM emission from cattle, swine, poultry, horses, sheep, goats and fur bearing animals (Table 5.10). The number of grazing days is taken into account. Some animal categories are divided into subcategories and for some categories (if applicable) distinction is made between solid and slurry based housing systems.

The PM emission is related to the annual average population (AAP) and to the time the animal is housed. The PM emission from grazing animals is considered as negligible.

Table 5.10 Livestock categories used in the PM emission inventory.

Livestock categories as given in NFR	Subcategories as given in Danish inventory the EMEP/EEA guidebook		Grazing days
Dairy Cattle	Dairy cattle	Dairy cattle	18
Non-Dairy Cattle	Calves	Calves < ½ yr	0
		Bulls	0
	Beef cattle	Heifers	132
		Suckling cattle	224
Swine	Sows	Sows (incl. weaners until 7 kg)	0
	Weaners	Weaners (7-32 kg)	0
	Fattening pigs	Fattening pigs (32-107 kg)	0
Poultry	Laying hens	Laying hens	0
	Broilers	Broilers	0
	Turkeys	Turkeys	0
	Other poultry	Ducks	0
		Geese	365
Horses	Horses	Horses	183
Sheep	Sheep	Sheep	265
Goats	Goats	Goats	265

#### Activity data

See Chapter 5.2.1

#### Emission factor

Emission factors for TSP, PM<sub>10</sub> and PM<sub>2.5</sub> are based on the EMEP/EEA guidebook (EMEP/EEA, 2013). The same emissions factors are used for all years. Estimation of TSP is based on the transformation factors between TSP and PM<sub>10</sub> as given in the EMEP/EEA emission inventory guidebook (2013).

Table 5.11 Emission factors for particle emission from animal housing system.

Livestock category	Housing system	Emission factor			Transformation factor
		PM <sub>10</sub>	PM <sub>2.5</sub>	TSP	PM <sub>10</sub> to TSP
kg per AAP per year					
<b>Cattle:</b>					
Dairy cattle	Slurry	0.83	0.54	1.81	0.46
	Solid	0.43	0.28	0.94	0.46
Calves < ½ yr	Slurry	0.15	0.10	0.34	0.46
	Solid	0.16	0.10	0.35	0.46
Beef cattle	Slurry	0.32	0.21	0.69	0.46
	Solid	0.24	0.16	0.52	0.46
Heifer <sup>1)</sup>	Slurry	0.49	0.32	1.07	0.46
	Solid	0.30	0.19	0.64	0.46
Suckling cattle <sup>2)</sup>	Slurry	0.32	0.21	0.69	0.46
	Solid	0.24	0.16	0.52	0.46
<b>Swine:</b>					
Sows	Slurry	0.61	0.11	1.36	0.45
	Solid	0.80	0.14	1.77	0.45
Weaners	Slurry	0.16	0.03	0.36	0.45
	Solid	0.16	0.03	0.36	0.45
Fattening pigs	Slurry	0.31	0.06	0.70	0.45
	Solid	0.37	0.07	0.83	0.45
<b>Poultry:</b>					
Laying hens, cages	Solid	0.025	0.003	0.02	1.00
Laying hens, perchery	Solid	0.119	0.023	0.12	1.00
Broilers	Solid	0.069	0.009	0.07	1.00
Ducks	Solid	0.139	0.018	0.14	1.00
Geese	Solid	0.243	0.032	0.24	1.00
Turkeys	Solid	0.521	0.068	0.52	1.00
Horses	Solid	0.22	0.14	0.48	0.46
Sheep	Solid	0.06	0.02	0.12	0.46
Goats	Solid	0.06	0.02	0.12	0.46
Fur bearing animals	Solid	0.008	0.004	0.01	1.00

<sup>1)</sup> Average of "calves" and "dairy cattle".

<sup>2)</sup> Assumed the same value as for "Beef cattle".

<sup>3)</sup> Same as slurry based systems.

### Emissions

Figure 5.6 shows the PM emission, given in TSP for each animal category in the period 2000 to 2013. It is seen that the main part of the emission originates from swine housings. See Annex 3D Table 3D-5 for the PM emission, given in TSP, PM<sub>10</sub> and PM<sub>2.5</sub>. In the period 2000 to 2013, the total agricultural emission of TSP from housings is decreased by 6 %. The decrease in the emission in 2003 is mainly due to change in housing type for non-dairy cattle. The decrease in 2008 is mainly due to change in housing types for swine. The increase in 2010 and following decrease in 2011 to 2013 is mainly due to change in number of swine.




Figure 5.6 PM emission from housings 2000 – 2013, Gg TSP.

### 5.3.4 NMVOC

#### Description

A first estimate of NMVOC from manure has been calculated and shows that 32 % of the total Danish NMVOC emission is related to animal husbandry and mainly to the cattle production.

#### Methodological issues

The estimation of NMVOC emission is based on the EMEP/EEA guidebook (2013). NMVOC emissions from animal husbandry comes from feed, degradation of feed in the rumen and from undigested fat, carbohydrate and protein decomposition in the rumen and in the manure. Silage is a major source of NMVOC emissions and therefore two sets of emission factor are introduced in the Guidebook; a high emission factor based on feeding with silage and a low emission factor based on feeding without silage.

The calculation of NMVOC emissions is based on Tier1 approach.

#### Activity data

The NMVOC emission is estimated on the number of animal multiplied with the NMVOC emission factor for each animal category. The number of animal is given as the average annual population (AAP) – see Table 5.5.

#### Emission factor

NMVOC emission factors recommended in EMEP/EEA Guidebook Table 3-3 is used (Table 5.12). For days on grass, the emission factor for feeding without silage is used for cattle, sheep, goats and horses (Table 5.10). However, all emissions are entered in NFR category 3B, while the notation key NE is used for NFR category 3Da3.

Same emissions factors are used during all years, which mean that changes of the emission over time depends on change in animal production or change in grazing days.

Table 5.12 NMVOC emission factors (EMEP/EEA Guidebook 2013, Tier1)

	EF NMVOC with silage	EF NMVOC without silage <sup>1</sup>
Dairy Cattle	17.937	8.047
Non-Dairy Cattle	8.902	3.602
Sheep	0.279	0.169
Swine – sows		1.704
Swine – other		0.551
Goats	0.624	0.542
Horses	7.781	4.275
Laying hens		0.165
Broilers		0.108
Turkeys		0.489
Other poultry		0.489
Fur bearing animals		1.941

<sup>1</sup>Emission factor is also used for time on grass.

### Emissions

The development of NMVOC emission from 1990 to 2013 shows a decrease from 38 Gg to 36 Gg with the highest fall in the beginning of the period (Figure 5.7). Back in 1990 two third of the emission originates from the cattle production, which is fallen to half the emission in 2013. A decrease of emission from cattle is a consequence of less number of animals due to higher milk yield and production ceiling due to the EU milk quota. An increase of the production of swine and fur bearing animals has resulted in an increase of the emission from 1990 to 2013




Figure 5.7 Emission of NMVOC from manure management, 1990-2013.

### 5.3.5 NO<sub>x</sub>

#### Description

A first estimate of NO<sub>x</sub> from manure management has been calculated and shows that less than 1 % of the total Danish NO<sub>x</sub> emission in 2013 is related to animal husbandry.

### Methodological issues

The estimation of NO<sub>x</sub> emission is based on the EMEP/EEA guidebook (2013) Tier1 approach. The Guidebook mentioned that “Emissions of NO are estimated to quantify the N mass balance for the Tier 2 methodology for calculating NH<sub>3</sub> emissions”. However, only two EF Tier2 is shown in Guidebook Chapter 3.3. Manure Management Appendix B – for dairy cattle and fattening pigs. Using of Tier 2 approach requires probably that the estimation of the NO<sub>x</sub> EF is based on a national N-balance.

### Activity data

The Tier 1 approach is based on number of animal given as the average annual population (AAP). The Number is showed in Table 5.5.

### Emission factor

Emission factor for estimation of NO emission from manure management is listed in Table 5.13. Some of the manure from the mink production is handled as slurry, but no EF for slurry is mentioned in the Guidebook. Therefore, the same emissions factor is used for both slurry and solid systems.

The emission factors in the Guidebook are given as NO. All emissions are converted from NO emission to NO<sub>2</sub> emission by multiply with a conversion factor of 46/30 due to the differences on molecular weight.

Table 5.13 NO emission factors (EMEP/EEA Guidebook 2013), kg NO per AAP.

NFR code	Livestock	slurry	solid
3B 1a	Dairy cows	0.007	0.154
3B 1b	Other cattle	0.002	0.094
3B 2	Sheep		0.005
3B 3	Sows	0.004	0.132
3B 3	Fattening pigs	0.001	0.045
3B 4d	Goats		0.005
3B 4e	Horses		0.131
3B 4gi	Laying hens	0.003	0.0001
3B 4gii	Broilers		0.001
3B 4giii	Turkeys		0.005
3B 4giv	Ducks		0.004
3B 4giv	Geese		0.001
3B 4h	Fur bearing animals	0.0002 <sup>1</sup>	0.0002

<sup>1</sup> Used the same EF as given for solid manure.

### Emissions

The NO<sub>x</sub> emission from 1990 to 2013 has decreased significantly from 0.48 Gg NO<sub>x</sub> to 0.22 Gg NO<sub>x</sub> corresponding to a 55 % reduction. The emission depends on number of animal and manure type and the decrease is mainly related to changes from solid based system to slurry based system for both the dairy cattle and the swine production. Thus, the allocation of solid manure was 23 % in 1990 and dropped to the half 11 % in 2013.






Figure 5.8 NO<sub>x</sub> emission from manure management 1990–2013.

## 5.4 Soils

At present, farmed area covers about 60 % of the total land area in Denmark. In recent decades, farmed area has decreased, being replaced by built-up areas, roads, forest and nature habitats.

### 5.4.1 Inorganic N-fertilizers

#### Description

For the sector inorganic N-fertilizer the emission of NH<sub>3</sub> and NO<sub>x</sub> are estimated.

The emission of NH<sub>3</sub> from inorganic fertilizer contributes in 2013 with 10 % of the emission from the agricultural sector. The emission of NO<sub>x</sub> contributes in 2013 with 54 % of the emission from the agricultural sector.

#### Methodological issues

Emission of NH<sub>3</sub> from inorganic fertilizer is based on the consumption of fertilizer of different types and emission factors. In Table 5.14 are shown emission factors and consumption for 2013. See Annex 3D Table 3D-6 for assumptions for fertilizer type.

Emission of NO<sub>x</sub> is based on the total consumption of N in inorganic N-fertilizer and emission factor.

Table 5.14 Inorganic N-fertilizer consumption 2013 and emission factors.

Fertilizer type	NH <sub>3</sub> Emission factor <sup>1</sup> , Consumption <sup>2</sup> ,	
	Kg NH <sub>3</sub> -N pr kg N	t N
Calcium and boron calcium nitrate	0.11	0.1
Ammonium sulphate	0.01	5.5
Calcium ammonium nitrate and other nitrate types	0.02	97.6
Ammonium nitrate	0.04	5.7
Liquid ammonia	0.01	6.7
Urea	0.24	0.4
Other nitrogen fertilizer	0.04	19.3
Magnesium fertilizer	0.11	0.0
NPK-fertilizer	0.04	49.5
Diammonphosphate	0.11	1.2
Other NP fertilizer types	0.11	5.6
NK fertilizer	0.04	1.8
Total consumption of N in inorganic N-fertilizer		193.6

<sup>1</sup> EMEP/EEA (2013), see Annex 3D Table 3D-6 for assumptions for fertilizer type.

<sup>2</sup> The Danish AgriFish Agency.

### Activity data

Data on the use of inorganic fertilizer is based on the annual sale estimations collected by the Danish AgriFish Agency (2013). The use of inorganic fertilizer includes fertilizer used in parks, golf courses and private gardens. Approximately 1 % of the inorganic fertilizer can be related to use outside the agricultural area.

### Emission factor

Emission factors for both NH<sub>3</sub> and NO<sub>x</sub> are based on the values given in EMEP/EEA guidebook (EMEP/EEA, 2013) and the same emission factors are used for all years 1985-2013. The implied emission factor for NH<sub>3</sub> is shown in Table 5.15 and it depends on consumption and type of fertilizer.

Table 5.15 Implied emission factor NH<sub>3</sub> for inorganic N-fertilizer, 1985-2013.

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Implied emission factor NH <sub>3</sub> , % of total N	3.24	3.47	3.98	3.24	3.12	3.10	3.08	3.11	3.10

### Emissions

Since 1985 there has been a significant decrease in use of inorganic N-fertilizer. This is due to requirements to utilising of nitrogen in manure as outlined in various environmental action plans. Another explanation for a reduction of emission is a decrease in use of urea as currently accounting for less than 1 % of the total nitrogen (Table 5.14). In Figure 5.9 are shown emission of NH<sub>3</sub> and NO<sub>x</sub>.




Figure 5.9 Emission of NH<sub>3</sub> and NO<sub>x</sub> for 1985-2013, Gg.

## 5.4.2 Animal manure applied to soils

### Description

For the sector animal manure applied to soils the emission of NH<sub>3</sub> and NO<sub>x</sub> are estimated.

Emission of NH<sub>3</sub> from animal manure applied to soils contributes in 2013 with 28 % of the NH<sub>3</sub> emission from the agricultural sector. Emission of NO<sub>x</sub> from animal manure applied to soils contributes in 2013 with 42 % of the NO<sub>x</sub> emission from the agricultural sector.

### Methodological issues

To calculate both emissions of NH<sub>3</sub> and NO<sub>x</sub> from animal manure applied to soils an emission factor are estimated and multiplied with TAN ex storage for liquid manure and N ex storage for solid manure for each animal type.

### Activity data

Based on the normative figures (Poulsen, 2014) the amount of TAN ex storage for liquid manure and the amount of N ex storage for solid manure are estimated.

### Emission factor NH<sub>3</sub>

The emission factor are based on background estimates of time of application, application methods, application in growing crops or on bare soil and the time from application to ploughing in soil. The amount of manure a acidified is also taken into account. The emission factor differs between solid manure and liquid manure and also between manure from cattle and swine. For all other animals same emission factor as for cattle is used.

The emission factors will vary from year to year depending on changes in the practice of application. In Table 5.16 background information for 2013 are given. This estimate is based on information from the Danish Agricultural Advisory Service.

Table 5.16 Estimate for application method, time of application and time before the manure is incorporated in the soil for 2013.

Liquid manure		Percentage distribution of manure		Length of time before incorporation into soil, hours							
Application methods	Application time			0		4, and then harrowed		4, and then Ploughed		Not incorporated	
		Cattle	Pigs	Cattle	Pigs	Cattle	Pigs	Cattle	Pigs	Cattle	Pigs
Incorporated	winter-spring	61	24	61	24	-	-	-	-	-	-
Incorporated	summer-autumn	18	4	18	4	-	-	-	-	-	-
Trailing hoses	winter-spring	17	64	-	-	-	3	-	2	17	59
Trailing hoses	spring-summer	2	2	-	-	-	-	-	-	2	2
Trailing hoses	late summer-autumn	2	6	-	-	-	2	-	1	2	3
<b>Total</b>		<b>100</b>	<b>100</b>	<b>79</b>	<b>28</b>	<b>-</b>	<b>5</b>	<b>-</b>	<b>3</b>	<b>21</b>	<b>64</b>

Solid manure		Percentage distribution of manure		Length of time before incorporation into soil, hours							
Application methods	Application time			0		4		6		Not incorporated	
		Cattle	Pigs	Cattle	Pigs	Cattle	Pigs	Cattle	Pigs	Cattle	Pigs
Broad spreading	winter-spring	90	76	-	-	70	60	20	16	-	-
Broad spreading	spring-summer	5	5	5	5	-	-	-	-	-	-
Broad spreading	late summer-autumn	5	19	-	-	5	19	-	-	-	-
<b>Total</b>		<b>100</b>	<b>100</b>	<b>5</b>	<b>5</b>	<b>75</b>	<b>79</b>	<b>20</b>	<b>16</b>	<b>-</b>	<b>-</b>

Acidification of manure in the housings, storage or just before application is an increasing used technique in Denmark. The acidification of the manure lowers the emission of NH<sub>3</sub> from the treated manure by 49 % (VERA, 2010). The amount of manure acidified is estimated by Knowledge Centre for Agriculture for the years 2011, 2012 and 2013 (Vestergaard, 2014). It is mainly cattle manure which is acidified.

Table 5.17 Share of liquid manure acidified, 2011-2013.

	2011	2012	2013
Share of cattle manure, %	3	6	10
Share of swine manure, %	1	1	1

In 2013 the emission factor for cattle is for solid manure estimated to 7 % of N ex storage and for liquid manure estimated to 13 % TAN ex storage, for swine the emission factors are 6 % and 11 %, respectively.

#### Emissions factor NO<sub>x</sub>

The emission factor for NO<sub>x</sub> is based on EMEP/EEA guidebook (2013). Only one emission factor regarding the NO<sub>x</sub> emission for 3D is mentioned in the EMEP/EEA Guidebook (refer to Table 3-1). The background reference for the Tier 1 emission factor is based on a literature study, which do not distinguish between different kinds of fertilizer types. This indicate that the same emission factor can be used independent of the crops are fertilized with mineral fertilizer or manure. The NO<sub>x</sub> emission is estimated based on the Tier 1 emission factor at 0.026 kg NO per kg N fertilized. Because the emission factor in the Guidebook is given as NO, the emission is converted from NO emission to NO<sub>2</sub> emission by multiply with a conversion factor of 46/30 due to the differences on molecular weight.

#### Emissions

The emission of NH<sub>3</sub> from manure applied to soils has decreased by 51 % from 1985 to 2013, this is due to decrease of N excreted by animals and by changes in the way manure is handled during application. Based on the ac-

tion plans various initiatives has been implemented and include for example requirement for a minimum 9-month manure storage capacity, requirement that manure applied to soil be ploughed down within six hours, a ban on the application of manure in winter and broad spreading is no longer allowed.

Emission of NO<sub>x</sub> from manure applied to soils has decreased by 7 % from 1985 to 2013 this is mainly due to decrease of N excreted.




Figure 5.10 Emission of NH<sub>3</sub> and NO<sub>x</sub> from manure applied to soils, 1985-2013, Gg NH<sub>3</sub> and NO<sub>x</sub>.

### 5.4.3 Sewage sludge applied to soils

#### Description

For the sector sewage sludge applied to soils the emission of NH<sub>3</sub> and NO<sub>x</sub> are estimated.

Emission of NH<sub>3</sub> and NO<sub>x</sub> from sewage sludge applied to soils contributes in 2013 with less than 1 % and 2 % from the agricultural sector, respectively.

#### Methodological issues

Amount of N applied are multiplied with the emission factor.

#### Activity data

Information on amount of sewage sludge, N-content and NH<sub>3</sub> emission factor is obtained from reports prepared by the Danish Environmental Protection Agency and based on data from the fertilizer accounts controlled by The Danish AgriFish Agency. Farmers with more than 10 animal units have to be registered and keep accounts of the use of N content in manure, received manure or other organic fertilizer.

Table 5.18 Activity data used to estimate NH<sub>3</sub> and NO<sub>x</sub> from sewage sludge, 1985-2013.

		1985	1990	1995	2000	2005	2010	2011	2012	2013
Amount of sludge applied on soil	Tonnes of dry matter	50 000	77 883	112 235	83 727	45 739	56 665	54 564	52 002	51 731
N-content	%	4.00	4.00	4.13	4.33	4.75	4.75	4.75	4.75	4.75
N applied on soil	Tonnes N	2 000	3 115	4 635	3 625	2 173	2 692	2 592	2 470	2 457

### Emission factor NH<sub>3</sub>

The emission factor for NH<sub>3</sub> emission from sewage sludge applied to soil is based on information from the Danish Environmental Protection Agency. It is estimated to 0.019 kg NH<sub>3</sub>-N per kg N and the same for all years 1985-2013.

### Emission factor NO<sub>x</sub>

The emission factor for NO<sub>x</sub> is based on EMEP/EEA guidebook (2013). Only one emission factor regarding the NMVOC emission from 3 D is mentioned in the EEA/EMEP Guidebook (refer to Table 3-1). The background reference for the Tier 1 emission factor is based on a literature study, which do not distinguish between different kinds of fertilizer types. Emission from N in sewage sludge could be lower or higher, but no information on this is available and therefore the emission factor from the Guidebook at 0.026 kg NO per kg N sewage sludge applied to soils is used.

Because the emission factor in the Guidebook is given as NO, the emission is converted from NO emission to NO<sub>2</sub> emission by multiply with a conversion factor of 46/30 due to the differences on molecular weight.

### Emissions

Emission of NH<sub>3</sub> and NO<sub>x</sub> from sewage sludge is shown in Figure 5.11. The emission follow the amount of N applied.




Figure 5.11 Emission of NH<sub>3</sub> and NO<sub>x</sub> from sewage sludge, 1985-2013, Gg.

### 5.4.4 Urine and dung deposited by grazing animals

#### Description

It is assumed that 5 % of the manure from dairy cattle is deposited in the field, which corresponding to 18 days per year (Aaes, 2008). For heifers 36 % of the nitrogen in the manure is estimated deposited during grazing (Aaes, 2008), 61 % for suckling cows (Poulsen et al, 2001), 50 % for horses (Clausen, 2008) and 73 % for sheep and goats (Poulsen et al, 2001).

Emission of NH<sub>3</sub> from urine and dung deposit by grazing animals contributes in 2013 with 3 % of the emission from the agricultural sector.

### Methodological issues

Emission of urine and dung deposited by grazing animals is based on N excreted ab animal, number of days the animals are on grass and the emission factor.

### Activity data

The activity data are number of animals (see Chapter 5.2.1), N excreted ab animal and number of days on grass (see Table 5.10) which combined gives the N deposit on grass, see Table 5.19.

Table 5.19 N deposit on grass, 1985-2013, M kg N.

	1985	1990	1995	2000	2005	2010	2011	2012	2013
N deposited on grass	37	34	36	34	26	22	21	22	22

### Emission factor

Study of grazing cattle indicates that 7 % of the total nitrogen content is assumed to evaporate as NH<sub>3</sub> (Jarvis *et al.* 1989a, Jarvis *et al.* 1989b and Bus-sink 1994). This emission factor is used for all animal categories.

### Emissions

The emission of NH<sub>3</sub> from urine and dung deposit by grazing animals has decreased by 41 % from 1985 to 2013 and this is mainly due to decrease in number of dairy cattle and decrease in number days on grass for dairy cattle.

Table 5.20 Emission of NH<sub>3</sub> from urine and dung deposit by grazing animals, 1985-2013, Gg NH<sub>3</sub>.

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Grazing animals	3.12	2.91	3.02	2.92	2.21	1.87	1.81	1.84	1.86

## 5.4.5 Farm-level agricultural operations including storage, handling and transport of agricultural products

### Description

During agricultural operations such as soil cultivation, harvesting, cleaning, drying and transport an emission PM occur. In the EMEP/EEA guide-book are only method and emission factors for the operations done in the field, that is soil cultivation, harvesting, cleaning and drying.

The emission of PM TSP from field operations contributes with 62 % of the total emission of TSP in 2013.

### Methodological issues

The emission of PM from field operations is calculated by area of cultivated crops multiplied with number of operations and emission factor, for each crop type and type of operation.

### Activity data

For activity data are used area of cultivated crops and number of operations for each crop. The area of crops is estimated by Statistic Denmark (DSt, 2014) and number of operations are based on budget estimates made by Knowledge Centre for Agriculture. See Annex 3D Table 3D-7 for area of cultivated crops and Annex 3D Table 3D-8a-8d for number of operations divided in soil cultivation, harvesting, cleaning and drying.

### Emission factor

The emission factors used are given in Table 5.21 and they are based on EMEP/EEA guidebook (EMEP/EEA, 2013) and van der Hoek (2007).

Table 5.21 Emission factors for field operations, kg per ha.

PM <sub>10</sub>	Soil cultivation	Harvesting	Cleaning	Drying
Wheat	0.25 <sup>a</sup>	0.27 <sup>b</sup>	0.19 <sup>a</sup>	0.56 <sup>a</sup>
Rye	0.25 <sup>a</sup>	0.2 <sup>b</sup>	0.16 <sup>a</sup>	0.37 <sup>a</sup>
Barley	0.25 <sup>a</sup>	0.23 <sup>b</sup>	0.16 <sup>a</sup>	0.43 <sup>a</sup>
Oat	0.25 <sup>a</sup>	0.34 <sup>b</sup>	0.25 <sup>a</sup>	0.66 <sup>a</sup>
Other arable	0.25 <sup>a</sup>	0.26 <sup>c</sup>	0.19 <sup>c</sup>	0.51 <sup>c</sup>
Grass	0.25 <sup>a</sup>	0.25 <sup>a</sup>	0 <sup>a</sup>	0 <sup>a</sup>
<b>PM<sub>2.5</sub></b>				
Wheat	0.015 <sup>a</sup>	0.011 <sup>b</sup>	0.009 <sup>a</sup>	0.168 <sup>a</sup>
Rye	0.015 <sup>a</sup>	0.008 <sup>b</sup>	0.008 <sup>a</sup>	0.111 <sup>a</sup>
Barley	0.015 <sup>a</sup>	0.009 <sup>b</sup>	0.008 <sup>a</sup>	0.129 <sup>a</sup>
Oat	0.015 <sup>a</sup>	0.014 <sup>b</sup>	0.0125 <sup>a</sup>	0.198 <sup>a</sup>
Other arable	0.015 <sup>a</sup>	0.010 <sup>c</sup>	0.009 <sup>c</sup>	0.152 <sup>c</sup>
Grass	0.015 <sup>a</sup>	0.01 <sup>a</sup>	0 <sup>a</sup>	0 <sup>a</sup>
<b>TSP<sup>d</sup></b>				
Wheat	2.5	2.7	1.9	5.6
Rye	2.5	2	1.6	3.7
Barley	2.5	2.3	1.6	4.3
Oat	2.5	3.4	2.5	6.6
Other arable	2.5	2.6	1.9	5.1
Grass	2.5	2.5	0	0

<sup>a</sup> EMEP/EEA (2013).

<sup>b</sup> van der Hoek (2007).

<sup>c</sup> average of wheat, rye, barley and oat.

<sup>d</sup> PM<sub>10</sub> multiplied by 10 (van der Hoek, 2007).

### Emissions

The emission of PM<sub>10</sub>, PM<sub>2.5</sub> and TSP are shown in Table 5.22. The emission of TSP has decreased 11 % from 2000 to 2013 due to decrease in the area of cultivated crops and number of treatments of the fields.

Table 5.22 Emissions of PM<sub>10</sub>, PM<sub>2.5</sub> and TSP from field operations, 1985-2013, tonnes.

	2000	2005	2010	2011	2012	2013
PM <sub>10</sub>	6 238	5 415	5 665	5 654	5 559	5 522
PM <sub>2.5</sub>	479	436	468	457	445	448
TSP	62 382	54 146	56 655	56 541	55 587	55 218

### 5.4.6 Cultivated crops

#### Description

For the sector cultivated crops the emission of NH<sub>3</sub> and NMVOC are estimated.

The Danish emission inventory includes NH<sub>3</sub> emission from crops, despite the uncertainties related to this emission source. Literature research shows



that the volatilisation from crop types differs considerably. However, as for the emission ceiling given in the Gothenburg-Protocol and the EU NEC Directive the emission from crops is not taken into account.

#### Methodological issues

The emission is calculated based on area of agricultural land and emission factors.

#### Activity data

Activity data are obtained from Statistics Denmark, see Annex 3.D Table 3D-7.

#### Emission factor NH<sub>3</sub>

EF's for crops are estimated to 2 % for crops and 0.5 % for grass based on a literary survey (Gyldenkærne and Albrektsen, 2009).

Table 5.23 EF used to estimate the emission of NH<sub>3</sub> from crops.

Crops	kg NH <sub>3</sub> -N per ha
Cash crops, beets and silage maize	2
Grass/clover in rotation	0.5
Permanent grass	0.5
Set-a side	0

#### Emission factor NMVOC

The calculation of the NMVOC emission is based on emission factors recommended in EMEP/EEA Guidebook 2013 Table A3-2 for cultivation of wheat, rye, rape and grass land (15 C). A Tier 2 IEF is estimated corresponding to Danish yield level dry matter content (DM) for these crop types. The emission from other crop types is not available in the Guidebook. However, the total NMVOC emission is estimated as the Tier 2 IEF multiplied with the total cultivated area.

The NMVOC emission from cultivated crops is estimated to 1.83 Gg in 2013 based on an IEF at 0.70 and a cultivated area of 2 628 thousand hectare. The IEF varies annually from 0.51 -0.70 kg NMVOC per hectare depending on the allocation of the four mentioned crop types. Higher allocation of rape and rye result in higher IEF due to a higher emission factor for these two crop types.

Table 5.24 Estimation of a Tier 2 NMVOC emission factor.

2013	EEA/EMEP, Emission factor	Fraction of year emitting	Total	Mean dry matter of crop	NMVOC EF	Cultivated area	NMVOC emission	Tier 2 DK
Crop	Kg NMVOC /kg DM/yr		Kg/kg DM//yr	kg DM/ha	Kg/ha/yr	ha	Kg/ha/yr	IEF, kg NMVOC/ha
Wheat	2.60E-08	0.3	6.82E-05	6 299	0.43	542 051	232 831	
Rye	1.41E-07	0.3	3.70E-04	5 134	1.87	88 181	167 576	
Rape	2.02E-07	0.3	5.30E-04	3 587	1.84	175 117	332 837	
Grass land*	1.03E-08	0.5	4.51E-05	7 974	0.42	515 615	185 497	
Total						1 320 964	918 741	0.70

\*Grass land 15 C

## Emissions

Emission of NH<sub>3</sub> and NMVOC are shown in Figure 5.12. The emission of NH<sub>3</sub> has decreased by 10 % from 1985 to 2013 and the emission of NMVOC has decreased by 2 %. This is mainly due to decrease in the agricultural area.




Figure 5.12 Emission of NH<sub>3</sub> and NMVOC from cultivated crops, 1985-2013, Gg.

## 5.4.7 Use of pesticides

### Description

A range of pesticides are used in the Danish agricultural sector and some of them contain Hexachlorobenzene (HCB), but pure HCB used as pesticide is banned. HCB is a poisonous substance, which is dangerous to human and animal health but is used as agent in pesticides.

The emission of HCB from use of pesticides contributes with less than 1 % of the Danish total HCB emission.

### Methodological issues

Emission of HCB from use of pesticides is based on amount of effectual substance used and emission factors for each type of pesticides.

### Activity data

A range of pesticides are used in Denmark. In the period from 1990 to 2011 six types of pesticides containing HCB have been identified as used in Denmark. These are atrazine, chlorothalonil, clopyralid, lindane, pichloram and simazine. Data of amounts of effectual substance used in Denmark are collected from Environmental Protection Agency (EPA), see Table 5.25. The use of atrazine and lindane stopped in 1994 and the use of chlorothalonil and simazine ceased in 2000 and 2004, respectively.

Table 5.25 Amounts of effectual substance used in Denmark, 1990-2013, kg.

	1990	1995	2000	2005	2010	2011	2012	2013*
Atrazine	91 294	-	-	-	-	-	-	-
Chlorothalonil	10 512	10 980	7 340	-	-	-	-	-
Clopyralid	16 461	22 587	7 446	5 874	9 122	11 840	8 170	8 170
Lindane	8 356	-	-	-	-	-	-	-
Pichloram	-	-	-	-	723	1 349	206	206
Simazine	30 234	19 865	23 620	-	-	-	-	-

\*Same as 2012 due to lack of data.

### Emission factor

No default emission factors are given in EMEP/EEA Guidebook. Emission factors given in Yang (2006) are used in the calculation of the emissions, see Table 5.26.

Table 5.26 Emission factors for HCB from pesticides, 1990-2013, g per tonnes.

	1990	1995	2000	2005	2010	2011	2012	2013
Atrazine	100	1	1	1	1	1	1	1
Chlorothalonil	500	40	40	10	10	10	10	10
Clopyralid	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
Lindane	100	50	50	1	1	1	1	1
Pichloram	100	50	50	8	8	8	8	8
Simazine	100	1	1	1	1	1	1	1

### Emissions

Table 5.27 shows the emission of HCB from the agricultural sector for the years 1990-2013. The emission has decreased significantly from 1990 to 2013 due to decrease in use of pesticides containing HCB.

Table 5.27 Emission of HCB, 1990-2013, kg.

	1990	1995	2000	2005	2010	2011	2012	2013
Pesticides	18.28	0.50	0.33	0.01	0.02	0.03	0.02	0.02

## 5.5 Field burning of agricultural residues

### Description

Field burning of agricultural residues has been prohibited in Denmark since 1990 and may only take place in connection with production of grass seeds on fields with repeated production and in cases of wet or broken bales of straw.

Emissions of NH<sub>3</sub>, NO<sub>x</sub>, CO, NMVOC, SO<sub>2</sub>, PM, BC, heavy metals, dioxin, PAHs, HCB and PCB are included under the NFR category 3F. The emission of NH<sub>3</sub> from field burning contributes in 2013 with less than 1 % of the agricultural emission. Emissions of PM and NMVOC from field burning contributes with less than 1 % TSP, 2 % PM<sub>10</sub>, 13 % PM<sub>2.5</sub> and less than 1 % NMVOC of the agricultural emission. The emission of NO<sub>x</sub>, BC, CO, SO<sub>2</sub>, heavy metals, dioxin and PCB from field burning contribute with less than or around 1 % of the total national emission, while the emission of PAHs and HCB contribute with around 4-6 % of the national emission. From 1989 to 1990 all emissions decrease significantly due to the ban on field burning.

### Methodological issues

Emissions from field burning of agricultural residues are calculated based on the amount of burnt straw given in tons dry matter and emission factors given in the EMEP/EEA guidebook (EMEP/EEA, 2013).

### Activity data

The amount of burnt straw from the grass seed production is estimated as 15-20 % of the total amount produced. The amount of burnt bales of wet straw is estimated as 0.1 % of total amount of straw. Both estimates are based on expert judgement by the Danish Agricultural Advisory Service.

The total amounts are based on data from Statistics Denmark. See Annex 3D Table 3D-9 for activity data.

### Emission factor

The EMEP/EEA guidebook (EMEP/EEA, 2013) default values for the emission factors for field burning of agricultural residues are used (Table 5.28).

Table 5.28 EF for field burning of agricultural residues.

Pollutant	EF	Unit
NO <sub>x</sub> <sup>1</sup>	2.4	g/kg DM
CO <sup>1</sup>	58.9	g/kg DM
NMVOC <sup>1</sup>	6.3	g/kg DM
SO <sub>x</sub> <sup>1</sup>	0.3	g/kg DM
NH <sub>3</sub> <sup>1</sup>	2.4	g/kg DM
TSP <sup>1</sup>	5.8	g/kg DM
PM <sub>10</sub> <sup>1</sup>	5.8	g/kg DM
PM <sub>2.5</sub> <sup>1</sup>	5.5	g/kg DM
BC <sup>1</sup>	0.5	g/kg DM
PCDD/F <sup>1</sup>	500	ng TEQ/t
Pb <sup>1</sup>	0.865	mg/kg DM
Cd <sup>1</sup>	0.049	mg/kg DM
Hg <sup>1</sup>	0.008	mg/kg DM
As <sup>1</sup>	0.058	mg/kg DM
Cr <sup>1</sup>	0.22	mg/kg DM
Ni <sup>1</sup>	0.177	mg/kg DM
Se <sup>1</sup>	0.036	mg/kg DM
Zn <sup>1</sup>	0.028	mg/kg DM
Cu <sup>2</sup>	0.0003	mg/kg DM
Benzo(a)pyrene <sup>2</sup>	2 787	mg/kg DM
benzo(b)fluoranthene <sup>2</sup>	2 735	mg/kg DM
benzo(k)fluoranthene <sup>2</sup>	1 073	mg/kg DM
Indeno(1,2,3-cd)pyrene <sup>2</sup>	1 017	mg/kg DM
HCB (broken bales) <sup>3</sup>	0.003	g/tonnes
HCB (seed production) <sup>3</sup>	0.002	g/tonnes
PCB (broken bales) <sup>4</sup>	3	ng TEQ/t
PCB (seed production) <sup>4</sup>	0.05	ng TEQ/t

<sup>1</sup> EMEP/EEA, 2013.

<sup>2</sup> Jenkins, 1996.

<sup>3</sup> Yang (2006).

<sup>4</sup> Black et al. (2012).

### Emissions

See Annex 3D Table 3D-10 for emissions of all pollutants 1985 to 2013.

## 5.6 Agriculture other

### 5.6.1 NH<sub>3</sub> treated straw

#### Description

NH<sub>3</sub> is used for conservation of straw for feeding. As for the emission ceiling given in the Gothenburg-Protocol and the EU NEC Directive the emission from NH<sub>3</sub> treated straw is not taken into account.

### Methodological issues

Emissions are calculated as  $\text{NH}_3$  used for treatment of straw multiplied the emission factor.

### Activity data

Information on  $\text{NH}_3$  used for treatment of straw is collected from the suppliers.  $\text{NH}_3$  treated straw has been prohibited from 2006, but in 2010 and 2011 an exemption were given due to wet weather.

Table 5.29 Activity data for  $\text{NH}_3$  treated straw 1985 to 2013.

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Tonnes $\text{NH}_3\text{-N}$	8 285	12 912	8 406	3 125	329	300	300	0	0

### Emission factor

Investigations show that up to 80-90% of the supplied  $\text{NH}_3$  (given in  $\text{NH}_3\text{-N}$ ) can emit (Andersen et al., 1999). However, the emissions can be reduced particularly if the right dose is used. It is assumed that the emission factor is 65 % of the applied  $\text{NH}_3\text{-N}$ .

### Emissions

Emission of  $\text{NH}_3$  from  $\text{NH}_3$ -treated straw is shown in Figure 5.13.




Figure 5.13 Emission of  $\text{NH}_3$  from  $\text{NH}_3$ -treated straw, 1985-2013.

## 5.7 Uncertainties

Table 5.30 shows the estimated uncertainties for activity data and emissions factor for each pollutant.

### $\text{NH}_3$

#### 3B Manure management

It is defined that activity for manure management covers both the number of animals and housing type. The allocation of animal on different the housing types determines if the manure handled as slurry or solid manure.

The number of animals for the most important animal categories is estimated by Statistic Denmark and in 2013 the uncertainties for swine is estimated to 1.3 % and 0.9 % for cattle. The uncertainty is higher for less important animal groups, e.g. fur bearing animals (3.4 %), poultry, horses, sheep (10.4 %) and goats is. The uncertainty for number of animals overall is estimated to 2 %. The allocation of housing system is based on infor-

mation from the farm nitrogen budgets handled and controlled by the Danish AgriFish Agency. All farmers have to submit the information regarding the housing type annually and the uncertainty is assumed as relatively low.

The uncertainties for the activity data is thus a combination of low uncertainty in animal numbers, a relatively low uncertainty for housing type, which assumed to result in an overall uncertainty by 5 %.

The uncertainty for the emission factor covers nitrogen excretion, grazing days and NH<sub>3</sub> emission factors from housing and during storage of the manure. The Danish Normative System for animal excretions is based on data from the Danish Agricultural Advisory Services (DAAS), which is the central office for all Danish agricultural advisory services. DAAS engages in a great deal of research as well as the collection of efficacy reports from Danish farmers for dairy production, meat production, pig production, etc., to optimise productivity in Danish agriculture. Feeding plans from 15-18 % of the Danish dairy productions, 25-30 % of swine productions, 80-90 % of poultry productions and approximately 100 % of fur productions are collected annually. These basic feeding plans are used to develop the standard values of the "Danish Normative System". However, due to the large number of farms included in the norm figures, the arithmetic mean can be assumed as a very good estimate with a low uncertainty.

Regarding the uncertainties for the emission factor, it has to be included that the emission comes from three different places in the livestock production; from manure in housing, from stored manure and from application of manure. The uncertainties for emission measurements in housing, which are the basement for the normative standards varies from 15 -25 % (Poulsen et al., 2001). But there is no specified uncertainty estimates for emission factors for storage and application of manure. The overall uncertainty value for NH<sub>3</sub> emission factor for manure management is assumed to be around 25%.

### **3Da1 Inorganic fertilizers**

The activity data for the emission from inorganic N-fertilizer depends on the amount of sold fertilizer and the N-content for each fertilizer type, which is based on annually information given by the Danish AgriFish Agency. Uncertainty is considered to be low; 3 % based on expert judgement.

No uncertainty values for the emission factor are given in the EMEP/EEA guidebook. The Danish inventory assume an uncertainty value at 25 %, which indicated a uncertainty in the translation of the Danish fertilizer types to types specified in the guidebook but also indicate an uncertainty of the emission factors specified in the guidebook.

### **3Da2a Animal manure applied to soils**

Besides the number of animal, the uncertainty for activity data covers N-excretion, grazing days and the NH<sub>3</sub> emission from housing and storage. It is assumed that the most important variables are the number of animal which has a low uncertainty 2 %. However, the uncertainty is also affected by the other variable which has a higher uncertainty estimate. Thus, the uncertainty for the activity data is assumed to be around 15 %.

The emission factor depends on the uncertainty regarding the information on application time, application technics and plant cover. The uncertainty is estimated to 25 %.

### **3Da2b Sewage sludge applied to soils**

From 2005 and onwards the amount of N applied from wastewater treatment is based on the fertilizer accounts controlled by the Danish AgriFish Agency. Farmers with more than 10 animal units<sup>1</sup> have to be registered and have to keep accounts of the N content in manure, received manure or other organic fertilizer. The uncertainty for the activity data is assumed to be 15 %.

The emission factor depends on the application of time, application technic and the climate conditions and the uncertainty is assumed to be relatively high – around 50 %.

### **3Da3 Urine and dung deposited by grazing animals**

The overall uncertainty for the activity is estimated to 5 %. Besides the number of animals, the uncertainty depends on number of grassing days.

Regarding the uncertainty for the emissions factor, this depends on the N excretion and the climate conditions as temperature, wind and precipitation. The uncertainty value is estimated to 25 %.

### **3De Cultivated crops**

The activity data covers the cultivated area which is based on Statistics Denmark. For the major crops, the uncertainty is relatively low – e.g. winter wheat it is 1.1 % in 2013. The overall uncertainty for the activity is estimated to 2 %. Knowledge concerning the emission is relatively limited and therefore the uncertainty is assumed to be 50 %.

### **3F Field burning of agricultural residues**

An uncertainty of 25 % for the activity for field burning of agricultural residues is used. The uncertainty is a combination of the uncertainty for area of grass for seed production, which has a low uncertainty, amount of burned straw and yield, which have a high uncertainty. The uncertainties for the emission factor are based on the EMEP/EEA Guidebook (EMEP/EEA, 2013) and Jenkins et al. (1996).

### **3I Agriculture other**

Under NFR category 3I emissions from NH<sub>3</sub> treated straw is entered. NH<sub>3</sub> treated straw was until 2006 used as cattle feed. By law in 2006 the NH<sub>3</sub> treatment of straw was banned. However, due to wet weather conditions a dispensation to the law was given in 2010 and 2011. The activity depends on the amount of ammonia used in the second half of the year and is based on information from the AgriFish Agency. The uncertainty value is assumed to be 20 %. The uncertainty level for the emission factor is assumed to be 50 %.

### **PM**

Uncertainty estimates due to the activity data is estimated to 7 %. Besides number of animal and housing type, also uncertainty related to the production cycles plays a role.

The activity data covers the cultivated crops and number of operations for each crop type. The area of crops is estimated by Statistic Denmark and number of operations is based on budget estimates made by Knowledge Centre for Agriculture. The uncertainty is assumed to be 10 %.

The uncertainties for the PM emission factors have been considered to be very high and especially for animal husbandry and manure management. The uncertainty estimates regarding the PM emission factors for manure management and farm level agricultural operations are based on the EMEP/EEA guidebook.

### **Other pollutants**

For both the NO<sub>x</sub> and NMVOC emission the activity data is based on the same conditions as mentioned in NH<sub>3</sub> chapter and therefore the same uncertainty estimates is used.

The uncertainty for the NO<sub>x</sub> and NMVOC emission factor is based on expert judgment and is considered to be very high; 100 - 500 % based on the on the EMEP/EEA guidebook.

Emission of BC, CO, SO<sub>2</sub>, heavy metals, dioxin, PAHs, HCB and PCB from the agricultural sector originates from field burning of agricultural residues. The uncertainty for activity data for these emissions is a combination of the uncertainty for crop production which is low and the uncertainty of the amount of burned straw which is high. The uncertainties for the emission factors are based on EMEP/EEA guidebook. All uncertainties for field burning are relatively high. The uncertainty for activity data for the emission of HCB from pesticides are estimated to 5 % and the uncertainty for the emission factor are relatively high.



Table 5.30 Estimated uncertainty associated with activities and emission factors for the agricultural sector.

Compound	NFR sector	Emission	Activity data, %	Emission factor, %	Combined Uncertainty, %	Total Uncertainty, %
NH <sub>3</sub> , Gg	3.B Manure management	36.31	5	25	25	24
	3.Da1 Inorganic fertilizers	7.29	3	25	25	
	3.Da2a Animal manure applied	19.53	15	25	29	
	3.Da2b Sewage sludge applied	0.06	15	50	52	
	3.Da3 Deposited by grazing	1.86	5	25	25	
	3.De Cultivated crops	5.37	2	50	50	
	3.F Field burning	0.11	25	50	56	
	3.G Agriculture other	NO	20	50		
TSP, Gg	3.B Manure management	10.23	7	300	300	257
	3.Dc Farm-level agri. operations	55.22	10	300	300	
	3.F Field burning	0.27	25	50	56	
PM <sub>10</sub> , Gg	3.B Manure management	5.52	7	300	300	207
	3Dc Farm-level agri. operations	5.52	10	300	300	
	3.F Field burning	0.27	25	50	56	
PM <sub>2.5</sub> , Gg	3.B Manure management	1.31	2	300	300	254
	3Dc Farm-level agri. operations	0.45	10	300	300	
	3.F Field burning	0.26	25	50	56	
NMVOC, Gg	3 B Manure management	36.13	2	300	300	284
	3.De Cultivated crops	1.83	5	500	500	
	3.F Field burning	0.29	25	100	103	
NO <sub>x</sub> , Gg	3.B Manure management	0.22	5	100	100	214
	3.Da1 Inorganic fertilizers	7.72	3	400	400	
	3.Da2a Animal manure applied	6.07	15	400	400	
	3.Da2b Sewage sludge applied	0.28	20	400	400	
	3.F Field burning	0,11	25	25	36	
HCB, kg	3.F Field burning	0.12	25	500	501	501
HCB, kg	3 G Agriculture other	0.02	5	500	500	
PCB, kg	3.F Field burning	<0.01	25	500	501	501
SO <sub>2</sub> , Gg	3.F Field burning	0.01	25	100	103	103
BC, Gg	3.F Field burning	0.02	25	100	103	103
CO, Gg	3.F Field burning	2.75	25	100	103	103
Pb, Mg	3.F Field burning	0.04	25	50	56	56
Cd, Mg	3.F Field burning	<0.01	25	100	103	103
Hg, Mg	3.F Field burning	<0.01	25	200	202	202
As, Mg	3.F Field burning	<0.01	25	100	103	103
Cr, Mg	3.F Field burning	0.01	25	200	202	202
Cu, Mg	3.F Field burning	<0.01	25	200	202	202
Ni, Mg	3.F Field burning	0.01	25	200	202	202
Se, Mg	3.F Field burning	<0.01	25	100	103	103
Zn, Mg	3.F Field burning	<0.01	25	200	202	202
Dioxin, g I-Teq	3.F Field burning	0.03	25	500	501	501
Benzo(a)pyrene, Mg	3.F Field burning	0.13	25	500	501	501
Benzo(b)fluoranthene, Mg	3.F Field burning	0.13	25	500	501	501
Benzo(k)fluoranthene, Mg	3.F Field burning	0.05	25	500	501	501
Indeno(1,2,3 cd)pyrene, Mg	3.F Field burning	0.05	25	500	501	501

## 5.8 Quality assurance and quality control (QA/QC)

A general QA/QC and verification plan for the agricultural sector is continuously under development and will be improved and developed in line with the deficiencies are identified and corrected. The objectives for the quality planning, as given in the CLRTAP Emission Inventory Guidebook,

which is closely related to the IPCC Good Practice Guidance, are to improve the transparency, consistency, comparability, completeness and confidence.

To ensure consistency a procedure for internal quality check are provided. Input of external data is checked and certain time series have been prepared for both the activity data, the emission factors and implied emission factors, 1985 - 2013. The annual change for each emission source on activity will be checked for significant differences and if necessary explained. Considerable variation between years can reveal miscalculations or changes in methods. All checks of all activity data, emission factor, implied emission factor and other important key parameters are provided and achieved in excel spread sheet.

Activity data and emission factors are collected and discussed in cooperation with specialists and researchers at different institutes and research departments. As a consequence, both data and methods are evaluated continuously according to latest knowledge and information. A more detailed description of quality assurance and quality control is given in the Denmark's National Inventory Report 2014 - submitted under the United Nations Framework Convention on Climate Change (<http://dce2.au.dk/pub/SR101.pdf>).

## 5.9 Recalculations

Compared with the previous NH<sub>3</sub>, NMVOC and PM emissions inventory (submission 2014), some changes and updates have been made, see Table 5.31. These changes cause a low (less than 1 %) decrease/increase in the total NH<sub>3</sub> emission for all years (1985–2012) and a very high increase in the PM emission 2000-2012 by more than 550 % due to implementation of new sectors.

Table 5.31 Changes in NH<sub>3</sub>, NMVOC and PM emission in the agricultural sector compared to NFR reported last year.

NH <sub>3</sub> emission, Gg NH <sub>3</sub>	1985	1990	1995	2000	2005	2010	2011	2012
2014 submission	125.98	123.74	106.85	93.16	83.85	75.53	74.70	73.43
2015 submission	125.97	123.71	106.83	93.15	83.84	75.52	74.72	73.30
Difference, %	-0.01	-0.02	-0.02	-0.01	-0.01	-0.002	0.03	-0.19

NMVOC emission, Gg	1990	1995	2000	2005	2010	2011	2012
2014 submission	2.10	2.00	1.98	2.10	2.160	2.15	2.21
2015 submission	40.20	38.69	37.71	36.83	37.26	37.18	37.78
Difference, %	1 811	1 839	1 808	1 655	1 625	1 630	1 610

PM emission, Gg TSP	2000	2005	2010	2011	2012
2014 submission	11.11	11.50	11.16	10.98	10.58
2015 submission	78.85	79.88	73.65	74.39	74.43
Difference, %	610	595	560	577	604

For the total emission of NH<sub>3</sub> the changes are very low, but due to restructuring of the NFR tables some changes have been made for some sectors. Emission of NH<sub>3</sub> from manure management (NFR 3B) is decreased because the emission from manure applied to soils is now reported under NFR 3D. The emission of NH<sub>3</sub> from sewage sludge applied to soils and cultivated crops are now reported under sector 3D instead of under 3I.

The emission of NMVOC has increased a lot due to implementation of NMVOC for manure management (3B) for the years 1985 to 2013. The emission factors for NMVOC emission from cultivated crops (3D) has been updated to the emission factors given in the EMEP/EEA 2013 Guidebook.

The emission of PM has increased a lot due to implementation of emission from farm-level agricultural operations including storage, handling, and transport of agricultural products. The emission from manure management has only changes up to 2 % due to change in number of animals.

The emission of NO<sub>x</sub> has been estimated for manure management (3B), inorganic N-fertilizers (3Da1) and sewage sludge applied to soils (3Da2b) for the years 1985 to 2013.

Some changes in activity data has been made; Number of animals has been changed due to updated numbers for swine in 2011, broilers in 2007-2012 and laying hens in 2012. This affects the emission of NH<sub>3</sub> and PM. Numbers of sheep has been divided in two subcategories – sheep and lambs – with emission factors for each subcategories. This increases the number of animals and affects the emission of NH<sub>3</sub> from manure management (3B) and urine and dung deposited by grazing animals (3D). N from sewage sludge has been updated for 2006 and this affects the emission of NH<sub>3</sub>. The area for field burning of agricultural residues has been changed due to updated data for 2012 and this affects emission of all pollutants.

### **5.10 Planned improvements**

In recent years, there has been focus on reduction of the NH<sub>3</sub> emission and especially the possibilities for emission reduction in housings. A number of investigations to estimate the effects from technical measures on the emission have been initiated. Until now, still relatively few housings have implemented NH<sub>3</sub> reduction technologies, but there is no doubt, that the ammonia reducing technology will play an important role in the future. When data are available, it is planned to implement the effect of the reducing technology in the emission inventory.

The QA/QC plan for the agricultural sector is continually under development. Until now, the main focus has been on the internal procedure check. There is still a need to provide the procedure for control of the inventory data calculations. This means to identify the possibility to compare the calculations made by other institutions or organisations e.g. calculation of total N-excretion made by the DCA-Danish Centre for Food and Agriculture, Aarhus University. Furthermore, it is a need to consider how to ensure a quality assurance procedure for the entire inventory.

### **5.11 References**

Aaes, Ole, 2008: Personal communication. Knowledge Centre for Agriculture.

Andersen, J.M., Sommer, S.G., Hutchings, N., Kristensen, V.F. & Poulsen, H.D., 1999: Emission af ammoniak fra landbruget – status og kilde. National Environmental Research Institute and the Danish Institute of Agricultural Sciences. (In Danish).

Black, R.R., Meyer, C.P., Touati, A., Gullett, B.K., Fiedler, H. and Mueller, J.F., 2012: Emission factors for PCDD/PCDF and dl-PCB from open burning of biomass, *Environment International*, Volume 38, Issue 1, January 2012, Pages 62-66

Bussink, D.W., 1994: Relationship between ammonia volatilisation and nitrogen fertilizer application rate, intake and excretion of herbage nitrogen by cattle on grazed swards. *Fertil. Res.* 38. 111-121.

Clausen, Eric F., 2008: Personal communication. Knowledge Centre for Agriculture.

Danish AgriFish Agency, 2013: Danmarks salg af handelsgødning 2012/2013 (Consumption of inorganic N-fertilizer in Denmark 2012/13). (In Danish). Available at: [http://naturerhverv.dk/fileadmin/user\\_upload/NaturErhverv/Files/Laendbrug/Handelsgoedning/Danmarks\\_salg\\_af\\_handelsgoedning\\_2012\\_2013.pdf](http://naturerhverv.dk/fileadmin/user_upload/NaturErhverv/Files/Laendbrug/Handelsgoedning/Danmarks_salg_af_handelsgoedning_2012_2013.pdf) (Jan. 2015).

EMEP/EEA air pollutant emission inventory guidebook, 2013: Available at: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013> (Jan. 2015)

Gyldenkærne, S. & Albrektsen, R., 2009: Report on the emission of NH<sub>3</sub> and NMVOC emission from agricultural crops. In press.

Hartung, J. & Seedorf, J. 1999: Characterisation of Airborne Dust in Livestock Housing and its Effects on Animal and Environment. International Symposium on Dust control in animal production facilities. June 1999.

Jarvis, S.C., Hatch, D.J. & Roberts, D.H., 1989a: The effects of grassland management on nitrogen losses from grazed swards through ammonia volatilization; the relationship to extral N returns from cattle. *J. Agric. Sci. Camb.* 112.205-216.

Jarvis, S.C., Hatch, D.J. & Lockyer, D.R., 1989b: Ammonia fluxes from grazed grassland annual losses from cattle production systems and their relation to nitrogen inputs. *J. Agric. Camp.* 113. 99-108.

Jenkins, B.M., 1996: Atmospheric Pollutant Emission Factors from Open Burning of Agricultural and Forest Biomass by Wind Tunnel Simulations; Final Report (3 Vols.); CARB Project A932-126; California Air Resources Board, Sacramento, California.

Lundgaard. N.H., 2006: Personal Communication: Department of Farm buildings and Machinery. Danish Agricultural Advisory Centre.

Poulsen, H.D., 2008: Baggrundstal 2008. DCA, Aarhus University. Available at: <http://anis.au.dk/forskning/sektioner/husdyrernaering-og-miljoe/normtal/> (Jan. 2015).

Poulsen, H.D., 2014: Normative figures 2000-2014. DCA, Aarhus University. Available at:

<http://anis.au.dk/forskning/sektioner/husdyrernaering-og-miljoe/normtal/> (Jan. 2015).

Poulsen, H.D., Børsting, C.F., Rom, H.B. & Sommer, S.G., 2001: Kvælstof, fosfor og kalium i husdyrgødning – normtal 2000. DJF rapport nr. 36 – husdyrbrug. The Danish Institute of Agricultural Sciences. (In Danish).

Poulsen, H.D. & Kristensen, V.F., 1998: Standards Values for Farm Manure – A revaluation of the Danish Standards Values concerning the Nitrogen, Phosphorus and Potassium Content of Manure. DIAS Report No. 7 - Animal Husbandry. Danish Institute of Agricultural Sciences.

Rasmussen, J.B., 2006: Personal Communication. Department of Farm buildings and Machinery. Danish Agricultural Advisory Centre.

Statistics Denmark, Agricultural Statistics. Available at:

[www.dst.dk/en.aspx](http://www.dst.dk/en.aspx) (Jan. 2015).

Takai, H., Pedersen, S., Johnsen, J.O., Metz, J.H.M., Grott Koerkamp, P.W.G., Uenk, G.H., Phillips, V.R., Holden, M.R., Sneath, R.W., Short, J.L., White, R.P., Hartung, J., Seedorf, J., Schröder, M., Linkert, K.H. & Wathers, C.M., 1998: Concentrations and Emissions of Airborne Dust in Livestock Buildings in Northern Europe. Journal of Agricultural Engineering Research, Volume 70 no. 1. May 1998.

VERA, 2010: Available at: [http://www.etadanmark.dk/-/media/DS/Files/Downloads/Artikler/VERA/VERA\\_erklaering\\_2012\\_0kt\\_enkeltside.pdf](http://www.etadanmark.dk/-/media/DS/Files/Downloads/Artikler/VERA/VERA_erklaering_2012_0kt_enkeltside.pdf) (Jan. 2015).

Vestergaard, A.V., 2014: Personal Communication. Knowledge Centre for Agriculture.

Yang, C., 2006: Estimating HCB Releases from Pesticide Applications, HCB Releases from Pesticide Applications in USA and Canada – All Years. Environment Canada.

## 6 Waste

The waste sector consists of the four main NFR categories 5A Solid waste disposal, 5B Biological treatment of solid waste, 5C Incineration and open burning of waste, 5D Wastewater treatment and discharge and 5E Other waste. Table 6.1 below shows the relevant SNAP codes for the waste sector.

Table 6.1 Link between SNAP codes and NFR sectors.

SNAP code	SNAP name	NFR code
090401	Managed Waste Disposal on Land	5A
090402	Unmanaged Waste Disposal Sites	5A
090403	Other	5A
091001	Wastewater treatment in industry	5D
091007	Latrines	5D
091002	Wastewater treatment in residential/commercial sector	5D
090201	Incineration of domestic or municipal wastes	5C
090202	Incineration of industrial wastes (except flaring)	5C
090204	Flaring in chemical industries	5C
090205	Incineration of sludge from waste water treatment	5C
090207	Incineration of hospital wastes	5C
090208	Incineration of waste oil	5C
090901	Incineration of corpses	5C
090902	Incineration of carcasses	5C
090700	Open burning of agricultural wastes	5C
091003	Sludge spreading	5E
091005	Compost production	5B
091006	Biogas production	5B
091008	Other production of fuel (refuse derived fuel)	5E
091009	Accidental fires	5E

Incineration of waste (municipal, industrial, clinical and hazardous) in Denmark is done with energy recovery and therefore the emissions are included under the relevant sectors under NFR sector 1A. The documentation for waste incineration is included in Chapter 3.2.

### 6.1 Solid waste disposal

Major emissions from landfilling are emissions of greenhouse gases, i.e. CH<sub>4</sub>. It is assumed that landfilling also leads to emission of small quantities of NMVOC, CO, NH<sub>3</sub> and NO<sub>x</sub>. PM emissions are emitted from waste handling as well, but these have not been included in the current submission.

Currently, Denmark has not estimated emissions of air pollutants from solid waste disposal. The EMEP/EEA Guidebook contains default NMVOC and particle emission factor, however due to limited amounts of resources it has not been possible to estimate such emissions.

### 6.2 Biological treatment of solid waste

This sector covers two activities: composting and anaerobic digestion at biogas facilities. These are described in more detail below.

### 6.2.1 Compost production

This section covers the biological treatment of solid organic waste called composting. Pollutants that are emitted during composting are CO and NH<sub>3</sub>.

#### Methodology

Emissions from composting have been calculated according to a country specific Tier 1 method. However, a Tier 1 default methodological guidance is available in the 2006 IPCC Guidelines (IPCC, 2006).

In Denmark, composting of solid biological waste includes composting of:

- garden and park waste (GPW),
- organic waste from households and other sources,
- sludge,
- home composting of garden and vegetable food waste.

In 2001, 123 composting facilities treated only garden and park waste (type 2 facilities), nine facilities treated organic waste mixed with GPW or other organic waste (type 1 facilities) and 10 facilities treated GPW mixed with sludge and/or “other organic waste” (type 3 facilities). 92 % of these facilities consisted entirely of windrow composting, which is a simple technology composting method with access to only natural air. It is assumed that all facilities can be considered as using windrow composting (Petersen & Hansen, 2003).

Composting is performed with simple technology in Denmark; this implies that temperature, moisture and aeration are not consistently controlled or regulated. Temperature is measured but not controlled, moisture is regulated by watering the windrows in respect to weather conditions and aeration is assisted by turning the windrows (Petersen & Hansen, 2003).

During composting a fraction of the degradable organic carbon (DOC) in the waste material is converted into CO. Even though the windrows are occasionally turned to support aeration, anaerobic sections are inevitable and will cause a small emission of CH<sub>4</sub>. In the same manner, aerobic biological digestion of N leads to an emission of NO<sub>x</sub>, while the anaerobic decomposition leads to the emission of NH<sub>3</sub> (IPCC, 2006).

#### Activity data

All Danish waste treatment plants are obligated to statutory registration and reporting of all waste entering and leaving the plants. All waste streams are weighed, categorised with a waste type and a type of treatment and registered to the ISAG waste information system, which contain data for 1995-2009 (ISAG, 2010). For 2010 to 2013 data was from the new waste data system was obtained from the Danish EPA. For this year reporting, the total amount of composted waste was allocated according to the fractional distribution between the four waste types in 2009. The new waste data system (WDS) replaces ISAG from 2010 and forward. Activity data for 2010-2013 are provided in table x. Based on the approach of allocation the total amount of waste composted in 2010 to 2013, no visible data break are observed in the time series as visualised in Figure 6.1. It should however be mentioned that work is ongoing in regard to reallocating composted amounts according to EWC level 3 description, why adjustments in data and methodology may occur in the reporting year 2015.

Figure 6.1 illustrates the composted amount of waste divided in the four categories mentioned earlier.




Figure 6.1 Amount of composted waste divided in garden and park waste (GPW), organic municipal solid waste (MSW), sludge and home composting of garden and food waste, these data are also shown in Table 6.3.

Activity data for the years 1995-2009 are collected from the ISAG database for the categories: "sludge", "organic waste from households and other sources" and "garden and park waste". Activities for 2010-2013 are collected from the new WDS.

The Danish legislation on sludge (DEPA, 2006) was implemented in the summer of 2003. This stated that composted sludge may only be used as a fertilizer on areas not intended for growing foods of any kind for at least 2-3 years. This restriction caused the amount of composted sludge to drop drastically from 2003 to 2004.

The trend in composting of sludge does not demonstrate a convincing trend that can be used for estimation of activity data for previous years. Since this activity is insignificant for 1995-1997 (1-2 %) it is assumed to be "not occurring" for 1985-1994.

The amount of organic waste from households composted in the years 1985-1994 is estimated by multiplying the number of facilities treating this type of waste with the average amount composted per facility in the years 1995-2001 (2.6-3.8 Gg per facility per year). Table 6.2 shows the number of composting plants grouped into plant types composting organic waste mixed with GPW (type1), GPW only (type 2) and GPW mixed with sludge and/or "other organic waste" (Type 3) described in the methodology section above.



Table 6.2 Number of composting facilities in the years 1985-2001.

Plant type*	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
Type 1	2	2	3	3	4	5	6	7	8	9
Type 2	6	10	14	18	22	38	54	70	86	102
Type 3	0	0	0	0	0	1	2	2	3	4
Total	8	12	17	21	26	44	62	79	97	115

*Continued*

Facility type	1995	1996	1997	1998	1999	2000	2001
Type 1	13	14	13	14	13	11	9
Type 2	113	108	99	102	111	115	123
Type 3	9	9	11	10	10	7	10
Total	136	133	126	130	139	138	149

Type 1 waste treatment sites normally includes biogas producing facilities, but these are not included in this table.

\*Petersen, 2001 and Petersen & Hansen, 2003

The ISAG activity data for composting of GPW include wood chipping. Compost data for GPW provided by Petersen (2001) and Petersen & Hansen (2003) show that for 1997-2001, wood chipping accounts for about 3 % of the total chosen ISAG activity data for GPW. Activity data for GPW for the years 1985-1994 and 2010-2013 are estimated by extrapolating the trend.

The last waste type involved in composting is home composting of garden waste and vegetable waste. The activity data for this category are known from Petersen & Kielland (2003) to be 21.4 Gg in 2001. It is assumed that the following estimates made by Petersen & Kielland (2003) are valid for all years 1985-2013:

- 28 % of all residential buildings with private gardens (including summer cottages) are actively contributing to home composting.
- 14 % of all multi-dwelling houses are actively contributing to home composting.
- 50 kg waste per year will on average be composted at every contributing residential building.
- 10 kg waste per year will on average be composted at every contributing multi-dwelling house.

Multi-dwelling houses include apartment buildings and it is very uncommon for people in these types of buildings to compost their biowaste and the average amount of composted waste is therefore lower in spite of the higher number of residents. Statistics on the total number of occupied residential buildings, summer cottages and multi-dwelling houses are available at the Statistics Denmark's website.

The calculated activity data for home composting of garden and vegetable waste are shown in Table 6.3 and Annex 3E-5.

Table 6.3 Activity data composting, Gg.

	1985	1990	1995	2000	2005	2010	2011	2012	2013
Composting of garden and park waste	130	288	376	677	737	648	816	751	1061
Composting of organic waste from households and other sources	5	16	40	47	45	54	67	62	88
Composting of sludge	NO	NO	7	218	50	82	103	95	134
Home composting of garden and vegetable food waste	19	20	21	21	22	18	22	20	29
Total	154	324	444	963	854	800	1008	929	1312

NO = Not occurring.

### Emission factors

The emissions from composting strongly depend on both the composition of the treated waste and on process conditions such as aeration, mechanical agitation, moisture control and temperature pattern (Amlinger et al., 2008).

The emission factors provided in Table 6.4 are considered the best available for the calculation of Danish national emissions from composting.

Table 6.4 Composting emission factors, per Mg.

	Composting of garden and park waste (GPW)	Composting of organic waste	Composting of sludge	Home composting of garden and vegetable food waste
Unit	Kg	Kg	kg	kg
NO <sub>x</sub>	NAV	NAV	NAV	NAV
CO	0.56	NAV	NAV	0.08
NH <sub>3</sub>	0.66	0.24	0.31	0.63
Source	Boldrin et al., 2009	EEA, 2009	MST, 2013	Boldrin et al., 2009

Emissions from Boldrin et al. (2009) are given in percentage of total degraded carbon or nitrogen respectively. The factors shown in Table 6.4 are calculated by assuming 37.5 % DOC in dry matter, 2 % N in dry matter and 50 % moisture in the waste (Boldrin et al., 2009).

Boldrin et al. (2009) and MST (2013) do not directly provide any emission factors, the following assumptions were made to derive the factors shown in Table 6.4:

- 0.5 % N per dry matter waste water sludge
- 25 % moisture in waste water sludge.
- 2 % N per dry matter garden waste (incl. home composting)
- DOC is 25-50 % in garden waste (incl. home composting)
- 50 % moisture in garden waste (incl. home composting)

### Emissions

Table 6.5 show the total national emissions from composting. The full time series is shown in Annex 3E-6.

Table 6.5 National emissions from composting, Mg.

	1985	1990	1995	2000	2005	2010	2011	2012	2013
CO	74.6	163.5	213.1	382.4	416.2	365.6	460.5	424.2	599.0
NH <sub>3</sub>	99.0	206.5	273.2	538.8	526.5	476.6	600.4	553.1	781.0

## 6.2.2 Biogas production

Emissions from biogas production are divided and reported in different sectors according to waste type and method.

### Methodology

Emissions from the combustion of biogas regardless of the origin are included in the energy sector and are allocated to the appropriate subsector in the Danish energy statistics. See this IIR Chapter 3, Energy.

The reduced emissions from biogasified versus raw manure spread on agricultural soils are described in the agricultural sector in Chapter 5.

Fugitive emissions of NMVOC and NH<sub>3</sub> from anaerobic digestion of sludge from wastewater treatment should be included in the NFR source category 5D Wastewater treatment and discharge. However, NMVOC and NH<sub>3</sub> emission from anaerobic digestion of sludge from wastewater treatment are not presently included in the submission but should be investigated and possibly added to this chapter to the extent that emissions originates from the digester tank, while the emissions originating from wastewater treatment processes should be described and included in Chapter 6.4.

Emissions that may be presented in this section in the future include fugitive emissions from the digester tank at the wastewater treatment plants treating the sludge by anaerobic digestion and emissions from combustion of biogas at the biogas production plants (own production and use).

Fugitive emissions from anaerobic digestion of biowaste is to be included in this chapter, but fugitive emissions from storage, pre- and post-treatment of the digestate should be included in the relevant chapters as appropriate.

Emissions from combustion of biogas at site of production are included for the years 1994-2004 for which years these are reported separately in the energy statistics produced by the Danish Energy Agency. This activity is not reported from 2005 and forward. Pollutants from this activity are SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, particulate matter, heavy metals, HCB, PCDD/F and PCBs.

#### Activity data

Activity data for this source category are collected from the energy statistics (DEA, 2013). Combustion of biogas at biogas production plants occurred for the years 1994-2004, the full time-series is available in Annex 3E Table 3E-7.

Table 6.6 Combusted biogas at biogas production plants

	1995	2000	2004
GJ	4711	40990	28744




Figure 6.2 Combusted biogas at biogas production plants.

#### Emission factors

Emission factors for combustion of biogas at biogas production plants are presented in Table 6.7.

Table 6.7 Emission factors for combustion of biogas, per GJ.

Pollutant	Unit	Emission factor
SO <sub>2</sub>	g	25
NO <sub>x</sub>	g	28
NMVOG	g	2
CO	g	36
TSP	g	1.5
PM <sub>10</sub>	g	1.5
PM <sub>2.5</sub>	g	1.5
As	mg	0.04
Cd	mg	0.002
Cr	mg	0.18
Cu	mg	0.31
Hg	mg	0.12
Ni	mg	0.23
Pb	mg	0.005
Se	mg	0.21
Zn	mg	3.95
HCB	ng	190
PCDD/F	ng	0.025
PCBs	ng	90

### Emissions

Table 6.8 shows the total national emissions from combustion of biogas at biogas production plants. This activity is reported separately for the time period 1994 to 2004, the full time-series is available in Annex 3E Table 3E-8.

Table 6.8 Emissions from the combustion of biogas at biogas production plants.

		1995	2000	2004
SO <sub>2</sub>	Mg	0.1	1.0	0.7
NO <sub>x</sub>	Mg	0.1	1.1	0.8
NMVOG	Mg	0.01	0.1	0.1
CO	Mg	0.2	1.5	1.0
TSP	kg	7.1	61.5	43.1
PM <sub>10</sub>	kg	7.1	61.5	43.1
PM <sub>2.5</sub>	kg	7.1	61.5	43.1
As	g	0.2	1.6	1.1
Cd	g	0.01	0.1	0.1
Cr	g	0.8	7.4	5.2
Cu	g	1.5	12.7	8.9
Hg	g	0.6	4.9	3.4
Ni	g	1.1	9.4	6.6
Pb	g	0.02	0.2	0.1
Se	g	1.0	8.6	6.0
Zn	g	18.6	161.9	113.5
HCB	mg	0.9	7.8	5.5
PCDD/F	µg	0.1	1.0	0.7
PCBs	mg	0.4	3.7	2.6

### 6.3 Waste incineration

Incineration of municipal, industrial, clinical and hazardous waste takes place with energy recovery, therefore the emissions are included in the relevant subsectors under NFR sector 1A. For documentation please refer to Chapter 3.2. Flaring off-shore and in refineries are included under NFR sec-

tor 1B2c, for documentation please refer to Chapter 3.4. No flaring in chemical industry occurs in Denmark.

### 6.3.1 Human cremation

The incineration of human corpses is a common practice that is performed on an increasing percentage of the deceased. All Danish crematoria use optimised and controlled cremation facilities, with temperatures reaching 800-850 °C, secondary combustion chambers, controlled combustion air flow and regulations for coffin materials.

However, the emissions of especially Hg caused by cremations can still contribute to a considerable part of the total national emissions. In addition to the most frequently discussed emissions of Hg and PCDD/Fs (dioxins and furans), are the emissions of compounds like SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, other heavy metals (As, Cd, Cr, Cu, Ni, Pb, Se, Zn), particulate matter, HCB, PAHs and PCBs.

Crematoria are usually located within cities, close to residential areas and normally, their stacks are relatively low. Therefore environmental and human exposure is likely to occur as a result of emissions from cremation facilities.

#### Methodology

During the 1990es all Danish crematoria were rebuilt to meet new standards. This included installation of secondary combustion chambers and in most cases, replacement of old primary combustion chambers (Schleicher et al., 2001). All Danish crematoria are therefore performing controlled incinerations with a good burn-out of the gases, and a low emission of pollutants.

Following the development of new technology, the emission limit values for crematoria were lowered again in January 2011. These new standards were originally expected from January 2009 but were postponed two years for existing crematoria.

Table 6.9 shows a comparison of the emission limit values from February 1993 and the new standard limits.

Table 6.9 Emission limit values mg per Nm<sup>3</sup> at 11 % O<sub>2</sub>.

Component	1993 standard*	2011 standard**
Total dust	80	10
CO	50	50
Hg	No demands	0.1
Other demands:		
Stack height	3 m above rooftop	3 m above rooftop
Temperature in stack	Minimum 150 °C	Minimum 110 °C
Flue gas flow in stack	8 – 20 m/s	No demands
Temperature in after burner	850 °C	800 °C
Residence time in after burner	2 seconds	2 seconds
Odour	The crematory must not cause noticeable odour in the surroundings	The crematory must not cause odour nuisance outside the crematory perimeter, that is significant according to the supervisory authority

\* Schleicher et al., 2001;\*\*Schleicher & Gram, 2008.

To meet the new standards, some crematoria have been rebuilt to larger capacity while others are closed (MILIKI, 2006). In 2011, there were 26 operating crematoria in Denmark, some with multiple furnaces (DKL, 2014).

Crematoria that are not closed are equipped with flue gas cleaning (bag filters with activated carbon). The use of air pollution control devices, and activated carbon, for the removal of Hg will also reduce the flue gas concentration of dioxins, PAHs and odour. Existing knowledge on the reduction efficiencies justifies are presented in Schleicher & Gram (2008).

Around half of the Danish crematoria are currently connected to the district heating system and in addition, a few crematoria produce heat for use in their own buildings. The bag filter cleaning system requires that the flue gas is cooled down to 125-150 °C, and the cheapest way to do so is to use the surplus heat in the district heating system (DKL, 2009). The heat contribution from crematoria is negligible compared to the total district heat production and is not part of the Danish energy statistics.

### Activity data

Table 6.10 shows the time series of total number of deceased persons (Statistics Denmark, 2014), number of cremations and the fraction of cremations in relation to the total number of deceased (DKL, 2014). Annex 3E Table 3E-1 presents data for the entire time series.

Table 6.10 Data human cremations (DKL 2014, Statistics Denmark 2014).

	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
Nationally deceased	55939	58378	60926	63127	57998	54962	54368	52516	52325	52471
Cremations	33986	36705	40991	43847	41651	40758	42050	41248	40909	42349
Cremation fraction, %	60.8	62.8	67.3	69.5	71.8	74.2	77.3	78.6	79.6	80.7




Figure 6.3 Illustration of the development in cremations (DKL 2014), where the number of cremations,  $N_{\text{cremations}}$ , is shown at the left Y-axis. The cremation percentage,  $F_{\text{cremations}}$ , shows the percentage of cremated deceased of the total number of deceased for the years 1984 to 2013. Data for 1980-1983 are estimated values, for details on the estimation, see Annex 3E-1.

Even though the total number of annual cremations is fluctuating, the cremation percentage has been steadily increasing since 1984, and is likely to continue to increase.




Figure 6.4 Trends of the activity data for cremation of human corpses and the number of deceased persons.

Figure 6.4 presents the trend of the number of deceased persons together with the activity data for human cremation. The figure shows a direct connection between the number of deceased and the activity of human cremation as the two trends are quite similar. Figure 6.4 also shows the effect of the increasing fraction of cremations per deceased, as the number of cremations is not decreasing along with the number of deceased. The percentage of the deceased being cremated has increased from 67 % in 1990 to 81 % in 2013 as shown in Figure 6.3, Table 6.10 and Annex 3E Table 3E-1.

#### Emission factors

For crematoria, emissions are calculated by multiplying the total number of cremations by the emission factors. The emission factors are gathered from literature and are based on the measurements performed in countries that are comparable with Denmark. By comparable is meant countries that use similar incineration processes, similar cremation techniques including support fuel and have a similar composition of sources to lifetime exposure, lifetimes and coffins.

Table 6.11 lists the emission factors in the time period 1980-2010 and their respective references. As mentioned earlier, 2011 is year one after installation of bag filters with activated carbon at all Danish crematoria, causing the emission factors for particles, heavy metals, PAHs and PCDD/Fs to decrease quite drastically (Schleicher & Gram, 2008).

Table 6.11 Emission factors for human cremation with references.

Pollutant name	Unit	Emission factor <sup>*</sup>	Reference
SO <sub>2</sub>	kg/body	0.113	Santarsiero et al., 2005
NO <sub>x</sub>	kg/body	0.825	Santarsiero et al., 2005
NMVOG	kg/body	0.013	EEA, 1996
CO	kg/body	0.010	Schleicher et al., 2001
NH <sub>3</sub>		NA	
TSP	kg/body	0.039	Webfire, 2012
PM <sub>10</sub>	kg/body	0.035	Webfire, 2012
PM <sub>2.5</sub>	kg/body	0.031	Webfire, 2012
As	g/body	0.014	Webfire, 2012
Cd	g/body	0.005	Webfire, 2012
Cr	g/body	0.014	Webfire, 2012
Cu	g/body	0.012	Webfire, 2012
Hg	g/body	1.12	Kriegbaum et al., 2005
Ni	g/body	0.017	Webfire, 2012
Pb	g/body	0.030	Webfire, 2012
Se	g/body	0.020	Webfire, 2012
Zn	g/body	0.160	Webfire, 2012
HCB	mg/body	0.152	Toda, 2006
PCDD/F	µg I-TEQ/body <sup>**</sup>	0.350	Schleicher et al., 2001
Benzo(b)flouranthene	µg/body	7.21	Webfire, 2012
Benzo(k)flouranthene	µg/body	6.44	Webfire, 2012
Benzo(a)pyrene	µg/body	13.20	Webfire, 2012
Indeno(1,2,3-c-d)pyrene	µg/body	6.99	Webfire, 2012
PCBs	mg/body	0.414	Toda, 2006

<sup>\*</sup>NA = not applicable. <sup>\*\*</sup> I-TEQ: International Toxicity Equivalents.

The average body weight of cremated corpses is assumed to be 65 kg.

Flue gas cleaning efficiencies are based on measurements performed at Danish crematoria and expert judgements, and set equal to 99 % for PCDD/Fs, particles, PAHs and heavy metals. These abatement efficiencies are implemented from 2011. For all other pollutants the emission factors are as listed in Table 6.11.

It has not been possible to find data for ammonia. Ammonia might appear in lesser amounts, but will most likely be converted to NO<sub>x</sub> at the high incineration temperatures.

There might for some emission factors be included a small part of the support fuel (natural gas) if the measurements were taken early in the burning process. This would then be a double counting since fuel for cremation is reported under NFR code 1A4a, commercial and institutional. However, this double counting is considered miniscule.

### Emissions

Table 6.12 shows the total emissions from selected years. To view the entire time series 1980-2013, see Annex 3E Table 3E-3a-d. The dioxin emission is given in I-TEQ; i.e. International Toxicity Equivalents which is a weighted addition of congener toxicity with reference to 2,3,7,8-TCDD (Seveso-dioxin).

Emissions from human cremations have been steady over the last two decades but have decreased strongly for the pollutants TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, As, Cd,



Cr, Cu, Hg, Ni, Pb, Se, Zn, PCDD/Fs and PAHs from 2010 to 2011 because of the installation of bag filters with activated carbon.

Table 6.12 Total national emissions from human cremations.

	Unit	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
SO <sub>2</sub>	Mg	3.83	4.14	4.62	4.94	4.70	4.60	4.74	4.65	4.61	4.77
NO <sub>x</sub>	Mg	28.04	30.28	33.82	36.17	34.36	33.63	34.69	34.03	33.75	34.94
NMVOG	Mg	0.442	0.477	0.533	0.570	0.541	0.530	0.547	0.536	0.532	0.551
CO	Mg	0.340	0.367	0.410	0.438	0.417	0.408	0.421	0.412	0.409	0.423
TSP	Mg	1.31	1.42	1.58	1.69	1.61	1.57	1.62	0.02	0.02	0.02
PM <sub>10</sub>	Mg	1.18	1.27	1.42	1.52	1.45	1.41	1.46	0.01	0.01	0.01
PM <sub>2.5</sub>	Mg	1.18	1.27	1.42	1.52	1.45	1.41	1.46	0.01	0.01	0.01
As	kg	0.46	0.50	0.56	0.60	0.57	0.55	0.57	0.01	0.01	0.01
Cd	kg	0.17	0.18	0.21	0.22	0.21	0.21	0.21	0.002	0.002	0.002
Cr	kg	0.46	0.50	0.56	0.59	0.56	0.55	0.57	0.01	0.01	0.01
Cu	kg	0.42	0.46	0.51	0.55	0.52	0.51	0.52	0.01	0.01	0.01
Hg	kg	38.03	41.07	45.87	49.06	46.61	45.61	47.05	0.46	0.46	0.47
Ni	kg	0.59	0.64	0.71	0.76	0.72	0.71	0.73	0.01	0.01	0.01
Pb	kg	1.02	1.10	1.23	1.32	1.25	1.22	1.26	0.01	0.01	0.01
Se	kg	0.67	0.73	0.81	0.87	0.82	0.81	0.83	0.01	0.01	0.01
Zn	kg	5.44	5.88	6.56	7.02	6.67	6.53	6.73	0.07	0.07	0.07
HCB	g	5.15	5.56	6.21	6.65	6.31	6.18	6.37	6.25	6.20	6.42
PCDD/F	mg	11.90	12.85	14.35	15.35	14.58	14.27	14.72	0.14	0.14	0.15
benzo(b)flouranthene	g	0.25	0.26	0.30	0.32	0.30	0.29	0.30	0.003	0.003	0.003
benzo(k)flouranthene	g	0.22	0.24	0.26	0.28	0.27	0.26	0.27	0.003	0.003	0.003
benzo(a)pyrene	g	0.45	0.48	0.54	0.58	0.55	0.54	0.56	0.005	0.005	0.006
indeno(1,2,3-c-d)pyrene	g	0.24	0.26	0.29	0.31	0.29	0.28	0.29	0.003	0.003	0.003
PCB	g	14.05	15.18	16.95	18.13	17.22	16.86	17.39	17.06	16.92	17.51

### 6.3.2 Animal cremation

The incineration of animal carcasses in animal crematoria follows much the same procedure as human cremation. Animal crematoria use similar two chambered furnaces and controlled incineration. However, animal carcasses are burned in special designed plastic (PE) bags rather than coffins. Emissions from animal cremation are similar to those from human cremation, with the exception of Hg which mainly stems from amalgam tooth fillings.

Animal cremations are performed in two ways, individually where the owner often pays for receiving the ashes in an urn or collectively which is most often the case with animal carcasses that are left at the veterinarian.

#### Methodology

Open burning of animal carcasses is illegal in Denmark and is not occurring and small-scale incinerators are not known to be used at Danish farms. Livestock that is diseased or in other ways unfit for consumption is disposed of through rendering plants, incineration of livestock carcasses is illegal and these carcasses are therefore commonly used in the production of fat and soap at Daka Bio-industries.

The only animal carcasses that are approved for cremation in Denmark are deceased pets and animals used for experimental purposes, where the incineration must take place at a specialised animal crematorium. There are four animal crematoria in Denmark but one of these is situated at a waste incineration company in northern Jutland called AVV. The specially designed

cremation furnaces are at this location connected to the flue gas cleaning equipment of the municipal waste incineration plant with energy recovery and the emission from the cremations are therefore included in the annual inventory from AVV. Consequently, this crematorium is included in Chapter 3.2 Stationary combustion. Therefore only three animal crematoria are included in this sector.

Animal by-products are regulated under the EU commission regulation no. 142/2011. This states that animal crematoria must be approved by the authority and comply either with the EU directive (2000/76/EC) on waste incineration or with Regulation (EC) No. 1069/2009. (EC, 2011).

The incineration of animal carcasses is, as the incineration of human corpses, performed in special incineration chambers. All Danish animal crematoria have primary combustion chambers with temperatures around 850 °C and secondary combustion chambers with temperatures around 1100 °C. The support fuel used at the Danish facilities is natural gas.

Emissions from pet cremations are calculated for SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, NH<sub>3</sub>, particles, heavy metals (As, Cd, Cr, Cu, Ni, Pb, Se, Zn), HCB, dioxins/furans, PAHs and PCBs. For the pollutants SO<sub>2</sub>, NO<sub>x</sub>, CO, As, Se, HCB, PAHs and PCBs, emissions are estimated by using the same emission factors as for human cremation.

#### Activity data

Activity data for animal cremation are gathered directly from the animal crematoria. There is no national statistics available on the activity from these facilities. The precision of activity data therefore depends on the information provided by the crematoria.

The following Table 6.13 lists the four Danish pet crematoria, their foundation year and provides each crematorium with an id letter.

Table 6.13 Animal crematoria in Denmark.

Id	Name of crematorium	Founded in
A	Dansk Dyrekremering ApS	May 2006
B	Ada's Kæledyrskrematorium ApS	Unknown, existed in more than 30 years, assumed 1980
C	Kæledyrskrematoriet	2006
D	Kæledyrskrematoriet v. Modtagestation Vendsyssel I/S	-

Crematoria D is situated at the AVV municipal waste incineration site and the emissions from this site are, as previously mentioned, included in the annual emission reporting from AVV and consequently included in the energy sector as waste incineration with energy recovery. Therefore, only crematoria A-C are considered in this chapter.

Table 6.14 lists the activity data for crematoria A-C. The entire dataset for 1980-2013 is available in Annex 3E Table 3E-2.

Table 6.14 Activity data. Source: direct contact with all Danish crematoria.

	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
Total, Mg	50	100	150	200	443	762	1449	1219	1238	1148

Crematorium B delivered exact annual activity data for the years 1998-2013. They were not certain about the founding year but believe to have existed since the early 1980es. It is assumed that crematorium B was founded at January 1<sup>st</sup> 1980 and activity data for 1980-1997 must therefore be estimated.

Statistical data describing the national consumption for pets including food and equipment for pets were evaluated as surrogate data. These statistical data show an increase of consumption of 6 % from 1998 to 2000, in the same period the amount of cremated animal carcasses increased with 89 % and no correlation seems to be present. Since there are no other available data on the subject of pets, it is concluded that there are no surrogate data available.

It is not possible to extrapolate data linearly back to 1980 because the activity, due to the steep increase, in this case would become negative from 1993 and back in time.

The activity data for animal cremation for the period of 1980-1997 are estimated by expert judgement. The estimated data are shown in Table 6.14, Figure 6.5 and Annex 3E Table 3E-2.




Figure 6.5 The amount of animal carcasses cremated, in Mg. Data from 1998-2013 are delivered by the crematoria and is considered to be exact; these data are marked as points. Data from 1980-1997 are estimated and are shown as the thick line in the figure.

### Emission factors

Concerning the incineration of animal carcasses in animal crematoria there is not much literature to be found. The EMEP/EEA Guidebook (EEA, 2009) is the only available source to emission factors for NMVOC, NH<sub>3</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and PCDD/F.

Chen et al. (2004) is the only available source to emission factors for the heavy metals Cd, Cr, Cu, Ni, Pb and Zn.

There is a good agreement between the emission factors for animal and human cremation for PCDD/F and a relatively good agreement for NMVOC, TSP, PM<sub>10</sub>, PM<sub>2.5</sub> and heavy metals.

The emission factors of the remaining pollutants SO<sub>2</sub>, NO<sub>x</sub>, CO, As, Se, HCB, PAHs and PCBs are collected from the literature search on human crema-

tion, and it is assumed that humans and animals are similar in composition for this purpose. Emission factors from human cremation are recalculated to match the activity data for animal cremation, emission per Mg.

No data were available for the emission of Hg in animal cremations. The Hg emission factor for human cremation is not transferable to animal cremations, because the Hg emission from human cremations primarily stems from tooth fillings.

Table 6.15 lists the emission factors and their respective references.

Table 6.15 Emission factors for animal cremation with references, per Mg.

Pollutant	Unit	Emission factor	Source
SO <sub>2</sub>	kg	1.73*	Santarsiero et al, 2005
NO <sub>x</sub>	kg	12.69*	Santarsiero et al, 2005
NMVOG	kg	2.00	EEA, 2009
CO	kg	0.15*	Schleicher et al., 2001
NH <sub>3</sub>	kg	1.90	EEA, 2009
TSP	kg	2.18	EEA, 2009
PM <sub>10</sub>	kg	1.53	EEA, 2009
PM <sub>2.5</sub>	kg	1.31	EEA, 2009
As	g	0.21*	Webfire, 2012
Cd	g	0.01	Chen et al., 2004
Cr	g	0.07	Chen et al., 2004
Cu	g	0.02	Chen et al., 2004
Hg	-	NAV	-
Ni	g	0.06	Chen et al., 2004
Pb	g	0.18	Chen et al., 2004
Se	g	0.30*	Webfire, 2012
Zn	g	0.19	Chen et al., 2004
HCB	mg	2.33*	Toda, 2006
PCDD/F	µg I-TEQ	10.00	EEA, 2009
Benzo(b)fluoranthene	mg	0.11*	Webfire, 2012
Benzo(k)fluoranthene	mg	0.10*	Webfire, 2012
Benzo(a)pyrene	mg	0.20*	Webfire, 2012
Indeno(1,2,3-c-d)pyrene	mg	0.11*	Webfire, 2012
PCB	mg	6.36*	Toda, 2006

\* Emission factors from human cremations.

### Emissions

For the incineration of animal carcasses, emissions are calculated by multiplying the amount of incinerated animals by the emission factors.

Emissions are summarised in Table 6.16, while emissions for the full time series are shown in Annex 3E Table 3E-4.

Table 6.16 Emissions from animal cremation.

	unit	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
SO <sub>2</sub>	Mg	0.09	0.17	0.26	0.35	0.77	1.32	2.51	2.11	2.15	1.99
NO <sub>x</sub>	Mg	0.63	1.27	1.90	2.54	5.63	9.68	18.39	15.47	15.71	14.57
NMVOOC	Mg	0.10	0.20	0.30	0.40	0.89	1.52	2.90	2.44	2.48	2.30
CO	Mg	0.01	0.02	0.02	0.03	0.07	0.12	0.22	0.19	0.19	0.18
NH <sub>3</sub>	Mg	0.10	0.19	0.29	0.38	0.84	1.45	2.75	2.32	2.35	2.18
TSP	Mg	0.11	0.22	0.33	0.44	0.97	1.66	3.16	2.66	2.70	2.50
PM <sub>10</sub>	Mg	0.08	0.15	0.23	0.31	0.68	1.17	2.22	1.86	1.89	1.76
PM <sub>2.5</sub>	Mg	0.07	0.13	0.20	0.26	0.58	1.00	1.90	1.60	1.62	1.50
As	kg	0.01	0.02	0.03	0.04	0.09	0.16	0.30	0.25	0.26	0.24
Cd	kg	0.001	0.001	0.002	0.002	0.004	0.01	0.01	0.01	0.01	0.01
Cr	kg	0.004	0.01	0.01	0.01	0.03	0.05	0.10	0.09	0.09	0.08
Cu	kg	0.001	0.002	0.003	0.004	0.01	0.02	0.03	0.02	0.02	0.02
Ni	kg	0.003	0.01	0.01	0.01	0.03	0.05	0.09	0.07	0.07	0.07
Pb	kg	0.01	0.02	0.03	0.04	0.08	0.14	0.26	0.22	0.22	0.21
Se	kg	0.02	0.03	0.05	0.06	0.13	0.23	0.44	0.37	0.38	0.35
Zn	kg	0.01	0.02	0.03	0.04	0.08	0.14	0.28	0.23	0.24	0.22
HCB	g	0.12	0.23	0.35	0.47	1.03	1.78	3.38	2.84	2.89	2.68
PCDD/F	mg	0.50	1.00	1.50	2.00	4.43	7.62	14.49	12.19	12.38	11.48
benzo(b)flouranthene	g	0.01	0.01	0.02	0.02	0.05	0.08	0.16	0.14	0.14	0.13
benzo(k)flouranthene	g	0.005	0.01	0.01	0.02	0.04	0.08	0.14	0.12	0.12	0.11
benzo(a)pyrene	g	0.01	0.02	0.03	0.04	0.09	0.15	0.29	0.25	0.25	0.23
indeno(1,2,3-c-d)pyrene	g	0.01	0.01	0.02	0.02	0.05	0.08	0.16	0.13	0.13	0.23
PCB	g	0.32	0.64	0.95	1.27	2.82	4.85	9.22	7.75	7.87	7.30

## 6.4 Wastewater handling

According to the EMEP/EEA Guidebook wastewater handling can be a source for emissions of POPs, NMVOC, NH<sub>3</sub> and CO. Of these pollutants only NMVOC is thought to be significant.

For the current submission Denmark has not estimated emissions of air pollutants from wastewater handling. The EMEP/EEA Guidebook contains a default NMVOC emission factor for latrines and wastewater handling, however due to limited resources it has not been possible to estimate such emissions.

## 6.5 Other waste

This category is a catch all for the waste sector. Emissions in this category could stem from e.g. sludge spreading, accidental fires and other combustion without energy recovery.

### 6.5.1 Sludge spreading

Sludge from wastewater treatment plants is only spread out in the open with the purpose of fertilising agricultural land. Emissions that derive from this activity are included in Chapter 5.

### 6.5.2 Accidental building fires

Emissions from accidental fires are categorised under the NFR category 5E Other waste. Pollutants that are emitted from building fires include SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, heavy metals (As, Cd, Cr, Cu, Hg, Pb), particles, PCDD/F and PAHs.

## Methodology

Emissions from building fires are calculated by multiplying the number of building fires with selected emission factors. Six types of buildings are distinguished with different emission factors: detached houses, undetached houses, apartment buildings, industrial buildings, additional buildings and containers.

## Activity data

In January 2005 it became mandatory for the local authorities to register every rescue assignment in the *online data registration- and reporting system called ODIN*, ODIN is developed and run by the Danish Emergency Management Agency (DEMA, 2007).

Activity data for accidental building fires are given by ODIN (DEMA, 2012). Fires are classified in four categories: full, large, medium and small. The emission factors comply for full scale fires and the activity data are therefore recalculated as a full scale equivalent where it is assumed that a full, large, medium and a small scale fire leads to 100 %, 75 %, 30 % and 5 % of a full scale fire respectively.

In practice, a full scale fire is defined as a fire where more than three fire hoses were needed for extinguishing the fire, a full scale fire is considered as a complete burnout. A large fire is in this context defined as a fire that involves the use of two or three fire hoses for fire extinguishing and is assumed to typically involve the majority of a house, an apartment, or at least part of an industrial complex. A medium size fire is in this context defined as a fire involving the use of only one fire hose for fire-fighting and will typically involve a part of a single room in an apartment or house. And a small size fire is in this context defined as a fire that was extinguished before the arrival of the fire service, extinguished by small tools or a chimney fire.

The total number of registered fires is known for the years 1989-2013. For the years 2007-2013 the total number of registered building fires is known with a very high degree of detail.

Table 6.17 shows the occurrence of all types of fires (registered for 1989-2012) and the occurrence of building fires (2007-2013) registered at DEMA. The 1980-1988 data for all fires are estimated to be the average of 1989-2010 data. In 2007-2010 the average per cent of building fires, in relation to all fires, was 60 %. The total numbers of building fires 1980-2006 are calculated using this percentage. The full time series is presented in Annex 3E-9.

Table 6.17 Occurrence of all fires and building fires.

	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
All fires	17751	17751	17025	19543	17174	16551	16728	16157	14084	14546
Building fires	10621	10621	10187	11694	10276	9903	9325	11447	9932	9893

The building fires that occurred in the years 2007-2012 are subcategorised into six building types; detached houses, undetached houses, apartment buildings, industrial buildings, additional buildings and container fires.

Table 6.18 states the average registered activity data for building fires for the years 2007-2010, divided in both damage size and building type. This describes the average share of building fires from 2007-2010 of a certain type and size, in relation to all building fires in the same four years period.

Table 6.18 Registered occurrence of building fires, average of 2007-2010 fires, %. (DEMA)

Size	Detached	Undetached	Apartment	Industry	Additional	Container	All building
Full	2.46	0.50	0.31	0.73	0.44	0.17	4.61
Large	4.01	1.14	1.09	1.69	3.08	1.92	12.93
Medium	5.24	2.33	6.15	2.92	4.30	18.46	39.40
Small	11.77	4.24	12.64	5.36	4.79	4.27	43.06
All	23.47	8.21	20.19	10.70	12.61	24.82	100.00

It is assumed that the average percentages provided by the years 2007-2010 shown in Table 6.18 are compliable for the years 1980-2006. Hereby, similar activity data for building fires can be estimated back to 1980.

By applying the damage rates of 100 %, 75 %, 30 % and 5 % corresponding to the damage sizes full, large, medium and small, a full scale equivalent can be determined. Table 6.19 shows the calculated full scale equivalents (FSE). The full time-series is presented in Annex 3E-10.

Table 6.19 Accidental building fires full scale equivalent activity data.

	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
Container fires	782	782	750	861	756	729	594	729	584	584
Detached house fires	810	810	777	892	784	755	833	818	742	761
Undetached house fires	240	240	231	265	233	224	194	206	181	162
Apartment building fires	383	383	367	421	370	357	348	362	327	316
Industry building fire	334	334	320	368	323	311	281	334	298	275
Additional building fires	455	455	437	501	440	424	429	740	610	619

#### Emission factors

For building fires, emissions are calculated by multiplying the number of full scale equivalent fires with the emission factors. The emission factors are produced from different measurements and assumptions from literature and expert judgements. When possible, emission factors are chosen that represent conditions that are comparable to Denmark. By comparable is meant countries that have similar building traditions, with respect to the materials used in building structure and interior.

In the process of selecting the best available emission factors for the calculation of the emissions from Danish accidental building fires, a range of different sources has been studied. Unfortunately it is difficult to do an interrelated comparison of the different sources because they all establish emission factors on different assumptions and many of these assumptions are not fully accounted for.

Table 6.20 lists the emission factors that are used for the year 2013 and their respective references.

Table 6.20 Emission factors building fires.

Compound	Unit /fire	Detached house	Undetached house	Apartment building	Industrial building	Additional building	Container	Reference
SO <sub>2</sub>	kg	266.6	214.4	124.5	802.9	32.1	2.4	Blomqvist et.al. 2002
NO <sub>x</sub>	kg	19.9	16.0	9.3	24.0	1.0	3.0	NAEI, 2009
NMVOC*	kg	99.6	80.1	46.5	120.0	4.8	0.7	NAEI, 2009
CO	kg	278.9	224.3	130.3	336.0	13.4	42.0	NAEI, 2009
TSP	kg	143.8	61.6	43.8	27.2	1.1	23.2	Aasestad, 2008**
PM <sub>10</sub>	kg	143.8	61.6	43.8	27.2	1.1	23.2	Aasestad, 2008**
PM <sub>2.5</sub>	kg	143.8	61.6	43.8	27.2	1.1	23.2	Aasestad, 2008**
As	g	1.35	0.58	0.41	0.25	0.01	0.22	Aasestad, 2008**
Cd	g	0.85	0.36	0.26	0.16	0.01	0.14	Aasestad, 2008**
Cr	g	1.29	0.55	0.39	0.24	0.01	0.21	Aasestad, 2008**
Cu	g	2.99	1.28	0.91	0.57	0.02	0.48	Aasestad, 2008**
Hg	g	0.85	0.36	0.26	0.16	0.01	0.14	Aasestad, 2008**
Pb	g	0.42	0.18	0.13	0.08	0.003	0.07	Aasestad, 2008**
PCDD/F*	mg	3.5	2.8	1.6	4.2	0.2	1.1	Hansen, 2000
Benzo[b]fluoranthene	g	12.6	10.1	5.9	15.2	0.6	1.9	NAEI, 2009
Benzo[k]fluoranthene	g	4.4	3.6	2.1	5.4	0.2	0.7	NAEI, 2009
Benzo[a]pyrene	g	8.0	6.4	3.7	9.6	0.4	1.2	NAEI, 2009
Indeno[1,2,3-cd]pyrene	g	8.6	6.9	4.0	10.4	0.4	1.3	NAEI, 2009

\*Container fires have a different source than the other five categories; Blomqvist et.al. 2002, \*\* Personal contact with Kristin Aasestad has provided a correction of the units which are inaccurate in the text of Aasestad (2008)

Emission factors for detached, undetached and apartment fires depend on the annual average floor space; see Table 6.21. Industrial, additional and container fires on the other hand are assumed to have a constant size/volume throughout the time series. Emission factors for all building fires for 1980-2013 are shown in Annex 3E-11.

Emission factors from Aasestad (2008) are already specified for four of the six building types; detached houses, undetached houses, apartment buildings and industrial buildings. Aasestad (2008) and all other sources considered were altered to match the six building types. This alternation was performed simply by adjusting the average floor space for each of the building types respectively, whereas factors like loss rate and mass of combustible contents per area are not altered.

The average floor space in Danish buildings is stated in Table 6.21. The data are collected from Statistics Denmark and takes into account possible multiple building floors but not attics and basements. For the full time series see Annex 3E-12. The average floor space in industrial buildings, schools etc. is estimated to 500 square meters for all years and the average floor space for additional buildings, sheds etc. is estimated to 20 square meters for all years.

Table 6.21 Average floor space in building types (Statistics Denmark, 2014).

	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
Detached houses	154	154	156	155	156	162	163	164	165	166
Undetached houses	130	130	129	129	131	131	134	132	134	133
Apartment buildings	74	75	75	75	75	76	77	78	78	78

Emission factors from literature are given in mass emission per mass burned. For the calculation of these emission factors to a unit that matches the activity data, the building masses are estimated using the same methodology as in Hansen (2000).



The total building masses are calculated using an average weight loss rate of 12.4 % (Persson et al., 1998) and data for the amount of combustible material in the building structure itself (Blomqvist et al., 2002) and the amount of combustible interior (Persson et al., 1998).

Emission factors for container fires cannot be calculated based on an average floor space but on an average mass. The average mass of a container is set to 1 Mg and covers all types of containers, from small residential garbage containers to large shipping containers and waste/goods in storage piles.

Building masses for 2013 are presented in Table 6.22.

Table 6.22 Building mass per building type.

	Unit	Detached house	Undetached house	Apartment building	Industry building	Additional building	Container
Average floor area*	m <sup>2</sup>	166	133	78	500	20	-
Building mass per floor area	kg/m <sup>2</sup>	40	40	35	30	30	-
Total building mass*	Mg/fire	6.6	5.3	2.7	15.0	0.6	1

\* 2013 numbers

For further detail on the emission factors and calculations, please refer to Hjelgaard (2013).

### Emissions

Table 6.23 shows the total emissions from building fires. The entire time series 1980-2013 is shown in Annex 3E-13.

Table 6.23 Emissions from building fires.

	unit	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
SO <sub>2</sub>	Mg	580.4	581.3	559.3	640.2	565.2	552.4	543.0	598.2	536.3	519.3
NO <sub>x</sub>	Mg	32.9	33.0	31.8	36.3	32.1	31.5	31.5	33.7	30.1	29.6
NMVOC	Mg	153.3	153.7	148.0	169.3	149.7	147.2	149.1	158.0	142.3	139.9
CO	Mg	460.7	461.6	444.5	508.4	449.5	441.2	441.3	471.6	421.9	415.1
TSP	Mg	175.8	175.8	168.6	193.5	170.1	163.9	168.8	173.0	154.5	155.0
PM <sub>10</sub>	Mg	175.8	175.8	168.6	193.5	170.1	163.9	168.8	173.0	154.5	155.0
PM <sub>2.5</sub>	Mg	175.8	175.8	168.6	193.5	170.1	163.9	168.8	173.0	154.5	155.0
As	kg	1.6	1.6	1.6	1.8	1.6	1.5	1.6	1.6	1.4	1.5
Cd	kg	1.0	1.0	1.0	1.1	1.0	1.0	1.0	1.0	0.9	0.9
Cr	kg	1.6	1.6	1.5	1.7	1.5	1.5	1.5	1.5	1.4	1.4
Cu	kg	3.7	3.7	3.5	4.0	3.5	3.4	3.5	3.6	3.2	3.2
Hg	kg	1.0	1.0	1.0	1.1	1.0	1.0	1.0	1.0	0.9	0.9
Pb	kg	0.5	0.5	0.5	0.6	0.5	0.5	0.5	0.5	0.5	0.5
PCDD/F	g I-TEQ	6.2	6.2	6.0	6.9	6.1	5.9	5.9	6.3	5.6	5.5
Benzo(b)fluoranthene	kg	20.8	20.9	20.1	23.0	20.3	20.0	20.0	21.3	19.1	18.8
Benzo(k)fluoranthene	kg	7.3	7.4	7.1	8.1	7.2	7.0	7.0	7.5	6.7	6.6
Benzo(a)pyrene	kg	13.2	13.2	12.7	14.5	12.8	12.6	12.6	13.5	12.1	11.9
Indeno(1,2,3-cd)pyrene	kg	14.3	14.3	13.8	15.7	13.9	13.7	13.7	14.6	13.1	12.8

### 6.5.3 Accidental vehicle fires

Pollutants that are emitted from accidental vehicle fires include SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, CO, particulate matter, heavy metals (As, Cd, Cr, Cu, Ni, Pb, Zn), PCDD/F and PAHs.

### Methodology

Emissions from vehicle fires are calculated by multiplying the mass of vehicle fires with selected emission factors. Emission factors are not available for different vehicle types, whereas it is assumed that all the different vehicle types leads to similar emissions. The activity data are calculated as an annual combusted mass by multiplying the number of different full scale vehicle fires with the Danish registered average weight of the given vehicle type.

### Activity data

As with accidental building fires, data for accidental vehicle fires are available through the Danish Emergency Management Agency (DEMA). DEMA provides very detailed data for 2007-2013; the remaining years back to 1980 are estimated by using surrogate data.

Table 6.24 shows the occurrence of fires in general and vehicle fires registered at DEMA. In 2007-2010 the average per cent of vehicle fires, in relation to all fires, was 20 %. The total numbers of vehicle fires in 1980-2006 are calculated using this percentage. The full time series is presented in Annex 3E-9.

Table 6.24 Occurrence of all fires and vehicle fires.

Year	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
All fires	17751	17751	17025	19543	17174	16551	16728	16157	14084	14546
Vehicle fires	3497	3497	3354	3850	3383	3260	3459	3255	2889	2841

There are fourteen different vehicle categories. The activity data are categorised in passenger cars (lighter than 3500 kg), buses, light duty vehicles (vans and motor homes), heavy duty vehicles (trucks and tankers), motorcycles/mopeds, other transport, caravans, trains, boats, airplanes, bicycles, tractors, combine harvesters and machines.

In the same manner as accidental building fires, the 2007-2013 data from DEMA can be divided in four categories according to damage size. It is assumed that a full scale fire is a complete burnout of the given vehicle, and that a large, medium and small scale fire corresponds to 75 %, 30 % and 5 % of a full scale fire respectively. The total number of full scale equivalent (FSE) fires can be calculated for each of the fourteen vehicle categories for 2007-2013.

The total number of registered vehicles is known from Jensen et al. (2012) and Statistics Denmark (2014). By assuming that the share of vehicle fires in relation to the total number of registered vehicles, of every category respectively, can be counted as constant, the number of vehicle fires is estimated for the years 1980-2006. The numbers of registered vehicles from 1980 to 1984 are extrapolated based on the years 1985 to 1989, where a clear trend has been visible this trend has been extrapolated (e.g. passenger cars), otherwise the average value of 1985 to 1989 has been used (e.g. buses).

Table 6.25 states the total number of national registered vehicles and the number of full scale equivalent vehicle fires. The full time series 1980-2013 is shown in Annex 3E-16.

Table 6.25 Number of nationally registered vehicles and full scale equivalent vehicle fires.

	Passenger Cars		Buses		Light Duty Vehicles		Heavy Duty Vehicles	
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires
1980	1475109	429	8070	12	99168	10	47428	60
1985	1564319	455	8010	11	147874	14	46962	60
1990	1645454	479	8109	12	192317	19	45664	58
1995	1733242	504	14371	21	228074	22	48077	61
2000	1916364	557	15051	22	272386	27	50227	64
2005	2012216	585	15131	22	372674	36	49311	63
2010	2246675	646	14577	23	362385	38	44813	60
2011	2281539	584	13915	13	343355	43	43640	54
2012	2326778	514	13177	11	318668	32	42326	53
2013	2373251	514	12829	11	306421	32	41999	53

*Continued*

	Motorcycles/Mopeds		Caravans		Train		Boat	
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires
1980	220273	78			7284	9	2222	25
1985	191478	68			7284	9	2222	25
1990	163133	58	86257	24	7156	9	2324	26
1995	165272	58	95831	26	6854	8	1911	21
2000	233309	82	106935	29	4907	6	1759	19
2005	273904	97	121350	33	3195	4	1792	20
2010	301562	83	142354	37	2740	2	1773	16
2011	295488	91	142764	34	2943	3	1768	21
2012	295798	82	142654	33	3055	2	1772	14
2013	296522	82	142667	33	3048	2	1781	14

*Continued*

	Airplane		Tractor		Combined Harvester		Bicycle	Other Transport	Machine
	Registered	FSE fires	Registered	FSE fires	Registered	FSE fires	FSE fires	FSE fires	FSE fires
1980	1060	1	139600	87	38781	64			
1985	1060	1	128700	80	35708	59			
1990	1055	1	131880	82	33594	56			
1995	1058	1	130028	81	27986	46			
2000	1070	1	111736	69	23272	39			
2005	1073	1	104551	65	20965	35			
2010	1152	1	89141	77	15986	32	4	58	94
2011	1132	0	85776	59	14990	21	3	50	111
2012	1111	0	82410	68	13994	18	2	50	115
2013	1069	0	79045	68	12998	18	-	-	-

The average weights of a passenger car, bus, light- and heavy commercial vehicle and motorcycle/moped are known for every year back to 1993 (Statistics Denmark, 2013), the weight of combined harvesters is based on an expert judgement. The corresponding weights from 1980 to 1992 and the average weight of the units from the remaining categories are estimated by an expert judgment; see Table 6.26 and Annex 3E-15.

Table 6.26 Average weight of different vehicle categories, kg.

	Cars	Buses	Light Duty Vehicles	Heavy Duty Vehicles	Motorcycles/ Mopeds	Combined Harvester
1980	850	10000	2000	15000	75	8000
1985	850	10000	2000	15000	75	8750
1990	850	10000	2000	15000	86	9500
1995	923	10807	2492	14801	97	10250
2000	999	11195	3103	15214	103	11000
2005	1068	11560	3793	13258	116	11750
2010	1144	11804	4498	11883	133	12500
2011	1154	11907	4296	11291	135	12650
2012	1160	11625	4150	10844	136	12800
2013	1162	11463	4046	10861	137	12950

It is assumed that the average weight of a boat equals that of a bus. That tractors and vans weigh the same and that trains and airplanes have the same average weight as trucks.

Bicycles, machines and other transport can only be calculated for the years 2007-2013 due to the lack of surrogate data (number of nationally registered vehicles). The average weight of a bicycle, caravan, machine and other transport is estimated as 12 kg, 90 % of a car, 50 % of a car and 40 % of a car respectively.

By multiplying the number of full scale fires with the average weight of the vehicles respectively, the total amount of combusted vehicle mass can be calculated. The result is shown in Table 6.27 and in Annex 3E-14.

Table 6.27 Burnt mass of different vehicle categories, Mg.

	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
Passenger cars	365	387	407	466	557	625	739	674	592	555
Buses	116	115	116	223	242	251	266	160	130	121
Light duty vehicles	19	29	37	55	82	138	171	185	133	118
Heavy duty vehicles	902	893	869	902	969	829	715	606	579	455
Motorcycle, moped	6	5	5	6	8	11	10	12	11	11
Other transport	-	-	-	-	-	-	33	29	29	26
Caravan	-	-	30	36	44	53	63	59	57	59
Train	130	130	128	121	89	51	24	28	23	18
Boat	246	246	257	228	218	229	189	249	160	100
Airplane	12	12	12	11	12	10	7	3	5	5
Bicycle	-	-	-	-	-	-	0	0	0	0
Tractor	174	160	164	202	216	247	347	254	283	330
Combine harvester	515	518	530	476	425	409	398	271	236	402
Machine	-	-	-	-	-	-	43	51	53	53
Total	2484	2495	2555	2727	2863	2858	3006	2580	2291	2253

### Emission factors

In the process of selecting the most reliable emission factors for the calculation of the emissions from Danish vehicle fires, a range of different sources have been studied. Unfortunately it is difficult to make an interrelated comparison of the different sources because they all establish emission factors on different assumptions and many of these assumptions are not fully accounted for. Table 6.28 lists the accepted emission factors and their respective references.

Table 6.28 Emission factors vehicle fires.

	Unit, per Mg	Emission factor	Source
SO <sub>2</sub>	kg	5	Lönnermark et al., 2004
NO <sub>x</sub>	kg	2	Lemieux et al., 2004
NMVOC	kg	8.5	Lönnermark et al., 2004
CO	kg	63	Lönnermark et al., 2004
TSP	kg	38	Lönnermark et al., 2004
PM <sub>10</sub>	kg	38	Lönnermark et al., 2004
PM <sub>2.5</sub>	kg	38	Lönnermark et al., 2004
As	g	0.26	Lönnermark et al., 2004
Cd	g	1.70	Lönnermark et al., 2004
Cr	g	3.80	Lönnermark et al., 2004
Cu	g	27.0	Lönnermark et al., 2004
Ni	g	2.80	Lönnermark et al., 2004
Pb	g	820	Lönnermark et al., 2004
Zn	g	3200	Lönnermark et al., 2004
PCDD/F	mg	0.04	Hansen, 2000
Benzo(b)fluoranthene	g	32.3	Lemieux et al., 2004
Benzo(k)fluoranthene	g		Lemieux et al., 2004
Benzo(a)pyrene	g	14.7	Lemieux et al., 2004
Indeno(1,2,3-cd)pyrene	g	23.3	Lemieux et al., 2004

No data are available for Hg, Se, HCB and PCBs. NH<sub>3</sub> is assumed not to be emitted.

### Emissions

Table 6.29 shows the total national emissions from vehicle. The entire time series is shown in Annex 3E-17.

Table 6.29 National emissions from vehicle fires.

	unit	1980	1985	1990	1995	2000	2005	2010	2011	2012	2013
SO <sub>2</sub>	Mg	12.42	12.47	12.77	13.64	14.32	14.29	15.12	13.12	11.59	11.27
NO <sub>x</sub>	Mg	4.97	4.99	5.11	5.45	5.73	5.72	6.05	5.25	4.64	4.51
NMVOC	Mg	21.11	21.21	21.72	23.18	24.34	24.29	25.71	22.30	19.71	19.15
CO	Mg	156.48	157.18	160.95	171.80	180.40	180.04	190.57	165.31	146.07	11.27
TSP	Mg	94.39	94.81	97.08	103.63	108.81	108.60	114.95	99.71	88.11	141.95
PM <sub>10</sub>	Mg	94.39	94.81	97.08	103.63	108.81	108.60	114.95	99.71	88.11	5.41
PM <sub>2.5</sub>	Mg	94.39	94.81	97.08	103.63	108.81	108.60	114.95	99.71	88.11	85.62
As	kg	0.65	0.65	0.66	0.71	0.74	0.74	0.79	0.68	0.60	85.62
Cd	kg	4.22	4.24	4.34	4.64	4.87	4.86	5.14	4.46	3.94	85.62
Cr	kg	9.44	9.48	9.71	10.36	10.88	10.86	11.49	9.97	8.81	0.59
Cu	kg	67.06	67.36	68.98	73.63	77.31	77.16	81.67	70.85	62.60	3.83
Ni	kg	6.95	6.99	7.15	7.64	8.02	8.00	8.47	7.35	6.49	8.56
Pb	Mg	2.04	2.05	2.09	2.24	2.35	2.34	2.48	2.15	1.90	60.84
Zn	Mg	7.95	7.98	8.18	8.73	9.16	9.14	9.68	8.40	7.42	6.31
PCDD/F	g I-TEQ	0.10	0.10	0.10	0.11	0.11	0.11	0.12	0.10	0.09	1.85
Benzo(b)fluoranthene	kg	40.11	40.29	41.26	44.04	46.25	46.15	48.85	42.38	37.44	7.21
Benzo(k)fluoranthene	kg	40.11	40.29	41.26	44.04	46.25	46.15	48.85	42.38	37.44	0.09
Benzo(a)pyrene	kg	36.51	36.68	37.56	40.09	42.09	42.01	44.47	38.57	34.08	36.39
Indeno(1.2.3-cd)pyrene	kg	57.87	58.13	59.53	63.54	66.72	66.59	70.48	61.14	54.02	36.39

### 6.5.4 Other

Other combustion sources include open burning of yard waste and bonfires.

Due to the cold and wet climatic conditions in Denmark wild fires very seldom occur. Controlled field burnings and the occasional wild fires are categorised under the sectors Agriculture and Land Use, Land Use Change and Forestry (LULUCF), respectively.

In Denmark, the open burning of private yard waste is under different restrictions according to the respective municipality. These restrictions involve what can be burned but also the quantity, and how, when and where, or in some cases a complete ban is imposed. The burning of yard waste is not allowed within urban areas (DEPA, 2011). There is no registration of private waste burning and the activity data on this subject are very difficult to estimate. Citizens are generally encouraged to compost their yard waste or to dispose of it through one of the many waste disposal/recycling sites.

The occurrence of bonfires at midsummer night and in general are likewise not registered. Therefore it has not been possible to obtain activity data and consequently, Bonfires are not included in this inventory.

## 6.6 Uncertainties and time series consistency

This section covers the uncertainty estimates

### 6.6.1 Input data

The uncertainty of the number of human cremations is miniscule. However, for the purpose of the calculation it has been set to 1 %.

The uncertainty of the activity data from animal cremations is also minimal for the most recent years (1998-2013) but is increasing back in time (to 200 % in 1980). The uncertainty is set to 40 % for all years.

Activity data for composting are estimated for the years 1990-1994 and 2010-2013 resulting in a higher level of uncertainty these years; this is set at 40 %.

The uncertainty of the total number of accidental fires is very small but the division into building and transportation types and also the calculation of full scale equivalents will lead to some uncertainty - partly caused by the category "other". The uncertainty for both building and vehicle activity data is therefore set to 10 % for all years. The uncertainty is however lowest for the most recent years (2007-2013).

Activity data for combustion of biogas at biogas production facilities are available from the national energy statistics; the uncertainty for this activity is set to 5 %.

The following Tables 6.30 lists the uncertainties for activity data in the waste sector.

Table 6.30 Estimated uncertainty rates for activity data.

	Human cremation	Animal cremation	Composting	Building fires	Vehicle fires	Biogas Production*
Activity data uncertainty, %	1	80	50	10	10	5*

\*This category only exists from 1994-2004 and is therefore (with the exception of particles) not included in the uncertainty calculations included in this report. Uncertainties for all pollutants are calculated for 1990 and 2013, except for those for particles which are calculated for 2000 and 2013.

The uncertainties for emission factors in the waste sector and at the present level of available information are listed in Table 6.31. The uncertainties are assumed valid for all years 1990-2013.

Table 6.31 Estimated uncertainty rates for emission factors. %.

Pollutant	Human cremation	Animal cremation	Compos-ting	Building fires	Vehicle fires	Biogas production
SO <sub>2</sub>	100	100		300	500	100
NO <sub>x</sub>	150	150		500	500	100
NM VOC	100	300		500	500	100
CO	150	150	100	500	500	100
NH <sub>3</sub>		300	100			
TSP	500	300		500	700	500
PM <sub>10</sub>	500	300		500	700	500
PM <sub>2.5</sub>	500	300		500	700	500
As	700	700		500	500	400
Cd	700	500		500	500	150
Cr	700	500		500	500	100
Cu	700	500		500	500	100
Hg	150			500		100
Ni	700	500			500	100
Pb	600	500		500	500	100
Se	700	700				100
Zn	700	500			500	150
HCB	500	500				1000
PCDD/F	300	300		100	100	1000
Benzo(b)flouranthene	1000	1000		500	500	
Benzo(k)flouranthene	1000	1000		500	500	
Benzo(a)pyrene	1000	1000		500	500	
Indeno(1.2.3-c.d)pyrene	1000	1000		500	500	
PCB	1000	1 000				1000

### 6.6.2 Uncertainty results

The Tier 1 uncertainty estimates for the waste sector are calculated from 95 % confidence interval uncertainties. Results are shown in Table 6.32.

Table 6.32 Tier 1 uncertainty results for waste.

Pollutant	Emission 2013. Mg	Total emission uncertainty. %	Trend 1990-2013. %	Trend Uncertainty. %-age points
SO <sub>2</sub>	537	±290.2	-7.0	±12.8
NO <sub>x</sub>	85	±189.3	16.7	±61.5
NM VOC	162	±435.6	-5.2	±13.2
CO	1156	±197.5	50.2	±187.5
NH <sub>3</sub>	783	±107.4	278.8	±213.7
TSP	243	±402.7	-13.5*	±26.8
PM <sub>10</sub>	243	±404.0	-13.7*	±26.7
PM <sub>2.5</sub>	242	±404.5	-13.7*	±26.7
As	2.30E-03	±349.2	-18.7	±128.5
Cd	4.76E-03	±413.8	-14.2	±26.5
Cr	1.00E-02	±432.0	-14.8	±32.5
Cu	6.41E-02	±475.4	-12.2	±12.7
Hg	1.39E-03	±333.5	-97.0	±9.9
Ni	6.39E-03	±493.8	-18.8	±60.9
Pb	1.85E+00	±499.9	-11.8	±12.5
Se	3.84E-04	±686.0	-55.1	±409.7
Zn	7.21E+00	±500.1	-11.9	±12.5
HCB	9.30E-06	±378.4	41.8	±257.3
PCDD/F	5.61E-06	±98.7	-8.0	±12.8
Benzo(b)flouranthene	2.76E-02	±500.1	-6.8	±13.2
Benzo(k)flouranthene	5.51E-02	±371.3	-10.2	±12.2
Benzo(a)pyrene	4.30E-02	±430.2	-11.1	±11.7
Indeno(1.2.3-c.d)pyrene	4.50E-02	±391.3	-10.5	±11.9
PCB	1.28E-02	±500.1	-6.8	±13.2

\*Trend 2000-2013. %

## 6.7 QA/QC and verification

A list of QA/QC tasks are performed directly in relation to the emissions from the waste sector part of the Danish emission inventories. The following procedures are carried out to ensure the data quality:

- Checking of time series in the NFR and SNAP source categories. Considerable changes are controlled and explained.
- Comparison with the inventory of the previous year. Any major changes are verified.
- A manual log table is applied to collect information about recalculations.
- Some automated checks have been prepared for the emission databases:
- Check of units for fuel rate and emission factors
- Additional checks on database consistency

The QC work will continue in future years.

### 6.7.1 Data deliveries

Table 6.33 lists the external data deliveries used for the waste emission inventory. Further the table holds information on the contacts at the data delivery companies.



Table 6.33 List of external data sources.

Category	Data description	Activity data. emission factors or emissions	Reference	Contact(s)	Data agreement/ Comment	http. file or folder name
Human cremation	Annual number of cremated persons	Activity data	Association of Danish Crematories	Hanne Ring	Public access	<a href="http://www.dkl.dk">http://www.dkl.dk</a>
Human cremation	Population statistics	Activity data	Statistics Denmark		Public access	<a href="http://www.statistikbanken.dk/BEF5">http://www.statistikbanken.dk/BEF5</a>
Animal cremation	Annual number of cremated carcasses	Activity data	Dansk Dyre-kremering ApS	Knud Ribergaard	Personal contact	
Animal cremation	Annual number of cremated carcasses	Activity data	Ada's Kæle- dyrskrematorium ApS	Frederik Møller	Personal contact	
Animal cremation	Annual number of cremated carcasses	Activity data	Kæledyrs- krematoriet	Annette Laursen	Personal contact	
Accidental building fires	Average floor space in buildings	Activity data	Statistics Denmark		Public access	<a href="http://www.statistikbanken.dk/BOL511">http://www.statistikbanken.dk/BOL511</a>
Accidental fires	Categorised fires	Activity data	The Danish Emergency Management Agency	Steen Hjere Nonnemann	Public access	<a href="https://statistikbanken.brs.dk">https://statistikbanken.brs.dk</a>
Accidental building fires	Building type statistics	Activity data	Statistics Denmark		Public access	<a href="http://www.statistikbanken.dk/BOL11.BOL3.BOL33.and.BYGB11">http://www.statistikbanken.dk/BOL11. BOL3. BOL33 and BYGB11</a>
Accidental vehicle fires	Weight categorisation of vehicles (passenger cars. busses. vans and trucks)	Activity data	Statistics Denmark		Public access	<a href="http://www.statistikbanken.dk/BIL10.BIL12.BIL15.and.BIL18">http://www.statistikbanken.dk/BIL10. BIL12. BIL15 and BIL18</a>
Composting	Waste categories for composting	Activity data	Waste Statistics (Affaldsstatistik)		Public access	<a href="http://www2.mst.dk/udgiv/publikationer/2010/978-87-92668-21-9/pdf/978-87-92668-22-6.pdf">http://www2.mst.dk/udgiv/publikationer/2010/978-87-92668-21-9/pdf/978-87-92668-22-6.pdf</a>

## 6.8 Source-specific recalculations and improvements

A new reporting template has been used for the 2015 submission in accordance with the reporting guidelines. The main change in the waste sector is that emissions from composting are now reported in a separate category rather than under 'Other waste' as in previous submissions.

Minor changes were made to the activity data for composting and accidental fires to incorporate new information and correct errors in the previous reporting.

## 6.9 Source-specific planned improvements

There are currently no planned improvements for this sector.

## 6.10 References

Aasestad, K. (eds.). 2008: Norwegian Emission Inventory 2008. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. Report 2008/48. Statistics Norway. Available at:  
[http://www.ssb.no/english/subjects/01/04/10/rapp\\_emissions\\_en/rapp\\_200848\\_en/rapp\\_200848\\_en.pdf](http://www.ssb.no/english/subjects/01/04/10/rapp_emissions_en/rapp_200848_en/rapp_200848_en.pdf) (30-12-2014).

Affaldsstatistik, 2006: (Waste Statistics). Orientering fra Miljøstyrelsen Nr. 2 2008. Miljøstyrelsen. Miljøministeriet. Available at:  
<http://www2.mst.dk/udgiv/publikationer/2008/978-87-7052-753-8/pdf/978-87-7052-754-5.pdf> (In Danish) (23-01-2015).

Affaldsstatistik, 2007 og 2008 (Waste Statistics). Orientering fra Miljøstyrelsen Nr. 5 2010. Miljøstyrelsen. Miljøministeriet. Available at:  
<http://www2.mst.dk/udgiv/publikationer/2010/978-87-92668-21-9/pdf/978-87-92668-22-6.pdf> (In Danish) (09-01-2014).

Amlinger, F., Peyr, S. & Cuhls, C. 2008: Greenhouse Gas Emissions from Composting and Mechanical Biological Treatment. Waste Management & Research 2008; 26; 47. Sage publications. International Solid Waste Association. Available at:  
<http://wmr.sagepub.com/cgi/reprint/26/1/47.pdf> (30-12-2014).

Blomqvist, P., Persson, B. & Simonson, M. 2002: Utsläpp från bränder till miljön. Utsläpp av dioxin. PAH och VOC till luften. Räddningsverket. Karlstad. FoU rapport. SP Brandteknik. Borås (In Swedish).

Boldrin, A., Andersen, J.K. & Christensen, T.H. 2009: Miljøvurdering af haveaffald i Århus kommune. Danmarks Tekniske Universitet - Miljø. 3R.

Chen, S.-J., Hung, M.-C., Huang, K.-L. & Hwang, W.-I. 2004: Emission of heavy metals from animal carcass incinerators in Taiwan. Chemosphere 55 (2004) 1197-1205. Elsevier.

Claire, S. 1999: Source inventory. Category #750. Miscellaneous emission sources accidental fires – Structural 1999 Emissions. Base year: 1999. pp. 10.3.1 – 2. Available at:  
[www.arb.ca.gov/ei/areasrc/districtmeth/BayArea/C750.pdf](http://www.arb.ca.gov/ei/areasrc/districtmeth/BayArea/C750.pdf) (30-12-2014).

Danish Energy Agency (DEA), 2014: The Danish Annual Energy Statistics. Basicdata 2013.xls. Available at:  
[http://www.ens.dk/en-US/Info/FactsAndFigures/Energy\\_statistics\\_and\\_indicators/Annual%20Statistics/Sider/Forside.aspx](http://www.ens.dk/en-US/Info/FactsAndFigures/Energy_statistics_and_indicators/Annual%20Statistics/Sider/Forside.aspx) (30-12-2014).

DEMA: The Danish Emergency Management Agency (DEMA). Beredskabsstyrelsen. Statistikbanken. Available at:  
<https://statistikbank.brs.dk/sb/main/p/a0109> (In Danish) (30-12-2014).

DEMA, 2007: The Danish Emergency Management Agency (DEMA). Beredskabsstyrelsen. Redningsberedskabets Statistiske Beretning 2007. Available at:  
[http://brs.dk/viden/publikationer/Documents/RBS\\_Statistisk\\_Beretning\\_2007.pdf](http://brs.dk/viden/publikationer/Documents/RBS_Statistisk_Beretning_2007.pdf) (In Danish) (30-12-2014).

Danish Environmental Protection Agency (DEPA), 2006: Legislation on sludge. Bekendtgørelse om anvendelse af affald til jordbrugsformål (Slambekendtgørelsen). BEK nr. 1650 af 13/12/2006. Available at: <https://www.retsinformation.dk/Forms/R0710.aspx?id=13056> (30-12-2014) (Danish).

Danish Environmental Protection Agency (DEPA), 2011: Legislation on waste (Affaldsbekendtgørelsen). BEK nr. 1415 af 12/12/2011 Gældende. Chapter 7. §46 and §47. Available at: <https://www.retsinformation.dk/forms/r0710.aspx?id=139654#Kap7> (30-12-2014) (Danish).

DKL, 2009: Danske Krematoriers Landsforening. Røggasrensning, miljøbehandling og varmeudnyttelse. Status juni 2009. Available at: <http://www.ft.dk/samling/20081/almdel/mpu/spm/591/svar/641398/714098/index.htm>

DKL, 2014: Danske krematoriers landsforening. Statistisk materiale. Available at: <http://dkl.dk/Info/statistik.php> (In Danish) (30-12-2014).

European Commission (EC), 2011: Regulations. Commission Regulation (EU) No 142/2011. of 25 February 2011. Available at: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2011:054:0001:0254:EN:PDF> (30-12-2014)

EEA, 1996: European Environmental Agency. Emission Inventory Guidebook 2007. SNAP 090901. December 2006.

EEA, 2009: EMEP/EEA air pollutant emission inventory guidebook — 2009. Technical report No 6/2009. Available at: <http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009> (30-12-2014).

EIIP, 2001: Structure Fires. Volume III: Chapter 18. Emission Inventory Improvement Program. Revised Final January 2001. Eastern Research Group. Inc.. Area Source Committee. Available at: [http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii18\\_apr2001.pdf](http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii18_apr2001.pdf) (30-12-2014).

Hansen, E. 2000: Substance Flow Analysis for dioxins in Denmark. Environmental Project No. 570. Miljøprojekt. the Danish Environmental Protection Agency. Available at: <http://www2.mst.dk/udgiv/publications/2000/87-7944-295-1/pdf/87-7944-297-8.pdf> (In Danish) (30-12-2014).

Hellebrand, H.J. 1998: Emission of Nitrous Oxide and other Trace Gases during Composting of Grass and Green Waste. Institute of Agricultural Engineering Bornim (ATB). Germany. Silsoe Research Institute. Journal of Agricultural Engineering Research (1998) 69. 365-375.

Henriksen, T.C., Illerup, J.B. & Nielsen, O.-K. 2006: Dioxin Air Emission Inventory 1990-2004. National Environmental Research Institute. Denmark. 90 pp. – NERI Technical report no 602. Available at: <http://www.dmu.dk/Pub/FR602.pdf>.

IPCC, 2006: Guidelines for National Greenhouse Gas Inventories. Chapter 4: Biological Treatment of Solid Waste. Available at: [http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/5\\_Volume5/V5\\_4\\_Ch4\\_Bio\\_Treat.pdf](http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/5_Volume5/V5_4_Ch4_Bio_Treat.pdf).

Hjelgaard, K. 2013: Danish Emission Inventory for Waste Incineration and Other Waste. Inventories until year 2011. Aarhus University. DCE – Danish Centre for Environment and Energy. 96 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 70. Available at: <http://www.dmu.dk/Pub/SR70.pdf>

Jensen, T.C. & Kveiborg, O. 2013: Dokumentation af konvertering af trafiktal til emissionsopgørelser. arbejdsnotat. DTU Transport. 2013.

Kriegbaum, M. & Jensen, E. 2005: Input til branchebilag for krematorie-anlæg. Kommentarer og begrundelser. Dato 23. september 2005. Available at: <http://mst.dk/media/mst/68544/060116Inputkrematorieanl%C3%A6g.pdf> (In Danish) (30-12-2014).

Lemieux, P.M., Lutes, C.C. & Santoianni, D.A. 2004: Emissions of organic air toxics from open burning: a comprehensive review. Elsevier. science direct; Progress in Energy and Combustion Science 30. Available at: <http://www.fire.uni-freiburg.de/vfe/Emissions-Open-Burning-Lemieux-et-al-2004.pdf> (09/01-2014).

Lönnermark, A. & Blomqvist, P. 2004: Emissions from an automobile fire. Elsevier. chemosphere 62 (2006); Dec 2004; p1043-1056.

Ministeriet for Ligestilling og Kirke (MILIKI) (Ministry for Gender Equality and Ecclesiastical Affairs), 2006: Arbejdsgruppe foreslår at der skal være færre krematorier. december 2006. Available at: <http://www.km.dk/kirke/nyhedsarkiv-kirke/nyhed-om-kirke/artikel/arbejdsgruppe-foreslaar-at-der-skal-vaere-faerre-krematorier/> (In Danish) (30-12-2014).

MST, 2013: Livscyklusvurdering og samfundsøkonomisk analyse for anvendelse af spildevandsslam. Miljøministeriet. Miljøstyrelsen (MST). Miljøprojekt nr. 1459. 2013. ISBN nr. 978-87-92903-81-5. Available at: <http://www.mst.dk/Publikationer/Publikationer/2013/Maj/978-87-92903-81-5.htm> (Danish) (30-12-2014).

NAEI, 2009: The UK National Atmospheric Emission Inventory. Available at: <http://www.naei.org.uk/emissions/selection.php>

Pagans, E., Barrena, R., Font, X. & Sánchez, A. 2006: Ammonia emissions from the composting of different organic wastes. Dependency on process temperature. Chemosphere 62. 1534-1542. Elsevier.

Persson, B. & Simonson, M. 1998: Fire Emissions Into the Atmosphere. Swedish National Testing and Research Institute. Birås. Sweden. Fire Technology. Vol. 34. No. 3. 1998.

Petersen, C. 2001: Statistik for behandling af organisk affald fra husholdninger – Revideret udgave. 1999 (Statistics for treatment of organic waste from households – revised version). ECONET A/S. Miljøprojekt Nr. 654. Miljøstyrelsen. Miljøministeriet. Available at: <http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-932-8/pdf/87-7944-933-6.pdf> (In Danish) (30-12-2014).

Petersen, C. & Hansen, V.L. 2003: Statistik for behandling af organisk affald fra husholdninger 2001 (Statistics for treatment of organic waste from households). ECONET A/S. Miljøprojekt Nr. 856. Miljøstyrelsen. Miljøministeriet. Available at:

<http://www2.mst.dk/Udgiv/publikationer/2003/87-7972-962-2/pdf/87-7972-963-0.pdf> (In Danish) (30-12-2014).

Petersen, C. & Kielland, M. 2003: Statistik for hjemmekompostering 2001 (Statistics on home composting). Econet A/S. Miljøprojekt Nr. 855. Miljøstyrelsen. Miljøministeriet. Available at:

<http://www2.mst.dk/udgiv/publikationer/2003/87-7972-960-6/pdf/87-7972-961-4.pdf> (In Danish) (30-12-2014).

Santarsiero, A., Settimo, G., Cappiello, G., Viviano, G., Dell'Andrea, E. & Gentilini, L. 2005: Urban crematoria pollution related to the management of the deceased. Microchemical Journal. Vol. 79. Issues 1-2. Pp 299-306.

Schleicher, O. & Gram, L.K. 2008: Miljøprojekt Nr. 1191. Analyse af omkostningerne for rensning for kviksølv på krematorier i Danmark. Miljøministeriet. Miljøstyrelsen. Available at:

<http://www2.mst.dk/common/Udgivramme/Frame.asp?http://www2.mst.dk/udgiv/publikationer/2008/978-87-7052-594-7/html/helepubl.htm> (In Danish) (30-12-2013).

Schleicher, O. & Jensen, A.A. 2004: Håndbog om vurdering af spredning af dioxin og andre miljøskadelige stoffer fra ukontrollerede brande. Miljøprojekt nr. 918. FORCE Technology. Miljøstyrelsen. Miljøministeriet. Available at:

<http://www.mst.dk/Publikationer/Publikationer/2004/05/87-7614-238-8.htm> (In Danish) (30-12-2014).

Schleicher, O., Jensen, A.A. & Blinksbjerg, P. 2001: Miljøprojekt Nr. 649. Måling af dioxinmissionen fra udvalgte sekundære kilder. Miljøministeriet. Miljøstyrelsen. Available at:

<http://www2.mst.dk/Udgiv/publikationer/2001/87-7944-868-2/pdf/87-7944-869-0.pdf> (In Danish) (30-12-2014).

Statistics Denmark, 2013: StatBank Denmark 2013. Available at:

<http://www.statistikbanken.dk/statbank5a/default.asp?w=1024> (In Danish and English) (30-12-2014).

Toda, E. 2006: POPs and heavy metals emission inventory of Japan. Ministry of the Environment. Japan.

UNEP, 2005: UNEP toolkit. United Nations Environment Programme. Standardized Toolkit for Identification and Quantification of Dioxin and Furan Releases. Edition 2.1 December 2005. Prepared by UNEP Chemicals. Geneva. Switzerland. Available at:

<http://chm.pops.int/Portals/0/Repository/toolkit1/UNEP-POPS-TOOLKIT.1-4.English.PDF> (30-12-2014).

WEBFIRE, 1992: United States Environmental Protection Agency database. Available at:

<http://cfpub.epa.gov/oarweb/index.cfm?action=fire.simpleSearch> (30-12-2014).

## **7 Other and natural emissions**

Denmark does not report emissions in the NFR category “Other” (NFR 7). Regarding natural emissions volcanoes do not occur in Denmark and hence the category is reported as NO (Not Occurring).

Emissions from forest fires are for most years negligible but have not been estimated. Any other natural emissions, to be reported under NFR category 11C, have also not been estimated.

## 8 Reporting spatially distributed emissions on grid

This chapter include descriptions on input data, methodology and results of the Danish gridded emissions for the years 2005 and 2010. A detailed methodological description is given in Plejdrup & Gyldenkærne (2011).

### 8.1 Background for reporting

According to the UNECE Convention on Long-Range Transboundary Air Pollution parties are obligated to report gridded emissions.

In December 2013 the Executive Body for the Convention on Long-range Transboundary Air Pollution adopted new reporting guidelines, which include requirement of four-year reporting of gridded emissions from 2017. The new reporting guidelines are yet not implemented for gridded emissions, why the data and methodology presented in this chapter refer to the latest reporting of gridded emissions according to the previous guidelines.

In the 2012 reporting Denmark reported gridded emissions for the years 2005 and 2010. The mandatory reporting of gridded emissions includes the following 13 pollutants: SO<sub>x</sub>, NO<sub>x</sub>, NH<sub>3</sub>, NMVOC, CO, PM<sub>10</sub>, PM<sub>2.5</sub>, Pb, Cd, Hg, benzo(b)flouranthene, benzo(k)flouranthene, benzo(a)pyrene, indeno(1,2,3-c,d)pyrene, HCB, and dioxins and furans. The reporting includes GNFR sectoral emissions as well as national total emissions disaggregated to the standard EMEP grid with a resolution of 50 km x 50 km. Table 9.1 lists the categories (sectors) used for reporting gridded emission data based on the Danish inventories.

Table 9.1 GNFR categories and corresponding NFR categories and SNAP IDs in the Danish gridded emission inventory

GNFR ID	GNFR (long name)	NFR	SNAP	Note
A_PublicPower	PublicPower	1A1a	0101, 0102	
B_IndustrialComb	IndustrialCombustion	1A1c, 1A2a, 1A2b, 1A2c, 1A2d, 1A2e, 1A2f i	0103, 0105, 0301, 0302, 0304, 0305, 0306, 0307, 0308, 0309, 0310 , 0311, 0312, 0313, 0314, 0315, 0316, 0320	
C_SmallComb	SmallCombustion	1A4a i, 1A4b i, 1A4c i	0201, 0202, 0203	
D_IndProcess	IndustrialProcesses	2	0303, 0402, 0404, 0405, 0406	
E_Fugitive	Fugitive emissions from fuels	1B1, 1B2	0401, 0501, 0502, 0505, 0506, 0902	
F_Solvents	Solvent and other product use	3		06
G_RoadRail	RoadRailway	1A3b, 1A3c		07, 0802
H_Shipping	Shipping	1A3d ii, 1A4c iii		0803, 080402, 080403
I_OffRoadMob	OffRoadMobile	1A2f ii, 1A4a ii, 1A4b ii, 1A4c ii, 1A5b	0801, 0806, 0807, 0808, 0809, 0811	
J_AviationLTO	AviationLTO	1A3 a i (i), 1A3 a ii (i)		080501, 080502
L_OtherWasteDisp	OtherWasteDisposal	6D		0910
M_WasteWater	WasteWater			NE
N_WasteIncin	WasteIncineration	6C		0909
O_AgriLivestock	AgricultureLivestock	4B		*
P_AgriOther	AgricultureOther	4D, 4G		*
Q_AgriWaste	AgricultureWaste	4F		*
R_Other	Other			NO
S_Natural	Natural			NO
K_CivilAviCruise	CivilAviationCruise	1A3a ii (ii)		080503
T_IntAviCruise	IntAviationCruise	1A3a i (ii)		080504
Z_memo	memo	1A3d i (i)		080404

\* The Danish national emission inventory system for agriculture builds on NFR categories and not SNAP categories as is the case for the remaining sectors in the Danish emission inventory system

The Guidelines used for this reporting are included in UNECE (2009). The methodology in Danish emission gridding model SPREAD follows the EMEP/EEA Guidebook (2009)<sup>1</sup>. The gridded emission data in the 2012 reporting are available at the EIONET Central Data Repository homepage:

[http://cdr.eionet.europa.eu/dk/Air\\_Emission\\_Inventories/Submission\\_EMEP\\_UNECE](http://cdr.eionet.europa.eu/dk/Air_Emission_Inventories/Submission_EMEP_UNECE)

Further, a detailed methodological description is given in Plejdrup & Gyldenkærne (2011).

## 8.2 Methods and data for disaggregation of emission data

A national model for high resolution spatial distribution of emissions to air, the SPREAD model, has been developed at Department of Environmental Science, Aarhus University. SPREAD includes all sources and pollutants in

<sup>1</sup> The SPREAD version used for the 2012 reporting follow the 2009 EMEP/EEA Guidebook. From the 2017 reporting, the gridded emissions will be prepared according to the 2013 EMEP/EEA Guidebook.



the Danish emission inventory system, and generates emissions on a resolution of 1 km x 1 km.

SPREAD covers the area defined by the Exclusive Economic Zone (EEZ) and the national boarder. Denmark is geographically the peninsula of Jutland and 443 named islands and islets, of which approximately 72 are inhabited. The country is located in Scandinavia neighbouring the sea (the Baltic Sea, Skagerrak, Kattegat and the North Sea) as well as Germany, which Jutland are adjacent to the south (Figure 9.1).

The spatial emission distribution is carried out on the most disaggregated level possible and therefore SPREAD includes a large number of distribution keys related to single sources, sub categories and in a single case to a whole sector. Gridded emissions reported to UNECE LRTAP are based on the results from SPREAD, aggregated on the 50 km x 50 km EMEP grid.

The spatial distribution in SPREAD is based on a number of national geographical data sets. As the model is very complex and include many spatial data, only the most important input data and methodology descriptions are included in the IIR report. For a more detailed description, please refer to Plejdrup & Gyldenkærne (2011).




Figure 9.1 Map of Denmark including names of regions and the Exclusive Economic Zone.

### 8.2.1 The SPREAD model

The distribution in SPREAD is made on SNAP category level to assure the most accurate distribution of the emissions. It has been aimed to use the most disaggregated SNAP level (SNAP 3 level) but for some categories and sectors SNAP 2 or SNAP 1 level has been applied in the distribution model, due to a lack of detailed geographical information. An exception is the agricultural sector, as this sector is not treated on SNAP level in the Danish inventory system. Instead the agricultural data processing is carried out for the relevant NFR categories, and the same approach is applied in SPREAD. The SPREAD model is prepared in order to be applicable for the mandatory reporting of gridded emissions under CLRTAP.

SPREAD includes a number of sub-models covering separate sources or groups of sources in the emission inventory; Large Point Sources, Stationary combustion for point sources, Stationary combustion for area sources, Mo-

bile sources, Aviation, Fugitive emissions from fuels, Industrial Processes, F-gases, Solvent and other product use, Waste, and a number of sub-models for the agricultural sector. All sub-models correspond to the methodology and groupings in the Danish inventory system. A number of sub-models include a higher disaggregation level compared to the NFR tables. Both SNAP and NFR categories are included in all SPREAD sub-models to enable a distribution in agreement with the international guidelines.

Emissions from all Large Point Sources (LPS) are treated in the LPS sub-model in SPREAD. LPSs represent emissions at all SNAP 1 categories except solvents (SNAP 06) and road traffic (SNAP 07). Further, LPSs in agriculture are included in a separate part of the emission database system covering agriculture and are not included in the LPS sub-model in SPREAD. The Point Sources sub-model covers emissions from stationary combustion from point sources, which refer to the large number of plants, for which the fuel consumption is known at plant level but emissions are calculated using standard emission factors.

### General methodology

The distribution of emissions in the Danish emission inventory is carried out in databases and in a geographical information system, GIS.

The methodology applied in the part of the distribution carried out in GIS is shortly described in this chapter. The description is made for the Industrial Processes sector as a case, as this distribution is rather simple.

The emission inventory for Industrial Processes covers both point sources and area sources. Emissions from point sources are allocated to the coordinates for the individual plants included in the Danish inventory system and are not relevant in relation to the GIS procedure. Emissions from area sources are calculated from production statistics and the resulting emissions are national totals as allocation of the sources (industrial plants) is not possible with the available data. Instead a proxy for the distribution is applied, in this case the location of industrial areas as given in the national topographic map KORT10 by the National Survey and Cadastre (Figure 9.2). The map of industrial areas is not reflecting differences in the location for different industries, but only holds industrial buildings (referred to as the industrial area as the buildings are treated as areas rather than units). The map is a shape file and the industrial areas are polygons.



Figure 9.2 Segment around Avedøre close to Copenhagen of the map of industrial areas (KORT10).

As SPREAD gives emissions on 1 km x 1 km, the map of industrial areas must be combined with the Danish 1 km x 1 km Grid Net. The grid is an orthogonal coordinate system and the cells are defined and named by their lower left corner coordinates. The grid net map is a shape file and the grid cells are polygons (Figure 9.3).



Figure 9.3 Segment around Avedøre in Copenhagen of the map of the Danish 1 km x 1 km grid net (KORT10).

To be able to distribute the emissions on 1 km x 1 km it is necessary to split the industrial polygons between the grid cells and thereby be able to calculate the industrial area in each grid cell (Figure 9.4). These functionalities are available in GIS, in this case ArcMAP. The split is made using the intersect tool, and afterwards the areas are applied to each cell using the Calculate Area function.



Figure 9.4 Segment around Avedøre in Copenhagen of the map of industrial areas and the Danish 1 km x 1 km grid net (KORT10).

The remaining part of the emission distribution for industrial processes is carried out in a database. The share of the national emissions that should be allocated to each grid cell is calculated as the industrial area of the cell divided by the total industrial area. The same distribution key is applied for all pollutants.

In the case of the Industrial Processes sector only one map is combined with the grid, but more maps or layers could be combined to make a distribution

key. This is the case for some sources in the agricultural sector, e.g. emissions from organic soils where the distribution key is based on a map of organic soils, a map of the agricultural fields and the Danish Grid Net. A number of area sources are distributed on line features, e.g. emissions from railways and road traffic. In these cases the lines are split into segments by intersection with the 1 km x 1 km grid net. The emission in each grid cell is calculated as the national emission multiplied by the length of the line segment(s) in the cell and divided by the total length of the line feature.

For some sources the same distribution key can be applied for more or all years, while other sources demands a separate distribution key for every year. For Industrial Processes the distribution key can be applied for more years, as the dataset is not available on annual basis. Further, the industrial area does not change much from year to year. In other cases the distribution keys must be set up on annual basis as large changes occur from year to year. This is the case for e.g. agricultural soils and point sources (PS) in the energy sector.

#### **National geographical data**

A large number of national geographical data sets are implemented in the SPREAD model in preparation of the various distribution keys. The data sets are listed in Table 9.2 with specification of data owner and a short description of the content of each data set.

Table 9.2 List of geographic data applied in the emission gridding.

Data owner	Data set	Contents
The National Survey and Cadastre	Topographic map	Geo-referenced basic map layers on administrative units, Land cover, territorial borders, coastline and infrastructure.
National Agency for Enterprise and Construction	Central Dwelling and Building Register (Danish abbreviation BBR)	Geo-referenced information on dwellings and buildings
Danish Ministry of the Environment	The Area Information System (AIS)	National maps of spatial data related to nature and environment (e.g. railways, industrial areas and one-storey settlements)
The Directorate for Food, Fisheries and Agri Business	The Central Husbandry Register (CHR)	Information on stock of livestock at farm level
	The General Agricultural Register (GLR)	Information on agricultural farms and crops on field level
Ministry of food, agriculture and fisheries	The fertilizer and husbandry register (Danish abbreviation GHI)	Information on manure and fertiliser amounts on farm level
	The Land Parcel Identification System (LPIS)	Geo-referenced data on agricultural land parcels, including field IDs for fields located in the parcels
The Central Business Register	Central Business Register (Danish abbreviation CVR)	Geo-referenced information on businesses with a CVR number, e.g. farms
The Central Office of the Civil Registration	The Civil Registration System (Danish abbreviation CPR)	Geo-referenced information on population on address level
The Department of Environmental Science, Aarhus University	National road and traffic database	Geo-referenced traffic load on the Danish road network
	Energy producer accountings	Geo-referenced information on fuel consumption for district heating and/or power producing plants
The Danish Energy Agency	The regional inventory	Regional inventory of energy consumption for heating for oil boilers, natural gas boilers and solid fuel installations on municipality level
	Large Point Sources (LPS)	Geo-referenced information on power plants, large industrial plants and offshore installations
DCE - Danish Centre for Environment and Energy		
Danish Petroleum association	Service stations	Geo-referenced information on addresses for all Danish service stations
Energinet.dk	Measurement and regulator stations	Geo-referenced information on location of measurement and regulator stations in the Danish natural gas transmission network
Danish Forest and Nature Agency	Military training terrain	Geo-referenced information on military training terrains
The Danish Environmental Protection Agency	Information system for waste and recycling (Danish abbreviation ISAG)	Data on waste treatment companies on address level
Miljøportalen.dk	Waste water treatment plants	Data on waste water treatment on facility level, including flow rates and organic matter content

### 8.3 Gridded emission data

In this section selected maps of gridded emissions are presented, all referring to the year 2010. The selected maps in Figure 9.5 illustrate the emissions included in the national total in the NFR table (all emissions excluding Civil Aviation - Domestic and International Cruise, and international Maritime Navigation). All figures illustrate the sum of all included GNFR sectors. The Danish high resolution gridded emissions are aggregated on the 50 km x 50 km EMEP grid for reporting to CLRTAP. The share of each 1 km x 1 km grid cell located in the relevant EMEP grid cells are calculated and the aggregated emissions are calculated as the weighted sum of emissions in the 1 km

grid cells intersecting each EMEP grid cell being partial or fully part of the Danish Exclusive Economic Zone, EEZ.




Figure 9.5 National total gridded emissions excluding civil aviation and international navigation of a)  $\text{NO}_x$ , b) NMVOC, c)  $\text{SO}_2$ , d)  $\text{NH}_3$ , e)  $\text{PM}_{2.5}$  and f) PAHs (the sum of benzo(b)flouranthen, benzo(k)flouranthen, benzo(a)pyrene and indeno(1,2,3-c,d)pyrene) for the year 2010.

Even on the 50 km x 50 km aggregated level spatial patterns from the major sectors are recognisable for different pollutants.

### 8.3.1 $\text{NO}_x$

The major GNFR source to  $\text{NO}_x$  emissions is RoadRail followed by Shipping, OffRoadMob, PublicPower and IndustrialComb contributing 36 %, 16 %, 15 %, 15 % and 11 %, respectively. The pattern of the gridded  $\text{NO}_x$  emissions reflect the major road and rail network located in the eastern part of Jutland

and across Funen and Zealand to Copenhagen (figure 9.5). Further, large emissions from PublicPower and IndustrialComb are seen around the major cities. Part of the fugitive emissions is located offshore due to extraction of oil and gas on the North Sea.

### 8.3.2 NMVOC

The major source of NMVOC is Solvents followed by SmallComb, RoadRail, OffRoadMob and Fugitive contributing 31 %, 18 %, 15 %, 11 % and 11 %, respectively. Both emissions from Solvents, SmallComb and OffRoadMob are to a large degree allocated according to population density and location of one-storey settlements. Part of the fugitive emissions is located offshore due to extraction of oil and gas on the North Sea.

### 8.3.3 SO<sub>2</sub>

The major sources of SO<sub>2</sub> are PublicPower and IndustrialComb followed by SmallComb and Shipping contributing 27 %, 26 %, 19 %, and 13 %, respectively. Even though the SO<sub>2</sub> emission has decreased over the years due to implementation of techniques for reduction of sulphur in the flue gas, it still produces a distinct pattern reflecting the localisation of large power plants in Denmark. The allocation of emissions from IndustrialComb reflect the location of a large number of CHP plants not reported as LPS due to no plant specific emission factors. The allocation of emissions from SmallComb reflects the areas with high population density and mainly one-storey settlements.

For the ferries operating between Copenhagen and Bornholm part of the route is outside the Danish EEZ. The emissions from all these ferries are included in Shipping and distributed on the part of the straight line between Copenhagen and Bornholm inside the Danish EEZ. This leads to an aggregation of the emissions in few EMEP cells, and thereby artificial high emissions at the part of the route inside the EEZ.

### 8.3.4 NH<sub>3</sub>

The agricultural sector is by far the major contributor to the NH<sub>3</sub> emission. 81 % of the national emissions excluding civil aviation and international navigation derive from AgriLivestock and another 15 % from AgriOther. Emission of NH<sub>3</sub> is mainly related to livestock farming and especially to manure management. Emissions are distributed according to very detailed data on animals and fields, and the geographical pattern is in good agreement with the localisation of the major Danish livestock farming in Jutland.

### 8.3.5 PM<sub>2.5</sub>

The major source of PM<sub>2.5</sub> emissions is SmallComb contributing 73 %. Road-Rail is the second largest source contributing 10 % of the PM<sub>2.5</sub> emission. Emissions from SmallComb are allocated rather evenly on the land area as a major source is residential wood combustion. Emissions from the residential sector are distributed on municipality level leading to equal emissions for larger areas. Further emissions from CHP plants are located in all parts of the country, also leading to a rather even distribution.

### 8.3.6 PAHs

Emissions of PAHs are the sum of benzo(b)flouranthene, benzo(k)flouranthene, benzo(a)pyrene and indeno(1,2,3-c,d)pyrene. The major source to

emissions of PAHs in Denmark is SmallComb and hereof the all-important source is residential wood combustion. As described for PM<sub>2.5</sub> the distribution are made on municipality level leading to a rather even distribution on the land area.

#### **8.4 References**

EMEP/EEA, 2009: EMEP/EEA air pollutant emission inventory guidebook 2009. Technical guidance to prepare national emission inventories. EEA Technical Report 9/2009. Available at:

<http://www.eea.europa.eu/publications/emep-eea-emission-inventoryguidebook-2009> (10-02-2011).

Plejdstrup, M.S. & Gyldenkerne, S. 2011: Spatial distribution of emissions to air – the SPREAD model. National Environmental Research Institute, Aarhus University, Denmark. 72 pp. – NERI Technical Report no. FR823. Available at: <http://www.dmu.dk/Pub/FR823.pdf>

UNECE, 2009: Guidelines for estimating and reporting emission data under the Convention on Long-range Transboundary Air Pollution (ECE/EB.AIR/97). Available at:

[http://www.ceip.at/fileadmin/inhalte/emep/reporting\\_2009/Rep\\_Guidelines\\_ECE\\_EB\\_AIR\\_97\\_e.pdf](http://www.ceip.at/fileadmin/inhalte/emep/reporting_2009/Rep_Guidelines_ECE_EB_AIR_97_e.pdf)



## 9 Recalculations and Improvements

In general, considerable work is being carried out to improve the inventories. Investigations and research carried out in Denmark and abroad produce new results and findings which are given consideration and, to the extent which is possible, are included as the basis for emission estimates and as data in the inventory databases. Furthermore, the updates of the EMEP/EEA Guidebook, and the work of the Task Force on Emission Inventories and its expert panels are followed closely in order to be able to incorporate the best scientific information as the basis for the inventories.

The implementation of new results in inventories is made in a way so that improvements, as far as possible, better reflect Danish conditions and circumstances. This is in accordance with good practice. Furthermore, efforts are made to involve as many experts as possible in the reasoning, justification and feasibility of implementation of improvements.

In improving the inventories, care is taken to consider implementation of improvements for the whole time series of inventories to make it consistent. Such efforts lead to recalculation of previously submitted inventories. This submission includes recalculated inventories for the whole time series. The reasoning for the recalculations performed is to be found in the sectoral chapters of this report. The text below focuses on recalculations, in general, and further serves as an overview and summary of the relevant text in the sectoral chapters. For sector specific planned improvements please also refer to the relevant sectoral chapters.

### 9.1 Energy

Improvements and updates of the Danish energy statistics are made regularly by the producer of the statistics, the Danish Energy Agency. In close cooperation with the DEA, these improvements and updates are reflected in the emission inventory for the energy sector. The Danish energy statistics have, for the most part, been aggregated to the SNAP categorisation.

The inventories are still being improved through work to increase the number of large point sources, e.g. power plants, included in the databases as individual point sources. Such an inclusion makes it possible to use plant-specific data for emissions, etc., available e.g. in annual environmental reports from the plants in question.

#### 9.1.1 Stationary Combustion

For stationary combustion plants, the emission estimates for the years 1990-2012 have been updated according to the latest energy statistics published by the Danish Energy Agency. The update included both end use and transformation sectors as well as a source category update. The changes in the energy statistics are largest for the years 2010, 2011 and 2012.

The consumption of wood in residential plants in 2012 is 4% lower in the revised energy statistics than in the energy statistics applied last year. This causes a lower emission of PM, PCDD/F, PAHs and PCB reported for 2012 this year.

A NH<sub>3</sub> emission factor for residential wood combustion has been added and this cause the higher NH<sub>3</sub> emission reported from stationary combustion this year.

The NMVOC emission factor for straw combusted in agricultural plants has been revised according to EEA (2013). This has caused a higher reported emission of NMVOC in this source category.

Double counting for SO<sub>2</sub> and NO<sub>x</sub> in mineral wool production plants in 1985-1994 have been corrected and cause a decrease for the industrial plants.

HM emission factors for residential wood combustion have been revised according to EEA (2013). This has caused higher reported emission values for Cd, Cr and Zn and lower reported emission values for As, Cu and Cu.

Emission factor time series for residential wood combustion have been estimated for 1990-2000. In previous inventories the emission factors were assumed constant in 1990-2000. This cause higher emission reported for residential plants for PAHs, PCDD/F, PCB, NMVOC and CO in 1990-2000.

An emission inventory for BC has been added.

### **9.1.2 Mobile sources**

#### **Road transport**

Based on the updated version of COPERT IV version 11 launched in 2014, fuel consumption and NO<sub>x</sub>, VOC, CO and PM emission factors for euro 5 and 6 gasoline and diesel passenger cars and light duty vehicles have been updated in the model. Updated Euro V and VI fuel consumption and NO<sub>x</sub>, VOC, CO and PM emission factors for heavy duty vehicles have been included in the calculations also.

For N<sub>2</sub>O and NH<sub>3</sub>, Euro 5/6 and V/VI emission factors are also updated for passenger cars/light duty vehicles (only N<sub>2</sub>O) and heavy duty vehicles.

Further a new Euro 6c technology class has been added for diesel passenger cars and light duty vehicles.

The amount of diesel sold for road transport reported by the Danish Energy Agency has been slightly changed in 2012.

Very small changes in mileage data has been made for the years 1985-2012 based on new information from DTU Transport.

The percentage emission change interval and year of largest percentage differences (low %; high %, year) for the different emission components are: SO<sub>2</sub> (0 %; 0.2 %, 2011), NO<sub>x</sub> (-1.4 %; 0 %, 2012), NMVOC (-0.5 %; 0 %, 2012), NH<sub>3</sub> (-2 %; 0.6 %, 2012) and TSP (-2.3 %; 0.1 %, 2012).

#### **Navigation**

Recreational craft have been regrouped in the Danish inventory. These vessels have now been removed from the navigation sector and included under Other (1A5b), the latter sector according to its sector subtitle also comprise recreational craft. Further, small amounts of LPG and kerosene previously included in the navigation sector has now been transferred to stationary sources.

The following largest percentage differences (in brackets) for domestic navigation are noted for: SO<sub>2</sub> (-5.6 %), NO<sub>x</sub> (-12.3 %), NMVOC (-80 %), NH<sub>3</sub> (-100 %) and TSP (-42 %).

#### **Agriculture/forestry**

The baseline emission factors of NO<sub>x</sub>, TSP, CO and VOC and the transient factors for fuel consumption, NO<sub>x</sub>, TSP, CO and VOC for diesel machinery has been slightly changed in the calculation model used to estimate the emissions from Danish non road mobile machinery. Further, the CH<sub>4</sub> fraction of VOC has been updated. The changes are made based on new emission knowledge published by IFEU (1999) for baseline emission factors and CH<sub>4</sub> fractions, and IFEU (2014) for transient factors.

The number of agricultural tractors has been regrouped into finer engine size intervals for all inventory years. The total number of agricultural tractors and harvesters has been updated for the years 2006-2012 based on new stock data from Statistics Denmark for the year 2013.

The following largest percentage differences (in brackets) for agriculture/forestry are noted for: SO<sub>2</sub> (-16.2 %), NO<sub>x</sub> (-17.7 %), NMVOC (-5.3 %), NH<sub>3</sub> (-13.4 %) and TSP (13.6 %).

#### **Fisheries**

Small amounts of LPG and kerosene previously included in the navigation sector has now been transferred to stationary sources.

The following largest percentage differences (in brackets) for fisheries are noted for: SO<sub>2</sub> (0.1 %), NO<sub>x</sub> (-0.8 %), NMVOC (-5.1 %), NH<sub>3</sub> (0.0 %) and TSP (-0.1 %).

#### **Industry**

The baseline emission factors of NO<sub>x</sub>, TSP, CO and VOC and the transient factors for fuel consumption, NO<sub>x</sub>, TSP, CO and VOC for diesel machinery has been slightly changed in the calculation model used to estimate the emissions from Danish non road mobile machinery. Further, the CH<sub>4</sub> fraction of VOC has been updated. The changes are made based on new emission knowledge published by IFEU (1999) for baseline emission factors and CH<sub>4</sub> fractions, and IFEU (2014) for transient factors.

The following largest percentage differences (in brackets) for industrial non road machinery are noted for: SO<sub>2</sub> (0.7 %), NO<sub>x</sub> (1.3 %), NMVOC (4.3 %), NH<sub>3</sub> (1.8 %) and TSP (5.7 %).

#### **Civil aviation**

The model used for calculating civil aviation emissions has been updated by replacing the previous fuel consumption and emission factors for representative aircraft types (46 types) with a new and more comprehensive list of aircraft types (79 types) provided by Eurocontrol and published in the EMEP/EEA guidebook (EMEP/EEA, 2014).

The following largest percentage differences (in brackets) for civil aviation are noted for: SO<sub>2</sub> (38 %), NO<sub>x</sub> (35 %), NMVOC (-39 %), and TSP (30 %).

### **Other (Military and recreational craft)**

Recreational craft have been regrouped in the Danish inventory. These vessels have now been removed from the navigation sector and included under Other (1A5b), the latter sector according to its sector subtitle also comprises recreational craft.

NO<sub>x</sub>, CO, VOC and TSP emission factors have been updated for diesel and gasoline boats complying with Directive 2003/44 based on reassessment of the emission factor data reported by IFEU (2004). The latter reference has all the time been the source of emission data for recreational craft in the Danish emission inventory.

The following largest percentage differences (in brackets) for military are noted for: SO<sub>2</sub> (651 %), NO<sub>x</sub> (218 %), NMVOC (2800 %), NH<sub>3</sub> (149 %) and TSP (1047 %).

### **9.1.3 Fugitive emissions**

#### **Exploration of oil and gas**

Exploration has been included as a new source in the emission inventory as activity data has now been made available by the Danish Energy Agency. Emissions only occur in years with exploration and appraisal wells (E/A wells) (1990-2000, 2002, 2005, 2009 and 2013). The largest E/A productions occurred in 1990 and 2002, contributing 2.4 % and 3.7 % to the fugitive NO<sub>x</sub> emission, respectively, while the corresponding shares for SO<sub>2</sub> and NMVOC are ≤ 0.1 %.

#### **Refineries**

In cooperation with one refinery, the SO<sub>2</sub> emissions have been examined. This work has led to a reallocation of part of the SO<sub>2</sub> emissions in the years 2005-2012. SO<sub>2</sub> emissions from the ammonium thiosulphate (ATS) unit have been reallocated to the flaring category for the years 2008-2012, as the refinery has informed that H<sub>2</sub>S from the ATS unit is being led to the flare. In previous inventory these emissions were allocated to the category refining/storage.

For 2012 the SO<sub>2</sub> emission has been updated according to supplementing information from one refinery, leading to a decrease of 8.1 ktonnes SO<sub>2</sub>. Also a small correction of the fuel gas consumption has been applied according to an update in the annual environmental report.

The methodology for estimating NMVOC from one refinery has been changed, resulting in an increase of the NMVOC emissions for the years 1994-2012. The refinery report annual VOC emissions based on results from measurement campaigns, the last in carried out in 2006. In previous inventories, the split between NMVOC and CH<sub>4</sub> given by the refinery has been used. This methodology has been changed, as the CH<sub>4</sub> share was much higher than for the other Danish refinery, and also much higher than corresponding shares found in a literature study. The CH<sub>4</sub> share has therefore been changed from 56 % to 10 % as given by the other Danish refinery and supported by shares of 10-20 % found for Swedish refineries. The largest increase of the NMVOC emissions are estimated for the years 2006-2012 (1611 tonnes per year), corresponding 10 % - 16 % of the total fugitive NMVOC emissions (16 % in 2011 and 2012).

### **Service stations**

The activity data has been updated for 2011-2012 according to the latest energy statistics published by the Danish Energy Agency. Also an error for the emission factors has been corrected for the years 1992-2012. The recalculation has changed the NMVOC emission by -0.711 ktonnes (2005) to +0.738 ktonnes (1992), which is insignificant compared to the total fugitive NMVOC emission (< 0.01%).

### **Natural gas transmission and distribution**

Activity data has been updated for one town gas distribution company for the year 2012. The change of 0.27 tonnes NMVOC has insignificant impact on the total fugitive emissions (< 0.01%).

### **Venting**

Activity data has been corrected for one gas storage plant for 2012 according to the annual environmental report. The change is insignificant for the total fugitive emissions (< 0.01%).

### **Flaring in refineries**

The NO<sub>x</sub> emission factor has been updated for the years 1994-1996 according to the 2013 EMEP/EEA Guidebook. The changes of the emissions are largest in 1994 with an increase of 2 ktonnes NO<sub>x</sub>, corresponding 0.4 % of the total fugitive emissions.

### **Flaring in gas storage and treatment plants**

NO<sub>x</sub> emissions are updated according to the environmental report for the gas treatment plant for 2012. The decrease of the NO<sub>x</sub> emission is due to installation of new low-NO<sub>x</sub> burner. NO<sub>x</sub> has been updated from 45 tonnes to 0.007 tonnes. The decrease in the NO<sub>x</sub> emission accounts for 0.02 % of the total fugitive NO<sub>x</sub> emission.

### **Flaring in gas transmission and distribution**

Flaring in gas transmission and distribution has been included as a new source in the emission inventory, only occurring in the years 2011-2013. The gas transmission company has provided information that they have started using a mobile flare in large construction works, and also one distribution company is flaring gas. The change of 0.4 tonnes NMVOC has insignificant impact on the total fugitive emissions (< 0.01%).

## **9.2 Industrial processes**

### **9.2.1 Lime production**

The activity data for lime no longer includes slacked lime. Personal communication with the industry made it clear that the inclusion of slacked lime resulted in a double counting because statistic on burned lime also includes lime that is later slacked.

An emission factor for black carbon has been added for lime production.

A factor 1000 error for the PCDD/F emission factor has been corrected for the years 1991-1999 and 2001-2003, resulting in increased emissions for these years.

### 9.2.2 Glass production

The activity data for container glass production has been altered for 1997-2012. Emissions were previously calculated based on fuel consumptions but this has now been changed to total production (mass product). This change necessitates a change and reassessment of the heavy metal emission factors. Where direct measured emissions are not available, Guidebook (2013) emission factors are now used to calculate emissions of heavy metals; As, Cd, Cr and Ni are new in this submission.

NMVOC from glass wool was not reported prior to 2010 causing a jump in the implied emission factor for NMVOC. In addition to the direct emissions of NMVOC known from the environmental reports, data on NMVOC has now been estimated for 1990-2006 using the average implied emission factor for 2007-2009.

Lesser changes were made for CO and NH<sub>3</sub> emissions from glass wool production for 1990-2012 and 1990-1995 respectively. Calculated emissions of these two pollutants were reassessed resulting in a small decrease of CO and increase of NH<sub>3</sub>. Particle size distribution for PM<sub>2.5</sub> was changed from 70 % of TSP to 80 % of TSP in accordance with the Guidebook (2013).

### 9.2.3 Other mineral products

All activity data for the three source categories in "Other mineral products" have been changed to mass of CaCO<sub>3</sub> equivalent consumption instead of mass product produced. This was done because data on the mass produced is impossible to obtain, and uncertainties on the estimated data were unreasonably high. The consumption of raw materials, such as lime, on the other hand are available from 2006 and are easily estimated back to 1990 using surrogate data. All emission factors are changed to match the new activity data.

A factor 1000 error in the bricks and expanded clay production emission factors for PCDD/F has been corrected for the years 1991-1999 and 2001-2003, resulting in increased emissions for these years. For the same categories, a calculation error has resulted in lesser corrections for SO<sub>2</sub>.

### 9.2.4 Chemical industry

For the production of nitric and sulphuric acid, activity data and emission factors are now completed for the time series back to 1980.

The available data for production of catalysts and tar products was reassessed, leading to minor changes for 1990-1996 and 2007-2012.

### 9.2.5 Metal industry

For the electro steelwork, emission factors for As, Cr, Cu and PCDD/F have been added to this year's inventory. Emission factors for Cd, Hg, Ni, Pb and Zn were reassessed and are now calculated based on the measurements provided by the factory's environmental reports, the change in emission factors have led to a strong increase in emissions. The emission factor for TSP was lowered for 2001 down to the measured level of 2000.

The source of secondary aluminium production has been reassessed for this year's inventory. The work on this source category has resulted in all new activity data (4-26 % increase) and a completion of the time series; meaning

that 1990-1994 and 2009-2013 are new years for this source. For the emission factors, particles are now calculated based on measurements from the industry, this change has caused a decrease. The PCDD/F emission factor now refers to the EMEP/EEA Guidebook (2013); this change has caused an increase in PCDD/F emissions. Cd, Pb, HCB and PCB from aluminium production are new in this year's inventory.

The source of secondary lead production has also been reassessed. The work on this source category has resulted in all new activity data (from 5 Gg in all years to 0.3-1.2 Gg in the new time series). Emission factors for As, Zn and PCDD/F are new in this year's inventory while Cd, Pb and particle emission factors now refer to the EMEP/EEA Guidebook (2013) resulting in large increases in emissions. The emission factor for Cu was removed due to the lack of a reference.

Secondary zinc production (SNAP 030308) does not occur in Denmark and never has. The reporting of emissions in this category was the result of a mistake that has now been corrected.

### **9.2.6 Non-energy products from fuels and solvent use**

The amount of solvent, which is added to the asphalt in "cutback", is comprised in Other solvent use (NFR 2D3i), above with an emission factor of approximately unity. This amount was previously included in Road paving with asphalt (NFR 2D3b) as "cutback". In the improved inventory NMVOC emissions from "cutback" asphalt in Road paving (NFR 2D3b) only include emissions from the asphalt fraction.

A change in allocation of amounts from Statistics Denmark (2014) has caused an increase in activity data for roofing, e.g. 2012: from 75.5 Gg to 131 Gg, and a relatively small increase for road paving, e.g. for 2012: from 3223 Gg to 3233 Gg.

Emission factors for NMVOC and CO have been updated with values from EMEP/EEA (2013) and US EPA (2004) for road paving and asphalt roofing.

### **9.2.7 Other production**

New source categories that were added in this year's inventory include:

- Palmin (added to Margarine and solid cooking fats)
- Caffeine free coffee (added to coffee roasting)
- Wine (separated in red and white)

1980-1989 data was reassessed for all categories. As a result of the assessment 1980-1984 data were added to this year's inventory and 1985-1989 are now actual statistical data and not estimates.

An error was corrected for beer production resulting in an increase in activity data for 1999 of 18 million litres.

The activity data for meat curing has been drastically reduced. Previously, the dataset included the entire meat production, but this has now been corrected to only meat that is fried, smoked or otherwise treated in ways that results in NMVOC emissions. The overall reduction for meat curing (including poultry and fish) is 82-89 %.

It was investigated whether it was possible to separate the production of spirits into categories depending on whether the spirit is matured or not. As a result, the inventory now separates between malt whisky and other spirits.

It was investigated whether the bread production could be subdivided to allow for the application of more disaggregated emission factors. This was however not possible with the available statistical data.

The activity data for slaughter house waste was reassessed with the following results: The dataset was extended back to 1980, some increases in activity for the beginning of the 90's and a few changes for other years (2001, 2002, 2006 and 2012).

### **9.3 Agriculture**

Emission of NH<sub>3</sub> from manure management (NFR 3B) is decreased because the emission from manure applied to soils is now reported under NFR 3D. The emission of NH<sub>3</sub> from sewage sludge applied to soils and cultivated crops are now reported under sector 3D instead of under 3I.

The emission of NMVOC has increased a lot due to implementation of NMVOC for manure management (3B) for the years 1985 to 2013. The emission factors for NMVOC emission from cultivated crops (3D) has been updated to the emission factors given in the EMEP/EEA 2013 Guidebook.

The emission of PM has increased a lot due to implementation of emission from farm-level agricultural operations including storage, handling, and transport of agricultural products. The emission from manure management has only changes up to 2 % due to change in number of animals.

The emission of NO<sub>x</sub> has been estimated for manure management (3B), inorganic N-fertilizers (3Da1) and sewage sludge applied to soils (3Da2b) for the years 1985 to 2013.

Some changes in activity data has been made; Number of animals has been changed due to updated numbers for swine in 2011, broilers in 2007-2012 and laying hens in 2012. This affects the emission of NH<sub>3</sub> and PM. Numbers of sheep has been divided in two subcategories – sheep and lambs – with emission factors for each subcategories. This increases the number of animals and affects the emission of NH<sub>3</sub> from manure management (3B) and urine and dung deposited by grazing animals (3D). N from sewage sludge has been updated for 2006 and this affects the emission of NH<sub>3</sub>. The area for field burning of agricultural residues has been changed due to updated data for 2012 and this affects emission of all pollutants.

### **9.4 Waste**

A new reporting template has been used for the 2015 submission in accordance with the reporting guidelines. The main change in the waste sector is that emissions from composting are now reported in a separate category rather than under 'Other waste' as in previous submissions.

Minor changes were made to the activity data for composting and accidental fires to incorporate new information and correct errors in the previous reporting.



## 10 Projections

Projections of emissions are carried out by DCE at irregular time intervals. The most recent projection was made in 2013, projecting the emissions of NO<sub>x</sub>, SO<sub>2</sub>, NMVOC, NH<sub>3</sub>, TSP, PM<sub>10</sub> and PM<sub>2.5</sub> to 2035.

The total projected emissions for these pollutants for 2015, 2020, 2025, 2030 and 2035 are shown in the table below. For further documentation, please refer to Nielsen et al. (2013).

Table 10.1 Projected emissions for 2015, 2020, 2025, 2030 and 2035, tonnes.

Pollutant	2015	2020	2025	2030	2035
SO <sub>2</sub>	13 656	12 270	12 207	12 435	14 061
NO <sub>x</sub>	105 244	83 006	73 265	69 251	67 095
NMVOC	73 384	68 386	65 760	65 283	64 894
NH <sub>3</sub>	71 442	70 236	67 294	64 696	63 617
TSP	34 576	31 526	30 658	30 296	29 261
PM <sub>10</sub>	26 296	23 029	21 912	21 173	20 169
PM <sub>2.5</sub>	20 218	16 734	15 411	14 371	13 340

### 10.1 References

Nielsen, O.-K., Plejdrup, M., Hjelgaard, K., Nielsen, M., Winther, M., Mikkelsen, M.H., Albrektsen, R., Fauser, P., Hoffmann, L. & Gyldenkærne, S. 2013. Projection of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and particle emissions - 2012-2035. Aarhus University, DCE – Danish Centre for Environment and Energy, 151 pp. Technical Report from DCE – Danish Centre for Environment and Energy No. 81. Available at: <http://dce2.au.dk/pub/SR81.pdf>

## 11 Adjustments

Decision 2012/3 of the Executive Body (UNECE, 2012a) decided that adjustments may be made under specific circumstances to the national emission inventories for the purpose of comparing the inventories with emission reduction commitments.

The circumstances under which an adjustment may be applied fall into three broad categories where:

- Emission source categories are identified that were not accounted for at the time when emission reduction commitments were set;
- Emission factors used to determine emissions levels for particular source categories for the year in which emissions reduction commitments are to be attained are significantly different than the emission factors applied to these categories when emission reduction commitments were set;
- The methodologies used for determining emissions from specific source categories have undergone significant changes between the time when emission reduction commitments were set and the year they are to be attained.

The supporting documentation required by Parties applying for an adjustment is set out in Decision 2012/12 (UNECE, 2012b) and is summarised below.

A Party's supporting documentation for an adjustment to its emission inventory or emission reduction commitments shall include:

- Evidence that the Party exceeds its emission reduction commitments;
- Evidence of to what extent the adjustment to the emission inventory reduces the exceedance and possibly brings the Party in compliance;
- An estimation of whether and when the reduction commitment is expected to be met based on emission projections without the adjustment, thereby using best available science;
- A full demonstration that the adjustment is consistent with one or more of the three broad categories above. Reference can be made, as appropriate, to relevant previous adjustments:
  - For new emission source categories:
    - Evidence that the new emission source category is acknowledged in scientific literature and/or the EMEP/EEA air pollutant emission inventory guidebook;
    - Evidence that this source category was not included in the relevant historic national emission inventory at the time when the emission reduction commitment was set;
    - Evidence that emissions from a new source category contribute to a Party being unable to meet its reduction commitments, supported by a detailed description of the methodology, data and emission factors used to arrive at this conclusion;
  - For significantly different emission factors used for determining emissions from specific source categories:
    - A description of the original emission factors, including a detailed description of the scientific basis upon which the emission factor was derived;

- Evidence that the original emission factors were used for determining the emission reductions at the time when they were set;
- A description of the updated emission factors, including detailed information on the scientific basis upon which the emission factor was derived;
- A comparison of emission estimates made using the original and the updated emission factors, demonstrating that the change in emission factors contributes to a Party being unable to meet its reduction commitments; and
- The rationale for deciding whether the changes in emission factors are significant;
- For significantly different methodologies used for determining emissions from specific source categories:
  - A description of the original methodology used, including detailed information on the scientific basis upon which the methodology was based;
  - Evidence that the original methodology was used for determining the emission reductions at the time when they were set;
  - A description of the updated methodology used, including a detailed description of the scientific basis or reference upon which it has been derived;
  - A comparison of emission estimates made using the original and updated methodologies demonstrating that the change in methodology contributes to a Party being unable to meet its reduction commitment; and
  - The rationale for deciding whether the change in methodology is significant;

## 11.1 Accepted adjustments

In the 2014 submission Denmark applied for two adjustments related to the emission of NH<sub>3</sub>, due to exceedance of the emission ceiling. One was related to ammonia from growing crops, which was a new emission source compared to when the emission reduction commitments were agreed. The other was related to the new NH<sub>3</sub> emission factors for inorganic fertilisers included in the 2013 EMEP/EEA Guidebook. The two adjustments were accepted during the technical review and approved by the EMEP Steering Body. The details on the two adjustments are included below.

### 11.1.1 NH<sub>3</sub> emissions from inorganic fertilisers

The 2013 EMEP/EEA Guidebook contained updated EFs for NH<sub>3</sub> from the use of synthetic fertilizer. These emission factors are unlike the emission factors in the previous version of the EMEP/EEA Guidebook not temperature dependent. This means that the current emission factors are significantly higher compared to the previous emission factors.

The NH<sub>3</sub> emission from inorganic fertilisers (NFR category 3Da1) using both the new emission factors and the original emission factors is shown in Table 11.1 below.

Table 11.1 Overview of the adjusted and unadjusted NH<sub>3</sub> emission from inorganic fertilisers

NH <sub>3</sub> emission, kt	2010	2011	2012	2013
NH <sub>3</sub> emission from inorganic fertilisers using new EFs	7.14	7.36	7.06	7.29
NH <sub>3</sub> emission from inorganic fertilisers using old EFs	3.47	3.94	3.77	3.54
Adjustment	-3.67	-3.42	-3.30	-3.75

The numbers presented in Table 11.1 for 2010 to 2012 are identical to the numbers included in the expert review report (CEIP, 2014). The 2013 value is estimated using the same methodology as the methodology presented to and approved by the expert review team.

### 11.1.2 NH<sub>3</sub> from cultivated crops

NH<sub>3</sub> emissions from cultivated crops are acknowledged in the EMEP/EEA Guidebook, but no default emission factor is provided. Denmark uses a country specific emission factor to estimate emissions from cultivated crops as documented in Chapter 5. This source was not included in the consideration when establishing the emission ceiling neither is it included in the GAINS model.

The NH<sub>3</sub> emission from cultivated crops (NFR category 3De) is shown in Table 11.2 below.

Table 11.2 NH<sub>3</sub> emission from cultivated crops (NFR category 3De).

NH <sub>3</sub> emission, kt	2010	2011	2012	2013
NH <sub>3</sub> emission from cultivated crops	5.41	5.42	5.40	5.37
Adjustment	-5.41	-5.42	-5.40	-5.37

The numbers presented in Table 11.2 for 2010 to 2012 are identical to the numbers included in the expert review report (CEIP, 2014). The 2013 value is estimated using the same methodology as the methodology presented to and approved by the expert review team.

### 11.1.3 Total effect of approved adjustments

The total effect of the approved adjustments is documented in Table 11.3 below. The emission ceiling for NH<sub>3</sub> for Denmark was 69 kt.

Table 11.3 Total effect of adjustments

Emission, kt	2010	2011	2012	2013
Total NH <sub>3</sub> adjustment	-9.08	-8.84	-8.70	-9.13
Unadjusted NH <sub>3</sub> emission	79.97	78.83	77.13	74.32
Adjusted NH <sub>3</sub> emission	70.90	69.99	68.43	65.19

## 11.2 Application for adjustment

In the 2015 inventory submission, NMVOC emissions from animal husbandry and manure management (NFR category 3B) have been estimated for the first time. This constitutes a new source in the Danish emission inventory.

The improvement is based on the new guidance contained in the 2013 edition of the EMEP/EEA Guidebook. Previously, no default emission factors had been included in the Guidebook.

The emission from this source is very significant as shown in Table 11.4 and means that Denmark exceeds the emission ceiling. With the proposed adjustment Denmark will be in compliance with our commitments, see Table 11.4.

The latest official Danish projections (Nielsen et al., 2013) did not include this source. However, since the NMVOC emissions from animal husbandry and manure management is strongly correlated with animal numbers there

is no indication that a significant decrease in NMVOC emissions from this source will take place. This implies that with the current projection for the other sources of NMVOC, the Danish NMVOC emissions will still be significantly higher than the emission ceiling even in 2035.

Table 11.4 NMVOC emissions with and without emissions from animal husbandry and manure management (3B).

Emissions, kt	2010	2011	2012	2013
NMVOC excluding 3B	89.86	83.12	79.85	78.30
NMVOC from 3B	35.52	35.45	35.85	36.13
NMVOC including 3B	125.38	118.57	115.70	114.43
Proposed adjustment	-35.52	-35.45	-35.85	-36.13

The emissions from category 3B are based on the default emission factors as contained in Table 3-3 of the EMEP/EEA Guidebook. These are replicated in Table 11.5 below.

Table 11.5 Emission factors used in the estimation.

	EF NMVOC with silage	EF NMVOC without silage
Dairy Cattle	17.937	8.047
Non-Dairy Cattle	8.902	3.602
Sheep	0.279	0.169
Swine – sows		1.704
Swine – other		0.551
Goats	0.624	0.542
Horses	7.781	4.275
Laying hens		0.165
Broilers		0.108
Turkeys		0.489
Other poultry		0.489
Fur animals		1.941

For animals that are fed with silage for part of the year, the emission factors for feeding with silage are assumed to be applicable when the animals are fed in the housing systems, whereas when they are grazing the emission factors without silage are used. More information on animal numbers, grazing days and a more in-depth description of the estimation methodology is available in Chapter 5.

### 11.3 References

CEIP, 2014: Review of the 2014 Adjustment Application by Denmark. CEIP/Adjustment RR/2014/DENMARK. 01/09/2014. Available at: [http://webdab.umweltbundesamt.at/download/adjustments2014/Adjustment\\_Review\\_Report\\_DENMARK\\_%202014.pdf?cgiproxy\\_skip=1](http://webdab.umweltbundesamt.at/download/adjustments2014/Adjustment_Review_Report_DENMARK_%202014.pdf?cgiproxy_skip=1) (29-12-2014).

UNECE, 2012a: Decision 2012/3 'Adjustments under the Gothenburg Protocol to emission reduction commitments or to inventories for the purposes of comparing total national emissions with them'. Available at: [http://www.unece.org/fileadmin/DAM/env/documents/2013/air/ECE\\_EB.AIR.111.Add.1\\_ENG\\_DECISION\\_3.pdf](http://www.unece.org/fileadmin/DAM/env/documents/2013/air/ECE_EB.AIR.111.Add.1_ENG_DECISION_3.pdf) (29-12-2014).

UNECE, 2012b: Decision 2012/12 'Guidance for adjustments under the 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone to emission reduction commitments or to inventories for the purposes of comparing total national emissions with them'. Available at: [http://www.unece.org/fileadmin/DAM/env/documents/2012/EB/Decision\\_2012\\_12.pdf](http://www.unece.org/fileadmin/DAM/env/documents/2012/EB/Decision_2012_12.pdf) (29-12-2014).

Nielsen, O.-K., Plejdrup, M., Hjelgaard, K., Nielsen, M., Winther, M., Mikkelsen, M.H., Albrektsen, R., Fauser, P., Hoffmann, L. & Gyldenkærne, S. 2013. Projection of SO<sub>2</sub>, NO<sub>x</sub>, NMVOC, NH<sub>3</sub> and particle emissions - 2012-2035. Aarhus University, DCE – Danish Centre for Environment and Energy, 151 pp. Technical Report from DCE – Danish Centre for Environment and Energy No. 81. Available at: <http://dce2.au.dk/pub/SR81.pdf>

## **Annex 1 – Key category analysis**

Due to a lack of resources a key category analysis has not been performed for this submission.

## **Annex 2 – Information on the energy balance**

The official Danish energy balance is prepared by the Danish Energy Agency (DEA). The DEA is responsible for reporting of energy data to Eurostat and the IEA. DCE uses the energy balance as published by the DEA. However, some reallocations between sectors are made in connection with the bottom-up modelling done at DCE for different subsectors within transport and mobile sources. For a more in-depth discussion of the energy statistics please see Annex 3A-9. For information on the reallocation of fuels please see Chapter 3.3.



## Annex 3A - Stationary combustion

Annex 3A-1:	Correspondence list for SNAP/NFR
Annex 3A-2:	Fuel rate
Annex 3A-3:	Default Lower Calorific Value (LCV) of fuels and fuel correspondance list
Annex 3A-4:	Emission factors
Annex 3A-5:	Implied emission factors for power plants and municipal waste incineration plants
Annex 3A-6:	Large point sources
Annex 3A-7:	Uncertainty estimates
Annex 3A-8:	Emission inventory 2013 based on SNAP sectors
Annex 3A-9:	Description of the Danish energy statistics
Annex 3A-10:	Time-series 1980/1985 - 2012
Annex 3A-11:	QA/QC for stationary combustion

## Annex 3A-1 Correspondence list for SNAP/CRF

Table 3A-1.1 Correspondence list for stationary combustion SNAP/NFR.

snap_id	snap_name	nfr_id EA	nfr_name
010100	Public power	1A1a	Public electricity and heat production
010101	Combustion plants >= 300 MW (boilers)	1A1a	Public electricity and heat production
010102	Combustion plants >= 50 and < 300 MW (boilers)	1A1a	Public electricity and heat production
010103	Combustion plants < 50 MW (boilers)	1A1a	Public electricity and heat production
010104	Gas turbines	1A1a	Public electricity and heat production
010105	Stationary engines	1A1a	Public electricity and heat production
010200	District heating plants	1A1a	Public electricity and heat production
010201	Combustion plants >= 300 MW (boilers)	1A1a	Public electricity and heat production
010202	Combustion plants >= 50 and < 300 MW (boilers)	1A1a	Public electricity and heat production
010203	Combustion plants < 50 MW (boilers)	1A1a	Public electricity and heat production
010204	Gas turbines	1A1a	Public electricity and heat production
010205	Stationary engines	1A1a	Public electricity and heat production
010300	Petroleum refining plants	1A1b	Petroleum refining
010301	Combustion plants >= 300 MW (boilers)	1A1b	Petroleum refining
010302	Combustion plants >= 50 and < 300 MW (boilers)	1A1b	Petroleum refining
010303	Combustion plants < 50 MW (boilers)	1A1b	Petroleum refining
010304	Gas turbines	1A1b	Petroleum refining
010305	Stationary engines	1A1b	Petroleum refining
010306	Process furnaces	1A1b	Petroleum refining
010400	Solid fuel transformation plants	1A1c	Oil and gas extraction
010401	Combustion plants >= 300 MW (boilers)	1A1c	Oil and gas extraction
010402	Combustion plants >= 50 and < 300 MW (boilers)	1A1c	Oil and gas extraction
010403	Combustion plants < 50 MW (boilers)	1A1c	Oil and gas extraction
010404	Gas turbines	1A1c	Oil and gas extraction
010405	Stationary engines	1A1c	Oil and gas extraction
010406	Coke oven furnaces	1A1c	Oil and gas extraction
010407	Other (coal gasification, liquefaction)	1A1c	Oil and gas extraction
010500	Coal mining, oil / gas extraction, pipeline compressors	1A1c	Oil and gas extraction
010501	Combustion plants >= 300 MW (boilers)	1A1c	Oil and gas extraction
010502	Combustion plants >= 50 and < 300 MW (boilers)	1A1c	Oil and gas extraction
010503	Combustion plants < 50 MW (boilers)	1A1c	Oil and gas extraction
010504	Gas turbines	1A1c	Oil and gas extraction
010505	Stationary engines	1A1c	Oil and gas extraction
010506	Pipeline compressors	1A3e i	Pipeline transport
020100	Commercial and institutional plants	1A4a i	Commercial/institutional: Stationary
020101	Combustion plants >= 300 MW (boilers)	1A4a i	Commercial/institutional: Stationary
020102	Combustion plants >= 50 and < 300 MW (boilers)	1A4a i	Commercial/institutional: Stationary
020103	Combustion plants < 50 MW (boilers)	1A4a i	Commercial/institutional: Stationary
020104	Stationary gas turbines	1A4a i	Commercial/institutional: Stationary
020105	Stationary engines	1A4a i	Commercial/institutional: Stationary
020106	Other stationary equipments	1A4a i	Commercial/institutional: Stationary
020200	Residential plants	1A4b i	Residential: Stationary
020201	Combustion plants >= 50 MW (boilers)	1A4b i	Residential: Stationary
020202	Combustion plants < 50 MW (boilers)	1A4b i	Residential: Stationary
020203	Gas turbines	1A4b i	Residential: Stationary
020204	Stationary engines	1A4b i	Residential: Stationary
020205	Other equipments (stoves, fireplaces, cooking)	1A4b i	Residential: Stationary
020300	Plants in agriculture, forestry and aquaculture	1A4c i	Agriculture/Forestry/Fishing: Stationary
020301	Combustion plants >= 50 MW (boilers)	1A4c i	Agriculture/Forestry/Fishing: Stationary
020302	Combustion plants < 50 MW (boilers)	1A4c i	Agriculture/Forestry/Fishing: Stationary
020303	Stationary gas turbines	1A4c i	Agriculture/Forestry/Fishing: Stationary
020304	Stationary engines	1A4c i	Agriculture/Forestry/Fishing: Stationary
020305	Other stationary equipments	1A4c i	Agriculture/Forestry/Fishing: Stationary
030100	Comb. in boilers, gas turbines and stationary	1A2g viii	Other manufacturing industry
030101	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
030102	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
030103	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
030104	Gas turbines	1A2g viii	Other manufacturing industry
030105	Stationary engines	1A2g viii	Other manufacturing industry
030106	Other stationary equipments	1A2g viii	Other manufacturing industry
030200	Process furnaces without contact (a)	1A2g viii	Other manufacturing industry
030203	Blast furnace cowpers	1A2a	Iron and steel
030204	Plaster furnaces	1A2g viii	Other manufacturing industry
030205	Other furnaces	1A2g viii	Other manufacturing industry
030400	Iron and Steel	1A2a	Iron and steel
030401	Combustion plants >= 300 MW (boilers)	1A2a	Iron and steel

snap_id	snap_name	nfr_id_EA	nfr_name
030402	Combustion plants >= 50 and < 300 MW (boilers)	1A2a	Iron and steel
030403	Combustion plants < 50 MW (boilers)	1A2a	Iron and steel
030404	Gas turbines	1A2a	Iron and steel
030405	Stationary engines	1A2a	Iron and steel
030406	Other stationary equipments	1A2a	Iron and steel
030500	Non-Ferrous Metals	1A2b	Non-ferrous metals
030501	Combustion plants >= 300 MW (boilers)	1A2b	Non-ferrous metals
030502	Combustion plants >= 50 and < 300 MW (boilers)	1A2b	Non-ferrous metals
030503	Combustion plants < 50 MW (boilers)	1A2b	Non-ferrous metals
030504	Gas turbines	1A2b	Non-ferrous metals
030505	Stationary engines	1A2b	Non-ferrous metals
030506	Other stationary equipments	1A2b	Non-ferrous metals
030600	Chemical and Petrochemical	1A2c	Chemicals
030601	Combustion plants >= 300 MW (boilers)	1A2c	Chemicals
030602	Combustion plants >= 50 and < 300 MW (boilers)	1A2c	Chemicals
030603	Combustion plants < 50 MW (boilers)	1A2c	Chemicals
030604	Gas turbines	1A2c	Chemicals
030605	Stationary engines	1A2c	Chemicals
030606	Other stationary equipments	1A2c	Chemicals
030700	Non-Metallic Minerals	1A2f	Non-metallic minerals
030701	Combustion plants >= 300 MW (boilers)	1A2f	Non-metallic minerals
030702	Combustion plants >= 50 and < 300 MW (boilers)	1A2f	Non-metallic minerals
030703	Combustion plants < 50 MW (boilers)	1A2f	Non-metallic minerals
030704	Gas turbines	1A2f	Non-metallic minerals
030705	Stationary engines	1A2f	Non-metallic minerals
030706	Other stationary equipments	1A2f	Non-metallic minerals
030800	Mining and Quarrying	1A2g viii	Other manufacturing industry
030801	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
030802	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
030803	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
030804	Gas turbines	1A2g viii	Other manufacturing industry
030805	Stationary engines	1A2g viii	Other manufacturing industry
030806	Other stationary equipments	1A2g viii	Other manufacturing industry
030900	Food and Tobacco	1A2e	Food processing, beverages and tobacco
030901	Combustion plants >= 300 MW (boilers)	1A2e	Food processing, beverages and tobacco
030902	Combustion plants >= 50 and < 300 MW (boilers)	1A2e	Food processing, beverages and tobacco
030903	Combustion plants < 50 MW (boilers)	1A2e	Food processing, beverages and tobacco
030904	Gas turbines	1A2e	Food processing, beverages and tobacco
030905	Stationary engines	1A2e	Food processing, beverages and tobacco
030906	Other stationary equipments	1A2e	Food processing, beverages and tobacco
031000	Textile and Leather	1A2g viii	Other manufacturing industry
031001	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
031002	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
031003	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
031004	Gas turbines	1A2g viii	Other manufacturing industry
031005	Stationary engines	1A2g viii	Other manufacturing industry
031006	Other stationary equipments	1A2g viii	Other manufacturing industry
031100	Paper, Pulp and Print	1A2d	Pulp, Paper and Print
031101	Combustion plants >= 300 MW (boilers)	1A2d	Pulp, Paper and Print
031102	Combustion plants >= 50 and < 300 MW (boilers)	1A2d	Pulp, Paper and Print
031103	Combustion plants < 50 MW (boilers)	1A2d	Pulp, Paper and Print
031104	Gas turbines	1A2d	Pulp, Paper and Print
031105	Stationary engines	1A2d	Pulp, Paper and Print
031106	Other stationary equipments	1A2d	Pulp, Paper and Print
031200	Transport Equipment	1A2g viii	Other manufacturing industry
031201	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
031202	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
031203	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
031204	Gas turbines	1A2g viii	Other manufacturing industry
031205	Stationary engines	1A2g viii	Other manufacturing industry
031206	Other stationary equipments	1A2g viii	Other manufacturing industry
031300	Machinery	1A2g viii	Other manufacturing industry
031301	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
031302	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
031303	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
031304	Gas turbines	1A2g viii	Other manufacturing industry
031305	Stationary engines	1A2g viii	Other manufacturing industry
031306	Other stationary equipments	1A2g viii	Other manufacturing industry
031400	Wood and Wood Products	1A2g viii	Other manufacturing industry

snap_id	snap_name	nfr_id_EA	nfr_name
031401	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
031402	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
031403	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
031404	Gas turbines	1A2g viii	Other manufacturing industry
031405	Stationary engines	1A2g viii	Other manufacturing industry
031406	Other stationary equipments	1A2g viii	Other manufacturing industry
031500	Construction	1A2g viii	Other manufacturing industry
031501	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
031502	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
031503	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
031504	Gas turbines	1A2g viii	Other manufacturing industry
031505	Stationary engines	1A2g viii	Other manufacturing industry
031506	Other stationary equipments	1A2g viii	Other manufacturing industry
031600	Cement production	1A2f	Non-metallic minerals
031601	Combustion plants >= 300 MW (boilers)	1A2f	Non-metallic minerals
031602	Combustion plants >= 50 and < 300 MW (boilers)	1A2f	Non-metallic minerals
031603	Combustion plants < 50 MW (boilers)	1A2f	Non-metallic minerals
031604	Gas turbines	1A2f	Non-metallic minerals
031605	Stationary engines	1A2f	Non-metallic minerals
031606	Other stationary equipments	1A2f	Non-metallic minerals
032000	Non-specified (Industry)	1A2g viii	Other manufacturing industry
032001	Combustion plants >= 300 MW (boilers)	1A2g viii	Other manufacturing industry
032002	Combustion plants >= 50 and < 300 MW (boilers)	1A2g viii	Other manufacturing industry
032003	Combustion plants < 50 MW (boilers)	1A2g viii	Other manufacturing industry
032004	Gas turbines	1A2g viii	Other manufacturing industry
032005	Stationary engines	1A2g viii	Other manufacturing industry
032006	Other stationary equipments	1A2g viii	Other manufacturing industry

## Annex 3A-2 Fuel rate

Table 3A-2.1 Fuel consumption rate of stationary combustion plants 1990-2013, PJ.

Sum of Fuel_rate_PJ			Year									
fuel_type	fuel_id	fuel_gr_abbr	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
SOLID	101A	ANODIC CARBON										
	102A	COAL	253.4	344.3	286.8	300.8	323.4	270.3	371.9	276.3	234.3	196.5
	103A	FLY ASH (COAL)										
	106A	BROWN COAL BRI.	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0
	107A	COKE OVEN COKE	1.3	1.4	1.2	1.2	1.2	1.3	1.2	1.3	1.3	1.4
LIQUID	110A	PETROLEUM COKE	4.5	4.4	4.3	5.7	7.5	5.3	5.9	6.0	5.3	6.8
	203A	RESIDUAL OIL	32.1	38.3	38.5	32.8	46.2	33.0	37.8	26.6	30.0	23.7
	204A	GAS OIL	61.4	64.9	56.0	62.0	53.9	53.6	58.0	51.0	48.4	47.4
	206A	KEROSENE	5.1	1.0	0.8	0.8	0.7	0.6	0.5	0.4	0.4	0.3
	225A	ORIMULSION						19.9	36.8	40.5	32.6	34.2
	303A	LPG	3.0	2.7	2.4	2.5	2.6	2.7	3.0	2.6	2.8	2.5
308A	REFINERY GAS	14.2	14.5	14.9	15.4	16.4	20.8	21.4	16.9	15.2	15.7	
GAS	301A	NATURAL GAS	76.1	86.1	90.5	102.5	114.6	132.7	156.3	164.5	178.7	187.9
WASTE	114A	WASTE	15.5	16.7	17.8	19.4	20.3	22.9	25.0	26.8	26.6	29.1
	115A	INDUSTR. WASTES										
BIOMASS	111A	WOOD	18.2	20.0	21.0	22.2	21.9	21.8	23.4	23.4	22.9	24.4
	117A	STRAW	12.5	13.3	13.9	13.4	12.7	13.1	13.5	13.9	13.9	13.7
	215A	LIQUID BIOFUELS	0.7	0.7	0.7	0.8	0.2	0.3	0.1	0.0	0.0	0.0
	309A	BIOGAS	0.8	0.9	0.9	1.1	1.3	1.8	2.1	2.4	2.7	2.6
	310A	BIO GASIF GAS					0.1	0.0	0.0	0.0	0.0	0.0
Grand Total			498.9	609.6	549.9	580.6	623.0	600.2	756.8	652.7	615.2	586.4

*Continued*

Sum of Fuel_rate_PJ			Year									
fuel_type	fuel_id	fuel_gr_abbr	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
SOLID	101A	ANODIC CARBON										0.0
	102A	COAL	164.7	174.3	174.7	239.0	182.5	154.0	232.0	194.1	170.5	167.7
	103A	FLY ASH (COAL)										
	106A	BROWN COAL BRI.	0.0	0.0	0.0	0.0					0.0	0.0
	107A	COKE OVEN COKE	1.2	1.1	1.1	1.0	1.1	1.0	1.0	1.1	1.0	0.8
LIQUID	110A	PETROLEUM COKE	6.8	7.8	7.8	8.0	8.4	8.1	8.5	9.2	6.9	5.9
	203A	RESIDUAL OIL	18.8	21.1	26.2	28.6	24.5	21.9	26.1	19.8	15.8	14.7
	204A	GAS OIL	41.2	43.6	38.6	38.8	35.7	31.5	26.4	21.6	21.2	24.5
	206A	KEROSENE	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.1	0.1	0.1
	225A	ORIMULSION	34.1	30.2	23.8	1.9	0.0					
	303A	LPG	2.4	2.2	2.0	2.1	2.2	2.2	2.3	1.9	1.7	1.5
308A	REFINERY GAS	15.6	15.8	15.2	16.6	15.9	15.3	16.1	15.9	14.8	15.4	
GAS	301A	NATURAL GAS	186.1	193.8	193.6	195.9	195.1	187.4	191.1	171.0	171.9	164.9
WASTE	114A	WASTE	29.8	31.3	33.3	35.1	35.3	35.8	36.9	38.1	39.6	37.6
	115A	INDUSTR. WASTES	0.5	1.4	1.9	1.5	2.0	2.0	1.5	1.6	2.0	1.7
BIOMASS	111A	WOOD	27.5	30.8	31.6	38.9	43.9	49.7	52.1	60.3	63.6	66.0
	117A	STRAW	12.2	13.7	15.7	16.9	17.9	18.5	18.5	18.8	15.9	17.4
	215A	LIQUID BIOFUELS	0.0	0.2	0.1	0.4	0.6	0.8	1.1	1.2	1.8	1.7
	309A	BIOGAS	3.0	3.1	3.5	3.7	3.9	4.0	4.1	4.1	4.2	4.4
	310A	BIO GASIF GAS	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3
Grand Total			544.4	570.8	569.2	628.8	569.3	532.6	618.1	558.9	531.1	524.6

Continued

Sum of Fuel_rate_PJ			Year			
fuel_type	fuel_id	fuel_gr_abbr	2010	2011	2012	2013
SOLID	101A	ANODIC CARBON	0.0	0.0	0.0	0.0
	102A	COAL	163.0	135.5	106.2	135.0
	103A	FLY ASH (COAL)		0.0	0.1	0.1
	106A	BROWN COAL BRI.	0.0	0.0	0.0	0.0
	107A	COKE OVEN COKE	0.7	0.7	0.6	0.6
LIQUID	110A	PETROLEUM COKE	5.1	6.5	6.7	6.1
	203A	RESIDUAL OIL	13.0	8.0	7.3	5.7
	204A	GAS OIL	23.2	16.9	13.0	10.6
	206A	KEROSENE	0.1	0.0	0.0	0.0
	225A	ORIMULSION				
	303A	LPG	1.5	1.4	1.5	1.3
	308A	REFINERY GAS	14.3	15.0	15.6	15.4
GAS	301A	NATURAL GAS	184.9	156.5	146.4	138.0
WASTE	114A	WASTE	36.4	36.7	35.8	37.1
	115A	INDUSTR. WASTES	1.4	1.7	1.5	1.8
BIOMASS	111A	WOOD	81.3	78.8	83.6	82.5
	117A	STRAW	23.3	20.2	18.3	20.6
	215A	LIQUID BIOFUELS	2.0	0.8	1.1	0.9
	309A	BIOGAS	4.6	4.3	4.7	4.9
	310A	BIO GASIF GAS	0.2	0.3	0.4	0.4
Grand Total			555.0	483.4	442.7	460.9

Table 3A-2.2 Detailed fuel consumption data for stationary combustion plants, PJ. 1990 – 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3A-3 Default Lower Calorific Value (LCV) of fuels and fuel correspondance list

Table 3A-3.1 Time-series for calorific values of fuels (DEA 2014a).

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Crude Oil, Average	GJ pr tonne	42.40	42.40	42.40	42.70	42.70	42.70	42.70	43.00	43.00	42.40
Crude Oil, Golf	GJ pr tonne	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80
Crude Oil, North Sea	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	43.00	43.00	42.70
Refinery Feedstocks	GJ pr tonne	41.60	41.60	41.60	41.60	41.60	41.60	41.60	42.70	42.70	41.60
Refinery Gas	GJ pr tonne	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00
LPG	GJ pr tonne	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00
Naphtha (LVN)	GJ pr tonne	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50
Motor Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Aviation Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
JP4	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Other Kerosene	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
JP1	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Gas/Diesel Oil	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70
Fuel Oil	GJ pr tonne	40.40	40.40	40.40	40.40	40.40	40.40	40.70	40.65	40.65	40.40
Orimulsion	GJ pr tonne	27.60	27.60	27.60	27.60	27.60	28.13	28.02	27.72	27.84	27.60
Petroleum Coke	GJ pr tonne	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40
Waste Oil	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
White Spirit	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Bitumen	GJ pr tonne	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80
Lubricants	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
Natural Gas	GJ pr 1000 Nm <sup>3</sup>	39.00	39.00	39.00	39.30	39.30	39.30	39.30	39.60	39.90	40.00
Town Gas	GJ pr 1000 m <sup>3</sup>							17.00	17.00	17.00	17.00
Electricity Plant Coal	GJ pr tonne	25.30	25.40	25.80	25.20	24.50	24.50	24.70	24.96	25.00	25.00
Other Hard Coal	GJ pr tonne	26.10	26.50	26.50	26.50	26.50	26.50	26.50	26.50	26.50	26.50
Coke	GJ pr tonne	31.80	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30
Brown Coal Briquettes	GJ pr tonne	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30
Straw	GJ pr tonne	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50
Wood Chips	GJ pr Cubic metre	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80
Wood Chips	GJ pr m <sup>3</sup>	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30
Firewood, Hardwood	GJ pr m <sup>3</sup>	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40
Firewood, Conifer	GJ pr tonne	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60
Wood Pellets	GJ pr tonne	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50
Wood Waste	GJ pr Cubic metre	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70
Wood Waste	GJ pr 1000 m <sup>3</sup>	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20
Biogas	GJ pr tonne								23.00	23.00	23.00
Wastes	GJ pr tonne	8.20	8.20	9.00	9.40	9.40	10.00	10.50	10.50	10.50	10.50
Bioethanol	GJ pr tonne	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70
Liquid Biofuels	GJ pr tonne	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60
Bio Oil	GJ pr tonne	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20



Continued		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Crude Oil, Average	GJ pr tonne	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00
Crude Oil, Golf	GJ pr tonne	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80	41.80
Crude Oil, North Sea	GJ pr tonne	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00	43.00
Refinery Feedstocks	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70
Refinery Gas	GJ pr tonne	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00	52.00
LPG	GJ pr tonne	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00	46.00
Naphtha (LVN)	GJ pr tonne	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50	44.50
Motor Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Aviation Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
JP4	GJ pr tonne	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80	43.80
Other Kerosene	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
JP1	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Gas/Diesel Oil	GJ pr tonne	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70	42.70
Fuel Oil	GJ pr tonne	40.65	40.65	40.65	40.65	40.65	40.65	40.65	40.65	40.65	40.65
Orimulsion	GJ pr tonne	27.62	27.64	27.71	27.65	27.65	27.65	27.65	27.65	27.65	27.65
Petroleum Coke	GJ pr tonne	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40	31.40
Waste Oil	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
White Spirit	GJ pr tonne	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50	43.50
Bitumen	GJ pr tonne	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80	39.80
Lubricants	GJ pr tonne	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90	41.90
Natural Gas	GJ pr 1000 Nm <sup>3</sup>	40.15	39.99	40.06	39.94	39.77	39.67	39.54	39.59	39.48	39.46
Town Gas	GJ pr 1000 m <sup>3</sup>	17.01	16.88	17.39	16.88	17.58	17.51	17.20	17.14	15.50	21.29
Electricity Plant Coal	GJ pr tonne	24.80	24.90	25.15	24.73	24.60	24.40	24.80	24.40	24.30	24.60
Other Hard Coal	GJ pr tonne	26.50	26.50	26.50	26.50	26.50	26.50	26.50	26.50	25.81	25.13
Coke	GJ pr tonne	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30	29.30
Brown Coal Briquettes	GJ pr tonne	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30	18.30
Straw	GJ pr tonne	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50	14.50
Wood Chips	GJ pr Cubic metre	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80	2.80
Wood Chips	GJ pr m <sup>3</sup>	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30	9.30
Firewood, Hardwood	GJ pr m <sup>3</sup>	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40	10.40
Firewood, Conifer	GJ pr tonne	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60	7.60
Wood Pellets	GJ pr tonne	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50	17.50
Wood Waste	GJ pr Cubic metre	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70	14.70
Wood Waste	GJ pr 1000 m <sup>3</sup>	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20	3.20
Biogas	GJ pr tonne	23.00	23.00	23.00	23.00	23.00	23.00	23.00	23.00	23.00	23.00
Wastes	GJ pr tonne	10.50	10.50	10.50	10.50	10.50	10.50	10.50	10.50	10.50	10.50
Bioethanol	GJ pr tonne	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70	26.70
Liquid Biofuels	GJ pr tonne	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.60	37.50	37.50
Bio Oil	GJ pr tonne	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20	37.20

Continued		2010	2011	2012	2013
Crude Oil, Average	GJ pr tonne	43.00	43.00	43.00	43.00
Crude Oil, Golf	GJ pr tonne	41.80	41.80	41.80	41.80
Crude Oil, North Sea	GJ pr tonne	43.00	43.00	43.00	43.00
Refinery Feedstocks	GJ pr tonne	42.70	42.70	42.70	42.70
Refinery Gas	GJ pr tonne	52.00	52.00	52.00	52.00
LPG	GJ pr tonne	46.00	46.00	46.00	46.00
Naphtha (LVN)	GJ pr tonne	44.50	44.50	44.50	44.50
Motor Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80
Aviation Gasoline	GJ pr tonne	43.80	43.80	43.80	43.80
JP4	GJ pr tonne	43.80	43.80	43.80	43.80
Other Kerosene	GJ pr tonne	43.50	43.50	43.50	43.50
JP1	GJ pr tonne	43.50	43.50	43.50	43.50
Gas/Diesel Oil	GJ pr tonne	42.70	42.70	42.70	42.70
Fuel Oil	GJ pr tonne	40.65	40.65	40.65	40.65
Orimulsion	GJ pr tonne	27.65	27.65	27.65	27.65
Petroleum Coke	GJ pr tonne	31.40	31.40	31.40	31.40
Waste Oil	GJ pr tonne	41.90	41.90	41.90	41.90
White Spirit	GJ pr tonne	43.50	43.50	43.50	43.50
Bitumen	GJ pr tonne	39.80	39.80	39.80	39.80
Lubricants	GJ pr tonne	41.90	41.90	41.90	41.90
Natural Gas	GJ pr 1000 Nm3	39.46	39.51	39.55	38.99
Town Gas	GJ pr 1000 m3	21.35	21.37	19.30	18.72
Electricity Plant Coal	GJ pr tonne	24.44	24.38	24.23	24.49
Other Hard Coal	GJ pr tonne	24.44	24.38	24.23	24.49
Coke	GJ pr tonne	29.30	29.30	29.30	29.30
Brown Coal Briquettes	GJ pr tonne	18.30	18.30	18.30	18.30
Straw	GJ pr tonne	14.50	14.50	14.50	14.50
Wood Chips	GJ pr Cubic metre	2.80	2.80	2.80	2.80
Wood Chips	GJ pr m3	9.30	9.30	9.30	9.30
Firewood, Hardwood	GJ pr m3	10.40	10.40	10.40	10.40
Firewood, Conifer	GJ pr tonne	7.60	7.60	7.60	7.60
Wood Pellets	GJ pr tonne	17.50	17.50	17.50	17.50
Wood Waste	GJ pr Cubic metre	14.70	14.70	14.70	14.70
Wood Waste	GJ pr 1000 m3	3.20	3.20	3.20	3.20
Biogas	GJ pr tonne	23.00	23.00	23.00	23.00
Wastes	GJ pr tonne	10.50	10.50	10.50	10.60
Bioethanol	GJ pr tonne	26.70	26.70	26.70	26.70
Liquid Biofuels	GJ pr tonne	37.50	37.50	37.50	37.50
Bio Oil	GJ pr tonne	37.20	37.20	37.20	37.20

Table 3A-3.2 Fuel category correspondence list, DEA, DCE and NFR.

<b>Danish Energy Agency</b>	<b>DCE Emission data-</b>	<b>IPCC fuel category base</b>
Other Hard Coal	Coal	Solid
Coke	Coke oven coke	Solid
Electricity Plant Coal	Coal	Solid
Brown Coal Briquettes	Brown coal briq.	Solid
-	Anode carbon	Solid
-	Fly ash	Solid
Orimulsion	Orimulsion	Liquid
Petroleum Coke	Petroleum coke	Liquid
Fuel Oil	Residual oil	Liquid
Waste Oil	Residual oil	Liquid
Gas/Diesel Oil	Gas oil	Liquid
Other Kerosene	Kerosene	Liquid
LPG	LPG	Liquid
Refinery Gas	Refinery gas	Liquid
Town Gas	Natural gas	Gas
Natural Gas	Natural gas	Gas
Straw	Straw	Biomass
Wood Waste	Wood and simil.	Biomass
Wood Pellets	Wood and simil.	Biomass
Wood Chips	Wood and simil.	Biomass
Firewood, Hardwood & Conifer	Wood and simil.	Biomass
Waste Combustion (biomass)	Municip. wastes	Biomass
Bio fuels	Liquid biofuels	Biomass
Biogas	Biogas	Biomass
Biogas, other	Biogas	Biomass
Biogas, landfill	Biogas	Biomass
Biogas, sewage sludge	Biogas	Biomass
(Wood applied in gas engines)	Biomass producer gas	Biomass
Waste Combustion (fossil)	Fossil waste	Other fuel

## Annex 3A-4 Emission factors

SO<sub>2</sub> emission factors are discussed in Nielsen et al. (2014). The emission factors applied for 2013 are shown in Table 3A-4.1.

Table 3A-4.1 SO<sub>2</sub> emission factors and references, 2013

Fuel type	Fuel	NFR	NFR_name	SNAP	SO <sub>2</sub> emission factor, g/GJ	Reference
SOLID	ANODE CARBON	1A2g	Industry - other	032002	574	Assumed equal to coal. DCE assumption.
	COAL	1A1a	Public electricity and heat production	0101	12	DCE estimate based on data reported by plant owners to the electricity transmission company, Energinet.dk (Energinet.dk, 2014)
				0102	574	DCE calculation based on DEPA (2010c), DEA (2012) and EMEP (2006)
		1A2a-g	Industry	03	574	DCE calculation based on DEPA (2010c), DEA (2012) and EMEP (2006)
		1A4b i	Residential	020200	574	DCE calculation based on DEPA (2010c), DEA (2012) and EMEP (2006)
		1A4c i	Agriculture/ Forestry	0203	574	DCE calculation based on DEPA (2010c), DEA (2012) and EMEP (2006)
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	010104	10	Assumed equal to the emission factor for coal in 2010. DCE assumption.
	BROWN COAL BRI.	1A4b	Residential	0202	574	Assumed equal to coal. DCE assumption.
	COKE OVEN COKE	1A2a-g	Industry	03	574	Assumed equal to coal. DCE assumption.
	LIQUID	PETROLEUM COKE	1A2a-g	Industry	03	605
RESIDUAL OIL		1A1a	Electricity and heat production	0101	100	DCE estimate based on plant specific data for 2008 and 2009.
				0102	344	DCE estimate based on EOF (2013) and DEA (2012)
		1A1b	Petroleum refining	010306	537	DCE calculation based on plant specific data for year 2003.
		1A2a-g	Industry	03	344	DCE estimate based on EOF (2013) and DEA (2012)
		1A4a	Commercial/ Institutional	0201	344	DCE estimate based on EOF (2013) and DEA (2012)
		1A4b	Residential	0202	344	DCE estimate based on EOF (2013) and DEA (2012)
		1A4c i	Agriculture/ Forestry	0203	344	DCE estimate based on EOF (2013) and DEA (2012)
		GAS OIL	1A1a	Public electricity and heat production	0101	23
0102					23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
1A1b			Petroleum refining	010306	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
1A1c			Other energy industries	0105	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
1A2a-g			Industry	03	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
1A4a			Commercial/ Institutional	0201	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
1A4b i			Residential	0202	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
1A4c			Agriculture/Forestry	0203	23	DCE estimate based on DEPA (1998), Miljø- og planlægningsudvalget (1998) and DEA (2012).
KEROSENE		1A2g	Industry - other	03	5	DCE estimate based on Tønder (2004) and Shell (2013).

Fuel type	Fuel	NFR	NFR_name	SNAP	SO <sub>2</sub> emission factor, g/GJ	Reference
		1A4a	Commercial/ Institutional	0201	5	DCE estimate based on Tønder (2004) and Shell (2013).
		1A4b i	Residential	0202	5	DCE estimate based on Tønder (2004) and Shell (2013).
		1A4c i	Agriculture/ Forestry	0203	5	DCE estimate based on Tønder (2004) and Shell (2013).
	LPG	1A1a	Public electricity and heat production	All	0.13	DCE estimate based on Augustesen (2003) and DEA (2012).
		1A2a-g	Industry	03	0.13	DCE estimate based on Augustesen (2003) and DEA (2012).
		1A4a	Commercial/ Institutional	0201	0.13	DCE estimate based on Augustesen (2003) and DEA (2012).
		1A4b i	Residential	0202	0.13	DCE estimate based on Augustesen (2003) and DEA (2012).
		1A4c i	Agriculture/ Forestry	0203	0.13	DCE estimate based on Augustesen (2003) and DEA (2012).
	REFINERY GAS	1A1b	Petroleum refining	0103	1	DCE estimate based on plant specific data for one plant, average value for 1995-2002.
GAS	NATURAL GAS	1A1a	Public electricity and heat production	0101, 0102, except engines	0.43	DCE estimate based on data from Energinet.dk (2013)
				010105, engines	0.5	Kristensen (2003)
		1A1c	Other energy industries	0105	0.43	DCE estimate based on data from Energinet.dk (2013)
		1A2a-g	Industry	03 ex- cept engines	0.43	DCE estimate based on data from Energinet.dk (2013)
				Engines	0.5	Kristensen (2003)
		1A4a	Commercial/ Institutional	0201 except engines	0.43	DCE estimate based on data from Energinet.dk (2013)
				Engines	0.5	Kristensen (2003)
		1A4b i	Residential	0202 except engines	0.43	DCE estimate based on data from Energinet.dk (2013)
				Engines	0.5	Kristensen (2003)
		1A4c i	Agriculture/ Forestry	0203 except engines	0.43	DCE estimate based on data from Energinet.dk (2013)
				Engines	0.5	Kristensen (2003)
WASTE	WASTE	1A1a	Public electricity and heat production	0101 0102	8.3	Nielsen et al. (2010a)
					14	DCE estimate based on plant specific data for four plants, 2009 data.
		1A2a-g	Industry	03	14	Assumed equal to district heating plants (DCE assumption).
		1A4a	Commercial/ Institutional	0201	14	Assumed equal to district heating plants (DCE assumption).
	INDU-STRIAL WASTE	1A2g	Industry - Other	031600	14	Assumed equal to waste. DCE assumption.
BIO- MASS	WOOD	1A1a	Public electricity and heat production	0101 0102	1.9	Nielsen et al. (2010a)
					11	EEA (2013)
		1A2a-g	Industry	03	11	EEA (2013)
		1A4a	Commercial/ Institutional	0201	11	EEA (2013)

Fuel type	Fuel	NFR	NFR_name	SNAP	SO <sub>2</sub> emission factor, g/GJ	Reference
		1A4b i	Residential	0202	11	EEA (2013)
		1A4c i	Agriculture/ Forestry	0203	11	EEA (2013)
	STRAW	1A1a	Public electricity and heat production	0101	49	Nielsen et al. (2010a)
				0102	130	Nikolaisen et al. (1998)
		1A4b i	Residential	0202	130	Assumed equal to district heating plants. DCE assumption.
		1A4c i	Agriculture/ Forestry	0203	130	Assumed equal to district heating plants. DCE assumption.
	BIO OIL	1A1a	Public electricity and heat production	0101	0.1	DCE estimate based on Folkecenter for Vedvarende Energi (2000) and DEA (2012).
		1A2a-g	Industry	03	0.1	DCE estimate based on Folkecenter for Vedvarende Energi (2000) and DEA (2012).
		1A4b i	Residential	0202	0.1	DCE estimate based on Folkecenter for Vedvarende Energi (2000) and DEA (2012).
	BIOGAS	1A1a	Public electricity and heat production	0101, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012).
				Engines	19.2	Nielsen & Illerup (2003)
				0102	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012).
		1A2a-g	Industry	03, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012).
				03, engines	19.2	Nielsen & Illerup (2003)
		1A4a	Commercial/ Institutional	0201, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012).
				020105	19.2	Nielsen & Illerup (2003)
		1A4c i	Agriculture/ Forestry	0203, except engines	25	DCE estimate based on Christiansen (2003), Hjort-Gregersen (1999) and DEA (2012).
				020304	19.2	Nielsen & Illerup (2003)
	BIO GASIF GAS	1A1a	Public electricity and heat production	010105	1.9	Assumed equal to wood. DCE assumption.

NO<sub>x</sub> emission factors are discussed in Nielsen et al. (2014). The emission factors applied for 2013 are shown in Table 3A-4.2.

Table 3A-4.2 NO<sub>x</sub> emission factors and references, 2013

Fuel type	Fuel	NFR	NFR_name	SNAP	NO <sub>x</sub> emission factor, g/GJ	Reference		
SOLID	ANODE CARBON	1A2g	Industry - other	032000	132	Assumed equal to coal. DCE assumption.		
	COAL	1A1a	Public electricity and heat production	0101	29	DCE estimate based on Energinet.dk (2014) and EU ETS (2014)		
				0102	95	DEPA (2001)		
		1A2a-g	Industry	03	132	DCE estimate based on plant specific data for 2011.		
				except cement production				
		1A2f	Industry, cement production	0316	153	DCE estimate based on plant specific data for 2013.		
		1A4b i	Residential	020200	95	DEPA (2001)		
		1A4c i	Agriculture/ Forestry	0203	95	DEPA (2001)		
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	0101	30	Assumed equal to the emission factor for coal in 2010.		
	BROWN COAL BRI.	1A4b	Residential	0202	95	Assumed equal to coal. DCE assumption.		
COKE OVEN COKE	1A2a-g	Industry	03	132	Assumed equal to coal. DCE assumption.			
LIQUID	PETROLEUM COKE	1A2a-g	Industry	03	129	Assumed equal to residual oil. DCE assumption.		
	RESIDUAL OIL	1A1a	Public electricity and heat production	0101	138	DCE estimate based on Energinet.dk (2009); Energinet.dk (2010); Energinet.dk (2011); EU ETS (2009-2011)		
				0102	142	DEPA (2001)		
		1A1b	Petroleum refining	010306	200	IPCC (1997)		
		1A2a-g	Industry	03	129	DCE estimate based on plant specific data from two plants, 2011		
		1A4a	Commercial/ Institutional	0201	142	DEPA (2001)		
		1A4b	Residential	0202	142	DEPA (2001)		
		1A4c i	Agriculture/ Forestry	0203	142	DEPA (2001)		
	GAS OIL	1A1a	Public electricity and heat production	010101, 010102	114	DCE estimate based on plant specific data for 2011. Data from Energinet.dk (2011) and EU ETS (2011).		
				010103	130	DEPA (2012b), DEPA (2003b) and DEPA (1990)		
				0102				
				010104	350	DCE estimate based on Eltra & Elkraft System, (2001) and DEA (2012b)		
				010105	942	Nielsen et al. (2010a)		
				1A1b	Petroleum refining	010306	65	DEPA (1990)
				1A1c	Other energy industries	010504	350	Assumed equal to gas turbines applied in CHP plants. DCE assumption.
				1A2a-g	Industry	03	130	DEPA (2012b), DEPA (2003b) and DEPA (1990)
					except engines			
				1A2a-g	Industry	Engines	942	Nielsen et al. (2010a)
	1A4a	Commercial/ Institutional	0201	52	DEPA (2001)			
			020105	942	Nielsen et al. (2010a)			
	1A4b i	Residential	0202	52	DEPA (2001a)			

Fuel type	Fuel	NFR	NFR_name	SNAP	NO <sub>x</sub> emission Reference factor, g/GJ
				Engines	942 Nielsen et al. (2010a)
		1A4c	Agriculture/Forestry	0203	52 DEPA (2001a)
				Engines	942 Nielsen et al. (2010a)
	KEROSENE	1A2g	Industry - other	03	50 EEA (2009)
		1A4a	Commercial/ Institutional	0201	50 EEA (2009)
		1A4b i	Residential	0202	50 EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	50 EEA (2009)
	LPG	1A1a	Public electricity and heat production	All	96 IPCC (1997)
		1A2a-g	Industry	03	96 IPCC (1997)
		1A4a	Commercial/ Institutional	0201	71 IPCC (1997)
		1A4b i	Residential	0202	47 IPCC (1997)
		1A4c i	Agriculture/ Forestry	0203	71 IPCC (1997)
	REFINERY GAS	1A1b	Petroleum refining	010304	170 DCE estimate based on plant specific data for a gas turbine in year 2000.
				010306	94 DCE estimate based on plant specific data for year 2011.
GAS	NATURAL GAS	1A1a	Public electricity and heat production	010101, 010102	55 DEPA (2003b)
				010103	42 Larsen (2009)
				010104	48 Nielsen et al. (2010a)
				010105	135 Nielsen et al. (2010a)
				0102	42 Larsen (2009)
		1A1c	Other energy industries	010504	250 Kristensen (2004)
		1A2a-g	Industry	03	42 Larsen (2009)
				Engines	135 Nielsen et al. (2010a)
				Turbines	48 Nielsen et al. (2010a)
		1A2f		030700	87 DCE estimate based on plant specific data for 11 clay production plants, EU ETS (2011-2012); DEPA (2012)
		1A4a	Commercial/ Institutional	0201	30 Larsen (2009); DEPA (2001a)
				Engines	135 Nielsen et al. (2010a)
		1A4b i	Residential	0202	26.9 Schweitzer (2014)
				Engines	135 Nielsen et al. (2010a)
		1A4c i	Agriculture/ Forestry	0203	30 Larsen (2009); DEPA (2001a)
				Engines	135 Nielsen et al. (2010a)
WASTE	WASTE	1A1a	Public electricity and heat production	0101	102 Nielsen et al. (2010a)
				0102	164 DCE estimate based on plant specific data for year 2000.
		1A2a-g	Industry	03	164 DCE estimate based on plant specific data for district heating plants in year 2000.
		1A4a	Commercial/ Institutional	0201	164 DCE estimate based on plant specific data for district heating plants in year 2000.
	INDUSTRIAL WASTE	1A2f	Industry - Other	031600	164 Assumed equal to waste. DCE assumption.
BIO-MASS	WOOD	1A1a	Public electricity and heat production	0101	81 Nielsen et al. (2010a)
				0102	90 Serup et al. (1999)
		1A2a-g	Industry	03	90 Serup et al. (1999)



Fuel type	Fuel	NFR	NFR_name	SNAP	NO <sub>x</sub> emission Reference factor, g/GJ
		1A4a	Commercial/ Institutional	0201	90 Serup et al. (1999)
		1A4b i	Residential	0202	74.95 DCE estimate based on DEA (2014a), DEPA (2013) and EEA (2013),
		1A4c i	Agriculture/ Forestry	0203	90 Serup et al. (1999)
	STRAW	1A1a	Public electricity and heat production	0101	125 Nielsen et al. (2010a)
				0102	90 Nikolaisen et al. (1998)
		1A4b i	Residential	0202	90 Assumed equal to district heating plants. DCE assumption.
		1A4c i	Agriculture/ Forestry	0203	90 Assumed equal to district heating plants. DCE assumption.
	BIO OIL	1A1a	Public electricity and heat production	0101	249 Assumed equal to gas oil. DCE assumption. The emission factor for gas oil have been changed and the emission factor for biooil will also be changed in future inventories.
				0102	65 Assumed equal to gas oil. DCE assumption.
		1A2a-g	Industry	03	65 Assumed equal to gas oil. DCE assumption.
				Engines	700 Assumed equal to gas oil. DCE assumption.
		1A4b i	Residential	0202	65 Assumed equal to gas oil. DCE assumption.
	BIOGAS	1A1a	Public electricity and heat production	0101, not engines	28 DEPA (2001a)
				Engines	202 Nielsen et al. (2010a)
				0102	28 DEPA (2001a)
		1A2a-g	Industry	03, not engines	28 DEPA (2001a)
				03, engines	202 Nielsen et al. (2010a)
				030902	59 DEPA (1990); DEPA (1995)
		1A4a	Commercial/ Institutional	0201, not engines	28 DEPA (2001a)
				020105	202 Nielsen et al. (2010a)
		1A4c i	Agriculture/ Forestry	0203, not engines	28 DEPA (2001a)
				020304	202 Nielsen et al. (2010a)
	BIO GASIF GAS	1A1a	Public electricity and heat production	010105	173 Nielsen et al. (2010a)

Table 3A-4.3 NMVOC emission factors and references, 2013.

Fuel type	Fuel	NFR	NFR_name	SNAP	NMVOC, Reference g/GJ										
SOLID	ANODE CARBON	1A2g	Industry - other	0320	10 Assumed equal to coal. DCE assumption.										
	COAL	1A1a	Public electricity and heat production	0101 0102	1.2 EEA (2009)										
		1A2a-g	Industry	03	10 EEA (2009)										
		1A4c i	Agriculture/ Forestry	0203	88.8 EEA (2009)										
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	0101	1.2 Assumed equal to coal. DCE assumption.										
	BROWN COAL BRI.	1A4b i	Residential	0202	484 EEA (2009)										
	COKE OVEN COKE	1A2a-g	Industry	03	10 EEA (2009)										
LIQUID	PETROLEUM COKE	1A2a-g	Industry	03	10 Assumed equal to coal. DCE assumption.										
	RESIDUAL OIL	1A1a	Public electricity and heat production	010101 010102 010103 010104 010105 010203	0.8 Nielsen et al. (2010)										
				1A1b	Petroleum refining	010306	2.3 EEA (2009)								
				1A2a-g	Industry	03 except engines Engines	0.8 Nielsen et al. (2010) 10 EEA (2009)								
				1A4a	Commercial/ Institutional	0201	5 EEA (2009)								
				1A4b	Residential	0202	15 EEA (2009)								
				1A4c i	Agriculture/ Forestry	0203	5 EEA (2009)								
				GAS OIL	1A1a	Public electricity and heat production	010101 010102 010103 010104 010105 0102 010306 010504	0.8 EEA (2009)							
							1A1b	Petroleum refining	010306	0.2 EEA (2009)					
							1A1c	Other energy industries	010504	37 EEA (2009)					
							1A2a-g	Industry	03 boilers > 50 MW 03 boilers < 50 MW Gas turbines Engines	5 EEA (2009) 10 EEA (2009) 0.2 EEA (2009) 37 EEA (2009)					
									1A4a	Commercial/ Institutional	0201 except engines Engines	5 EEA (2009) 37 EEA (2009)			
									1A4b i	Residential	0202	15 EEA (2009)			
							1A4c	Agriculture/Forestry	020302	5 EEA (2009)					
							KEROSENE	1A2a-g	Industry	03 0201 0202 0203	10 EEA (2009) 5 EEA (2009) 15 EEA (2009) 5 EEA (2009)				
										1A4a	Commercial/ Institutional	0201 0202 0203	5 EEA (2009) 15 EEA (2009) 5 EEA (2009)		
												1A4b i	Residential	0201 0202	5 EEA (2009) 15 EEA (2009)
														1A4c i	Agriculture/ Forestry
							LPG	1A1a	Public electricity and heat production	0101 0102	0.8 EEA (2009)				
										1A2a-g	Iron and steel	03	5 EEA (2009)		

Fuel type	Fuel	NFR	NFR_name	SNAP	NMVOC, Reference g/GJ
		1A4a	Commercial/ Institutional	0201	5 EEA (2009)
		1A4b i	Residential	0202	10 EEA (2009)
		1A4c i	Agriculture/ Forestry	0203	5 EEA (2009)
	REFINERY GAS	1A1b	Petroleum refining	0103	1.4 Assumed equal to natural gas fuelled gas turbines. DCE assumption.
GAS	NATURAL GAS	1A1a	Public electricity and heat production	010101 010102 010103 010104 010105 0102	2 Danish Gas Technology Centre (2001).   1.6 Nielsen et al. (2010) 92 Nielsen et al. (2010) 2 Danish Gas Technology Centre (2001).
		1A1c	Other energy industries	0105	1.6 Nielsen et al. (2010)
		1A2a-g	Industry	03 except engines and turbines Turbines Engines	2 Danish Gas Technology Centre (2001). 1.6 Nielsen et al. (2010) 92 Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines Engines	2 Danish Gas Technology Centre (2001). 92 Nielsen et al. (2010)
		1A4b i	Residential	0202 except engines Engines	4 Gruijthuijsen & Jensen (2000) 92 Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines Engines	2 Danish Gas Technology Centre (2001). 92 Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101 0102	0.56 Nielsen et al. (2010) 2 EEA (2009)
		1A2a-g	Industry	03	2 EEA (2009)
		1A4a	Commercial/ Institutional	0201	2 EEA (2009)
	INDISTRIAL WASTE	1A2f	Industry	0316	2 EEA (2009)
BIO- MASS	WOOD	1A1a	Public electricity and heat production	0101  0102	5.1 Nielsen et al. (2010)  7.3 EEA (2009)
		1A2a-g	Industry	03	10 EEA (2009)
		1A4a	Commercial/ Institutional	0201	146 EEA (2009)
		1A4b i	Residential	0202	338.2 DCE estimate based on DEA (2014a), DEPA (2013) and EEA (2013),
		1A4c i	Agriculture/ Forestry	0203	146 EEA (2009)
	STRAW	1A1a	Public electricity and heat production	0101 0102	0.78 Nielsen et al. (2010) 7.3 EEA (2009)
		1A4b i	Residential	0202	600 EEA (2009)
		1A4c i	Agriculture/ Forestry	0203 020302	600 EEA (2009). Plants are assued equal to residential plants. 10 EEA (2009)
	BIO OIL	1A1a	Public electricity and heat production	010102 010105 0102	0.8 EEA (2009) 37 EEA (2009) 0.8 EEA (2009)
		1A2a-g	Industry	03, not engines	0.8 EEA (2009)

Fuel type	Fuel	NFR	NFR_name	SNAP	NMVOC, Reference g/GJ
				Engines	37 EEA (2009)
		1A4b i	Residential	0202	15 EEA (2009)
	BIOGAS	1A1a	Public electricity and heat production	0101	2 Assumed equal to natural gas. DCE assumption.
				010105	10 Nielsen et al. (2010)
				0102	2 Assumed equal to natural gas. DCE assumption.
		1A2a-g	Industry	03 except engines	2 Assumed equal to natural gas. DCE assumption.
				Engines	10 Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	2 Assumed equal to natural gas. DCE assumption.
				Engines	10 Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines	2 Assumed equal to natural gas. DCE assumption.
				Engines	10 Nielsen et al. (2010)
	BIO GASIF GAS	1A1a	Public electricity and heat production	010105	2 Nielsen et al. (2010)

Table 3A-4.4 CO emission factors and references, 2013.

Fuel type	Fuel	NFR	NFR_name	SNAP	CO emis- sion factor g/GJ	Reference		
SOLID	ANODE CARBON	1A2a-g	Industry	03	10	Assumed the same emission factor as for coal. DCE assumption.		
	COAL	1A1a	Public electricity and heat production	0101 and 0102	10	Sander (2002)		
		1A2a-g	Industry	03	10	Assumed equal to boilers in public electricity and heat production. DCE assumption.		
		1A4c i	Agriculture/ Forestry	0203	931	EEA (2009)		
	FLY ASH FOSSIL	1A1a	Public electricity and heat production	0101	10	Assumed equal to coal. DCE assumption.		
	BROWN COAL BRI.	1A4b i	Residential	0202	2000	EEA (2009)		
	COKE OVEN COKE	1A2a-g	Industry	03	10	Assumed the same emission factor as for coal. DCE assumption.		
LIQUID	PETROLEUM COKE	1A2a-g	Industry	03	61	EEA (2013)		
	RESIDUAL OIL	1A1a	Electricity and heat production	010101	15	Sander (2002)		
				010104				
				010105				
				010102	2.8 -			
				010103				
			0102	30	EEA (2007)			
		1A1b	Petroleum refining	010306	30	EEA (2007)		
		1A2a-g	Industry	03 except engines	2.8	Nielsen et al. (2010)		
				Engines	100	EEA (2009)		
				1A4a	Commercial/Institutional	0201	30	EEA (2007)
				1A4b	Residential	0202	30	EEA (2007)
				1A4c i	Agriculture/ Forestry	0203	30	EEA (2007)
		GAS OIL	1A1a	Public electricity and heat production	0101 except engines	15	Sander (2002)	
					Engines	130	Nielsen et al. (2010)	
					0102	30	EEA (2007)	
	1A1b				Petroleum refining	010306	30	EEA (2007)
	1A2a-g				Industry	03 except gas turbines and engines	30	EEA (2007)
						Gas turbines	15	Sander (2002)
			Engines	130	Nielsen et al. (2010)			
	1A4a		Commercial/ Institutional	0201 except engines	30	EEA (2007)		
			Engines	130	Nielsen et al. (2010)			
	1A4b i		Residential	0202	43	EEA (2007)		
	1A4c		Agriculture/Forestry	0203	30	EEA (2007)		
	KEROSENE		1A2a-g	Industry	03	20	EEA (2007)	
		1A4a			Commercial/ Institutional	0201	20	EEA (2007)
		1A4b i			Residential	0202	20	EEA (2007)
1A4c i		Agriculture/ Forestry			0203	20	EEA (2007)	
LPG	1A1a	Public electricity and heat production	0101 and 0102	25	EEA (2007)			
			1A2a-g	Industry	03	25	EEA (2007)	
			1A4a	Commercial/ Institutional	0201	25	EEA (2007)	
			1A4b i	Residential	0202	25	EEA (2007)	

Fuel type	Fuel	NFR	NFR_name	SNAP	CO emis- sion factor g/GJ	Reference
		1A4c i	Agriculture/ Forestry	0203	25	EEA (2007)
	REFINERY GAS	1A1b	Petroleum refining	0103	6.2	Assumed same emission factor as for natural gas fuelled gas turbines. DCE assumption.
GAS	NATURAL GAS	1A1a	Public electricity and heat production	010101 and 010102	15	Sander (2002)
				010103	28	DEPA (2001)
				010104	4.8	Nielsen et al. (2010)
				010105	58	Nielsen et al. (2010)
				0102	28	DEPA (2001)
		1A1c	Other energy industries	0105	4.8	Nielsen et al. (2010)
		1A2a-g	Industry	03 except gas tur- bines and engines	28	DEPA (2001)
				Gas turbines	4.8	Nielsen et al. (2010)
				Engines	58	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	28	DEPA (2001)
				Engines	58	Nielsen et al. (2010)
		1A4b i	Residential	0202 except engines	20	Gruijthuijsen & Jensen (2000)
				Engines	58	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines	28	DEPA (2001)
				Engines	58	Nielsen et al. (2010)
WASTE	WASTE	1A1a	Public electricity and heat production	0101	3.9	Nielsen et al. (2010)
				0102	10	DCE calculation based on annual environmental reports for Danish plants year 2000.
		1A2a-g	Industry	03	10	Assumed equal to district heating plants. DCE as- sumption.
		1A4a	Commercial/ Institutional	0201	10	Assumed equal to district heating plants. DCE as- sumption.
	INDISTRIAL WASTE	1A2f	Industry	0316	10	Assumed equal to waste, district heating plants. DCE assumption.
BIO- MASS	WOOD	1A1a	Public electricity and heat production	0101	90	Nielsen et al. (2010)
				010203	240	DEPA (2001)
		1A2a-g	Industry	03	240	DEPA (2001)
		1A4a	Commercial/ Institutional	020100	240	DEPA (2001)
		1A4b i	Residential	0202	2491	DCE estimate based on DEA (2014a), DEPA (2013) and EEA (2013),
		1A4c i	Agriculture/ Forestry	020300	240	DEPA (2001)
	STRAW	1A1a	Public electricity and heat production	0101	67	Nielsen et al. (2010)
				0102	325	DEPA (2001); Nikolaisen et al (1998)
		1A4b i	Residential	0202	4000	EEA (2007); Jensen & Nielsen (1990) and Bjerrum (2002)
		1A4c i	Agriculture/ Forestry	0203	4000	EEA (2007); Jensen & Nielsen (1990) and Bjerrum (2002)
				020302	325	DEPA (2001); Nikolaisen et al (1998)
	BIO OIL	1A1a	Public electricity and heat production	0101 and 0102	15	Assumed same emission factor as for gas oil. DCE

Fuel type	Fuel	NFR	NFR_name	SNAP	CO emis- sion factor g/GJ	Reference
		1A2a-g				assumption.
		1A4b i	Residential	0202	15	Assumed same emission factor as for gas oil. DCE assumption.
	BIOGAS	1A1a	Public electricity and heat production	0101 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
				0102	36	DEPA (2001)
		1A2a-g	Industry	03 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
		1A4a	Commercial/ Institutional	0201 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
		1A4c i	Agriculture/ Forestry	0203 except engines	36	DEPA (2001)
				Engines	310	Nielsen et al. (2010)
	BIO GASIF GAS	1A1a	Public electricity and heat production	010105	586	Nielsen et al. (2010)

Table 3A-4.5 SO<sub>2</sub> emission factors time series, g per GJ for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.6 NO<sub>x</sub> emission factors time series, g per GJ for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.7 NMVOC emission factors time series, g per GJ for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.8 CO emission factors time series, g per GJ for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>



Table 3A-4.9 PM emission factors (in g per GJ) and references, 2013.

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	EA	snap_id	TSP	Ref	PM <sub>10</sub>	Ref	PM <sub>2.5</sub>	Ref					
SOLID	101A	ANODIC CARBON	1A2g	iii	0320	17	6	12	14	7	14					
	102A	COAL	1A1a		0101	3	12	2.6	12	2.1	12					
					0102	6	10	6	10	5	10					
					03	17	6	12	14	7	14					
					1A4c i	17	6	12	14	7	14					
	103A	FLY ASH FOSSIL	1A1a		0101	3	12	2.6	12	2.1	12					
	106A	BROWN COAL BRI.	1A4b	i	0202	17	16	12	16	7	16					
107A	COKE OVEN COKE	1A2	a-g	03	17	16	12	16	7	16						
LIQUID	110A	PETROLEUM COKE	1A2a-g		03	10	9	7	9	3	9					
	203A	RESIDUAL OIL	1A1a		010101	3	3	3	3	2.5	3					
					010102	9.5	18	9.5	13	7.9	13					
					010103	9.5	18	9.5	13	7.9	13					
					010104	3	9	3	9	2.5	9					
					010105	3	9	3	9	2.5	9					
					0102	3	9	3	9	2.5	9					
					1A1b	010306	50	9	40	9	35	9				
					1A2	a-g	03	9.5	18	7.1	13	4.8	13			
					1A4a	0201	14	6	10.5	13	7	13				
					1A4b	0202	14	6	10.5	13	7	13				
	1A4c i	0203	14	6	10.5	13	7	13								
	204A	GAS OIL	1A1a		0101	5	9	5	9	5	9					
					0102	5	9	5	9	5	9					
					010306	5	9	5	9	5	9					
					1A2a-g	03	5	9	5	9	5	9				
					1A4a i	0201	5	9	5	9	5	9				
					1A4b i	0202	5	9	5	9	5	9				
					1A4c i	0203	5	9	5	9	5	9				
	206A	KEROSENE	1A2	a-g	all	5	9	5	9	5	9					
					1A4a i	0201	5	9	5	9	5	9				
					1A4b i	0202	5	9	5	9	5	9				
					1A4c i	0203	5	9	5	9	5	9				
	303A	LPG	1A1a		0101, 0102	0.2	9	0.2	9	0.2	9					
					1A2	a-g	03	0.2	9	0.2	9	0.2	9			
					1A4a i	0201	0.2	9	0.2	9	0.2	9				
					1A4b i	0202	0.2	9	0.2	9	0.2	9				
					1A4c i	0203	0.2	9	0.2	9	0.2	9				
	308A	REFINERY GAS	1A1b		0103	5	9	5	9	5	9					
GAS	301A	NATURAL GAS	1A1a		0101	0.1	9	0.1	9	0.1	9					
					Gas turbines	0.1	3	0.061	3	0.051	3					
					Engines	0.76	3	0.189	3	0.161	3					
					0102	0.1	9	0.1	9	0.1	9					
					1A1c	0105	0.1	3	0.061	3	0.051	3				
					1A2a-g	Engines	0.76	3	0.189	3	0.161	3				
						Turbines	0.1	3	0.061	3	0.051	3				
						Other	0.1	9	0.1	9	0.1	9				
					1A4a i	0201	0.1	9	0.1	9	0.1	9				
						Engines	0.76	3	0.189	3	0.161	3				
					1A4b i	0202	0.1	9	0.1	9	0.1	9				
						Engines	0.76	3	0.189	3	0.161	3				
					1A4c i	0203	0.1	9	0.1	9	0.1	9				
						Engines	0.76	3	0.189	3	0.161	3				
					WASTE	114A	WASTE	1A1a		0101	0.29	18	0.29	3	0.29	3
0102	0.29	20	0.29	3						0.29	3					
1A2	a-g	03	4.2	20						3.2	20	2.1	20			
1A4a i	0201	4.2	20	3.2						20	2.1	20				
115A	INDUSTRIAL WASTE	1A2f	0316	4.2						20	3.2	20	2.1	20		
BIOMASS	111A	WOOD	1A1a							0101	10	18	7.45	8	4.82	8
										0102	19	1	13	2	10	1
					1A2	a-g	03	19	1	13	2	10	1			
					1A4a i	0201	143	1	143	9	135	9				
					1A4b i	0202	441	17	419	17	409	17				
					1A4c i	0203	143	1	143	9	135	9				
117A	STRAW	1A1a	i	0101	2.3	18	1.71	3	1.11	3						
				0102	21	1	15	2	12	2						

fuel_type	fuel_id	fuel_gr_abbr	nfr_id EA	snap_id	TSP	Ref	PM <sub>10</sub>	Ref	PM <sub>2.5</sub>	Ref
			1A4b i	0202	234	4	222	5	211	5
			1A4c i	0203	234	4	222	5	211	5
				020302	21	1	15	2	12	2
215A	BIO OIL		1A1a	0101	5	15	5	15	5	15
				0102	5	15	5	15	5	15
				0102	5	15	5	15	5	15
			1A2a-g	03	5	15	5	15	5	15
			1A4b i	0202	5	15	5	15	5	15
309A	BIOGAS		1A1a	0101, not engines	1.5	6	1.5	7	1.5	7
				010105	2.63	3	0.451	3	0.206	3
				0102	1.5	6	1.5	7	1.5	7
			1A2a-g	Engines	2.63	3	0.451	3	0.206	3
				Other	1.5	6	1.5	7	1.5	7
			1A4a i	0201	1.5	6	1.5	7	1.5	7
				Engines	2.63	3	0.451	3	0.206	3
			1A4c i	0203	1.5	6	1.5	7	1.5	7
				Engines	2.63	3	0.451	3	0.206	3
310A	BIO GASIF GAS		1A1a	010105	2.63	19	0.451	19	0.206	19

- Danish legislation, Miljøstyrelsen 2001. Luftvejledningen, Begrænsning af luftforurening fra virksomheder, Vejledning fra Miljøstyrelsen nr 2 2001 (DEPA, 2001).
- Particulate size distribution for wood and straw combustion in power plants refers to the TNO CEPMEIP emission factor database 2001 (wood). Available at: <http://www.air.sk/tno/cepmeip/> (05-02-2015).
- Nielsen, M. & Illerup, J.B: 2003. Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme. Eltra PSO projekt 3141. Kortlægning af emissioner fra decentrale kraftvarmeverker. Delrapport 6. Danmarks Miljøundersøgelser. 116 s. –Faglig rapport fra DMU nr. 442.(In Danish, with an English summary). Available at: [http://www.dmu.dk/1\\_viden/2\\_Publikationer/3\\_fagrappporter/rappporter/FR442.pdf](http://www.dmu.dk/1_viden/2_Publikationer/3_fagrappporter/rappporter/FR442.pdf) (05-02-2015).
- German, L., 2003. The Danish Technological Institute, Personal communication, expert judgement, rough estimate.
- Particulate size distribution for wood and straw combustion in residential plants refers to the TNO CEPMEIP emission factor database 2001 (wood). Available at: <http://www.air.sk/tno/cepmeip/> (05-02-2015).
- Danish legislation. Miljøstyrelsen 1990, Bekendtgørelse 689, 15/10/1990, Bekendtgørelse om begrænsning af emissioner af svovldioxid, kvælstofoxider og støv fra store fyringsanlæg. (and Bekendtgørelse 518/1995). (DEPA, 1990)
- All TSP emission is assumed to be <2,5µm (DCE assumption).
- Estimated based on the TSP emission factor.
- The TNO CEPMEIP emission factor database 2001. Available on the internet at: <http://www.air.sk/tno/cepmeip/> (05-02-2015).
- TNO CEPMEIP emission factor database 2001. Available at: <http://www.air.sk/tno/cepmeip/> (05-02-2015).
- 
- Livbjerg, H. Thellefsen, M. Sander, B. Simonsen, P., Lund, C., Poulsen, K. & Fogh, C.L., 2001. Feltstudier af Forbrændingsaerosoler, EFP -98 Projekt, Aerosollaboratoriet DTU, FLS Miljø, Forskningscenter Risø, Elsam, Energi E2 (in Danish).
- Particulate size distribution for residual oil combustion refers to the TNO CEPMEIP emission factor database 2001. Available at: <http://www.air.sk/tno/cepmeip/> (05-02-2015).
- Particulate size distribution for coal combustion refers to the TNO CEPMEIP emission factor database 2001. Available at: <http://www.air.sk/tno/cepmeip/> (05-02-2015).
- Assuming same emission factors as for gas oil (DCE assumption).
- Same emission factor as for coal is assumed (DCE assumption).
- DCE estimate based on DEA (2014a), DEPA (2013), Glasius et al. (2005), EEA (2013), Illerup et al. (2007), Nordic Swan label (2012)
- Nielsen, M., Nielsen, O.K. & Thomsen, M. 2010: Emissionskortlægning for decentral kraftvarme, Energinet.dk miljøprojekt nr. 07/1882. Delrapport 5. Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme, 2006. National Environmental Research Institute, Aarhus University.
- Same emission factor as for biogas assumed (DCE assumption).
- The emission factor have been estimated by DCE based on plant specific data from MSW incineration plants, district heating, 2008.

Table 3A-4.10 TSP emission factors, time series for the years 2000 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.11 PM10 emission factors, time series for the years 2000 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.12 PM2.5 emission factors, time series for the years 2000 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.13 HM emission factors (mg per GJ) and references 2013.

fuel_type	fuel_gr_abbr	nfr	nfr_name	snap	As mg/GJ	Cd mg/GJ	Cr mg/GJ	Cu mg/GJ	Hg mg/GJ	Ni mg/GJ	Pb mg/GJ	Se mg/GJ	Zn mg/GJ	Reference
SOLID	ANODE CARBON	1A2g	Industry	all	4	1.8	13.5	17.5	7.9	13	134	25	200	2
	COAL	1A1a	Public electricity and heat production	all	0.51	0.07	0.86	0.48	1.3	0.97	0.62	5.9	1.9	1
		All other	All other	All	4	1.8	13.5	17.5	7.9	13	134	25	200	2
	FLY ASH FOSSIL	1A1a	Public electricity and heat production		0.51	0.07	0.86	0.48	1.3	0.97	0.62	5.9	1.9	1
	BROWN COAL BRI.	1A4b i	Residential	0202	2.5	1.5	11.2	22.3	5.1	12.7	130	1	220	2
	COKE OVEN COKE	1A2 a-g	Industry	all	4	1.8	13.5	17.5	7.9	13	134	25	200	2
LIQUID	PETROLEUM COKE	all	All	all	4.3	1.3	2.7	5.7	0.4	362	4.9	2.2	94	2
	RESIDUAL OIL	1A1a	Public electricity and heat production	all	2.1	0.53	2.6	2.4	0.21	362	2.6	1.2	7.4	1
		All other	All other	all	4.3	1.3	2.7	5.7	0.4	362	4.9	2.2	94	2
	GAS OIL	-	Engines (reciprocating)	all	0.055	0.011	0.2	0.3	0.11	0.013	0.15	0.22	58	4
		-	All other	all	0.002	0.001	0.2	0.13	0.12	0.005	0.012	0.002	0.42	3
	KEROSENE	All	All	all	0.002	0.001	0.2	0.13	0.12	0.005	0.012	0.002	0.42	5
	LPG	All	All	all	0.002	0.001	0.2	0.13	0.12	0.005	0.012	0.002	0.42	2
REFINERY GAS	1A1b	Petroleum refining	all	1.8	1.4	1.4	2.7	1.4	1.4	4.1	6.8	1.8	2	
GAS	NATURAL GAS	-	Engines (reciprocating)	all	0.05	0.003	0.05	0.01	0.1	0.05	0.04	0.01	2.9	4
		-	All other	all	0.119	0.00025	0.00076	0.000076	0.1	0.00051	0.0015	0.01	0.0015	7 and 2
WASTE	WASTE	-	All	all	0.59	0.44	1.56	1.3	1.79	2.06	5.52	1.11	2.33	4
	INDUSTRIAL WASTE	1A2f	Industry - Other	all	0.59	0.44	1.56	1.3	1.79	2.06	5.52	1.11	2.33	4
BIOMASS	WOOD	-	All non-residential	all	1.4	0.27	2.34	2.6	0.4	2.34	3.62	0.5	2.3	2 and 4
		1A4b i	Residential	all	0.19	13	23	6	0.56	2	27	0.5	512	8
	STRAW	1A1a	Public electricity and heat production	all	1.4	0.32	1.6	1.7	0.31	1.7	6.2	0.5	0.41	2 and 4
		1A4b i	Residential	0202	1	1.4	2.9	8.6	0.5	4.4	40	0.5	130	2
		1A4c i	Agriculture/ Forestry	0203	1	1.4	2.9	8.6	0.5	4.4	40	0.5	130	2
	BIO OIL	-	Engines	engines	0.055	0.011	0.2	0.3	0.11	0.013	0.15	0.22	58	5
		-	All other	-	0.002	0.001	0.2	0.13	0.12	0.005	0.012	0.002	0.42	5
BIOGAS	-	All	all	0.04	0.002	0.18	0.31	0.12	0.23	0.005	0.21	3.95	4	
BIO GASIF GAS	-	All	all	0.12	0.009	0.029	0.045	0.54	0.014	0.022	0.18	0.058	4	

Reference:

1. Implied emission factor 2008 estimated by DCE based on plant specific emission data for power plants.
2. EMEP/EEA Emission inventory Guidebook, 2009 update (EEA 2009).
3. CONCAWE (Denier van der Gon & Kuenen, 2009).
4. Nielsen et al. 2010.
5. Assumed equal to gas oil. DCE assumption.
6. Assumed equal to natural gas fuelled engines.
7. Gruijthuijsen (2001).
8. EEA (2013)

Table 3A-4.14 As emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.15 Cd emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.16 Cr emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.17 Cu emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.18 Hg emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.19 Ni emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.20 Pb emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.21 Se emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.22 Zn emission factors time series, mg per GJ, for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3A-4.23 PAH emission factors 2013.

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	snap_id	Benzo(a)- pyrene		Benzo(b)- flouran-thene		Benzo(k)- flouran-thene		Indeno- (1,2,3-c,d)- pyrene		
					µg per GJ	Ref.	µg per GJ	Ref.	µg per GJ	Ref.	µg per GJ	Ref.	
SOLID	102A	ANODE CARBON	1A2g	0320	23	4	929	4	929	4	698	4	
			COAL	1A1a	All	0.14	4	0.29	4	0.29	4	0.28	4
			1A2 a-g	All	23	4	929	4	929	4	698	4	
			1A4c i	0203	59524	4	63492	4	1984	4	119048	4	
	103A	FLY ASH FOSSIL	1A1a	0101	0.14	4	0.29	4	0.29	4	0.28	4	
	106A	BROWN COAL BRI.	1A4b i	0202	59524	4 (8)	63492	4 (8)	1984	4 (8)	119048	4 (8)	
	107A	COKE OVEN COKE	1A2 a-g	all	23	4	929	4	929	4	698	4	
LIQUID	110A	PETROLEUM COKE	1A2 a-g	all	3184	5	9554	5	-	-	-	-	
	203A	RESIDUAL OIL	1A1a	All	109.6	4	475.41	4	93.21	4	177.28	4	
			1A1b	010306	109.6	4	475.41	4	93.21	4	177.28	4	
			1A2 a-g	all	80	4	42	4	66	4	160	4	
			1A4a i	all	80	4	42	4	66	4	160	4	
			1A4b i	all	80	4	42	4	66	4	160	4	
			1A4c i	all	80	4	42	4	66	4	160	4	
	204A	GAS OIL	1A1a	Not engines Engines	109.6 1.9	4 7	475.41 15	4 7	93.21 1.7	4 7	177.28 1.5	4 7	
			1A1b	010306	109.6	4	475.41	4	93.21	4	177.28	4	
			1A1c	010504	109.6	4	475.41	4	93.21	4	177.28	4	
			1A2 a-g	Not engines Engines	80 1.9	4 7	42 15	4 7	66 1.7	4 7	160 1.5	4 7	
			1A4a i	Not engines Engines	80 1.9	4 7	42 15	4 7	66 1.7	4 7	160 1.5	4 7	
			1A4b i	0202	80	4	42	4	66	4	160	4	
			1A4c i	0203	80	4	42	4	66	4	160	4	
			301A	NATURAL GAS	1A1a	010104 010105	1 1.2	8 7	1 9	8 7	2 1.7	8 7	3 1.8
	1A1c	010504	1		8	1	8	2	8	3	8		
	1A2 a-g	Turbines Engines	1 1.2		8 7	1 9	8 7	2 1.7	8 7	3 1.8	8 7		
1A4a i	020105	1.2	7		9	7	1.7	7	1.8	7			
1A4b i	020202 020204	0.133 1.2	6 6		0.663 9	6 6	0.265 1.7	6 6	2.653 1.8	6 6			
1A4c i	020304	1.2	6		9	6	1.7	6	1.8	6			
WASTE	114A	WASTE	1A1a		all	0.8	7	1.7	7	0.9	7	1.1	7
			1A4a i		0201	0.8	7	1.7	7	0.9	7	1.1	7
	115A	INDUSTRIAL WASTE	1A2f	0316	0.8	7	1.7	7	0.9	7	1.1	7	
BIOMASS	111A	WOOD	1A1a	0101 0102	11 6.46	7 4	15 1292.52	7 4	5 1292.52	7 4	0.8 11.56	7 4	
			1A2 a-g	all	6.46	4	1292.52	4	1292.52	4	11.56	4	
			1A4a i	0201	168707	4	221769	4	73469	4	119728	4	
			1A4b i	All	52204	9	52191	9	19084	9	29455	9	
			1A4c i	all	168707	4	221769	4	73469	4	119728	4	

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	snap_id	Benzo(a)-pyrene		Benzo(b)-flouran-thene		Benzo(k)-flouran-thene		Indeno-(1,2,3-c,d)-pyrene	
	117A	STRAW	1A1a	0101	0.5	7	0.5	7	0.5	7	0.5	7
				0102	1529	2	3452	2	1400	2	1029	2
			1A4b i	0202	12956	2	12828	2	6912	2	4222	2
			1A4c i	0203	12956	2	12828	2	6912	2	4222	2
	215A	BIO OIL	1A1a	all	109.6	3	475.41	3	93.21	3	177.28	3
			1A2 a-g	all	109.6	3	475.41	3	93.21	3	177.28	3
			1A4b i	0202	80	3	42	3	66	3	160	3
	309A	BIOGAS	Engines	All	1.3	7	1.2	7	1.2	7	0.6	7
	310A	BIO GASIF GAS	Engines	010105	2	7	2	7	2	7	2	7

1. -
2. Same emission factors as for gas oil is assumed (DCE assumption).
3. Berdowski J.J.M., Veldt C., Baas J., Bloos J.P.J., Klein A.E. 1995, Technical Paper to the OSPARCOM-HELCOM-UNECE Emission Inventory of heavy Metals and Persistent Organic Pollutants, TNO-report, TNO-MEP – R 95/247.
4. Finstad A., Haakonsen G., Kvingedal E. & Rypdal K. 2001. Utslipp til luft av noen miljøgifter i Norge, Dokumentasjon av metode og resultater, Statistics Norway Report 2001/17 (In Norwegian).
5. Jensen, J. 2001, Danish Gas Technology centre, personal communication, e-mail 11-10-2001.
6. Nielsen, M., Nielsen, O.K. & Thomsen, M. 2010c: Emissionskortlægning for decentral kraftvarme, Energinet.dk miljøprojekt nr. 07/1882. Delrapport 5. Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme, 2006. National Environmental Research Institute, Aarhus University.
7. Nielsen, M. & Illerup, J.B. 2003. Emissionsfaktorer og emissionsopgørelse for decentral kraftvarme. Eltra PSO projekt 3141. Kortlægning af emissioner fra decentrale kraftværker. Delrapport 6. Danmarks Miljøundersøgelser. 116 s. –Faglig rapport fra DMU nr. 442. (In Danish, with English summary). Available at : [http://www.dmu.dk/1\\_viden/2\\_Publikationer/3\\_fagrappporter/rapporter/FR442.pdf](http://www.dmu.dk/1_viden/2_Publikationer/3_fagrappporter/rapporter/FR442.pdf) (07-02-2011).
8. Same emission factor as for coal is assumed (DCE assumption).
9. Aggregated emission factor based on the technology distribution in the sector and guidebook (EEA 2009) emission factors. Technology distribution based on: DEPA (2013)

Table 3A-4.24 PAH emission factors time series, µg pr GJ for the years 1990 to 2013.

This table is available at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iiir/>

Table 3A-4.25 Emission factors for PCDD/F, 2013.

fuel_type	fuel_id	fuel_gr_abbr	nfr_id	snap_id	PCDD/F, ng per GJ			
SOLID	102A	ANODE CARBON COAL	1A2g	0320	1.32			
			1A1a	0101 and 0102	1.32			
			1A2 a-g	03	1.32			
			1A4c i	0203	300			
			103A	FLY ASH FOSSIL	1A1a	0101	1.32	
			106A	BROWN COAL BRI.	1A4b i	0202	800	
			107A	COKE OVEN COKE	1A2 a-g	03	1.32	
LIQUID	110A	PETROLEUM COKE	1A2 a-g	03	1.32			
			203A	RESIDUAL OIL	1A1a	All	0.882	
			1A1b		010306	0.882		
			1A2 a-g		03	0.882		
			1A4a i		0201	10		
			1A4b i		0202	10		
	1A4c i	0203	10					
	204A	GAS OIL	1A1a	Not engines Engines	0.882 0.99			
			1A1b	010306	0.882			
			1A1c	010504	0.882			
			1A2 a-g	Not engines Engines	0.882 0.99			
			1A4a i	Not engines Engines	10 0.99			
			1A4b i	0202	10			
			1A4c i	0203	10			
			206A	KEROSENE	1A2a-g	03	0.882	
					1A4a i	0201	10	
					1A4b i	0202	10	
					1A4c i	0203	10	
			303A	LPG	1A1a	0101 and 0102	0.025	
	1A2a-g	03			0.025			
	1A4a i	0201			2			
	1A4b i	0202			2			
	1A4c i	0203			2			
	308A	REFINERY GAS	1A1b	0103	0.025			
	GAS	301A	NATURAL GAS	1A1a	Not engines Engines	0.025 0.57		
				1A1c	010504	0.025		
				1A2 a-g	03, Not engines Engines	0.025 0.57		
1A4a i				0201 020105	2 0.57			
1A4b i				0202 020204	2 0.57			
1A4c i				0203 020304	2 0.57			
WASTE				114A	WASTE	1A1a	0101 and 0102	5
						1A4a i	0201	5
						115A	INDUSTRIAL WASTE	1A2f
BIOMASS				111A	WOOD	1A1a	0101 0102	14 1
	1A2 a-g	03	1					
	1A4a i	0201	400					
	1A4b i	0202	347					
	1A4c i	0203	400					
	117A	STRAW	1A1a			0101 0102	19 22	



fuel_type	fuel_id	fuel_gr_abbr	nfr_id	snap_id	PCDD/F, ng per GJ
			1A4b i	0202	500
			1A4c i	0203	400
	215A	BIO OIL	1A1a	0101 and 0102	0.882
			1A2 a-g	03	0.882
			1A4b i	0202	10
	309A	BIOGAS	1A1a	Engines	0.96
				Not engines	0.025
			1A2a-g	Not engines	0.025
				Engines	0.96
			1A4a i	Not engines	0.025
				Engines	0.96
			1A4c i	Not engines	0.025
				Engines	0.96
	310A	BIO GASIF GAS	Engines	010105	1.7

Table 3A-4.26 Emission factor time series for PCDD/F.

Year	Waste incineration, PCDD/F, ng/GJ	Residential wood combustion, PCDD/F, ng/GJ
1990	2095	696
1991	1746	696
1992	1396	694
1993	1047	693
1994	907	690
1995	767	685
1996	628	676
1997	488	668
1998	348	657
1999	253	618
2000	157	590
2001	157	537
2002	157	516
2003	157	512
2004	81	507
2005	5	487
2006	5	465
2007	5	470
2008	5	446
2009	5	416
2010	5	399
2011	5	383
2012	5	368
2013	5	347

References for HCB are discussed in Nielsen et al. (2014). Table 3A-4.27 presents the applied emission factors.

Table 3A-4.27 Emission factors for HCB, 2013.

<b>Fuel</b>	<b>NFR (SNAP)</b>	<b>Emission factor, Reference ng/GJ</b>
Coal	1A1, 1A2	6,700 Grochowalski & Koniecznyński (2008); EEA (2013)
Coal	1A4b	1,200,000 Syc et al. (2011)
Coal	1A4a and 1A4c	23,000 Syc et al. (2011)
Other solid fuels	1A1, 1A2	6,700 Assumed equal to coal.
Other solid fuels	1A4	1,200,000 Assumed equal to coal.
Liquid fuels <sup>1)</sup>	1A1, 1A2, 1A4	220 Nielsen et al. (2010)
Gaseous fuels	1A1, 1A2, 1A4	- Negligible
Waste	1A1, 1A2, 1A4	4,300 Nielsen et al. (2010). A time series have been estimated. The emission factor for 1990 (190,000 ng/GJ) refer to Pacyna et al. (2003).
Wood	1A1, 1A2	5,000 EEA (2013)
Wood	1A4	5,000 EEA (2013)
Straw	1A1, 1A2	113 Nielsen et al. (2010)
Straw	1A4	5,000 EEA (2013)
Biogas	1A1, 1A2, 1A4	190 Nielsen et al. (2010)
Bio gasification gas	1A1, 1A2, 1A4	800 Nielsen et al. (2010)

1) Except LPG and refinery gas.

Table 3A-4.28 Emission factor time series for HCB from waste incineration.

<b>Year</b>	<b>HCB, ng/GJ</b>
1990	190000
1991	158000
1992	127000
1993	95000
1994	82000
1995	70000
1996	57000
1997	45000
1998	32000
1999	23000
2000	14000
2001	12000
2002	10000
2003	8000
2004	6000
2005	4300
2006	4300
2007	4300
2008	4300
2009	4300
2010	4300
2011	4300
2012	4300
2013	4300

Table 3A-4.29 Emission factors for PCB.

Fuel	NFR (SNAP)	Emission factor, $\sum$ dl-PCB, ng/GJ	Reference
Coal	1A1	839	Grochowalski & Koniecznyński (2008)
Coal	1A2	5,700	Thistlethwaite (2001a)
Coal	1A4	7,403	Syc et al. (2011)
Other solid fuels	1A1	839	Assumed equal to coal.
Other solid fuels	1A2	5,700	Assumed equal to coal.
Other solid fuels	1A4	7,403	Assumed equal to coal.
Residual oil	1A1, 1A2, 1A4	839	The teq value refers to Dyke et al. (2003).
The TEQ value is equal to the emission factor for coal combustion in power plants and the sum of dioxin-like PCB congeners has been assumed equal to the corresponding factor for coal.			
Gas oil	1A1, 1A2, 1A4	93	Nielsen et al. (2010)
Other liquid fuels <sup>1)</sup>	1A1, 1A2, 1A4	93	Assumed equal to gas oil.
Gaseous fuels	1A1, 1A2, 1A4	-	Negligible
Waste	1A1, 1A2, 1A4	109 (time series)	Nielsen et al. (2010). A time series have been estimated. The emission factor for 1990 (46,000 ng/GJ / 117 ng WHO <sub>1996</sub> teq/GJ) have been estimated based on the assumption that the PCB emission factor time series follow the PCDD/F time series.
Wood	1A1, 1A2, 1A4a/c	2,800	Thistlethwaite (2001a)
Wood	1A4b	2,752 (time series)	Hedman et al. (2006). A time series have been estimated based on time series for technologies applied in Denmark.
Straw	1A1, 1A2	3,110	Assumed equal to residential plants.
Straw	1A4	3,110	Syc et al. (2011)
Biogas	1A1, 1A2, 1A4	90	Nielsen et al. (2010)
Bio gasification gas	1A1, 1A2, 1A4	144	Nielsen et al. (2010)

1. Except LPG and refinery gas

Emission factor time series for waste incineration and for residential wood combustion are shown in Table 3A-4.30.

Table 3A-4.30 PCB emission factor time series for waste incineration and for residential wood combustion.

Year	Waste incineration $\sum$ dl-PCB, ng/GJ	Residential wood combustion $\sum$ dl-PCB, ng/GJ
1990	45671	6649
1991	38063	6623
1992	30433	6589
1993	22825	6560
1994	19773	6511
1995	16721	6446
1996	13690	6329
1997	10638	6232
1998	7586	6103
1999	5515	5696
2000	3423	5390
2001	3423	4834
2002	3423	4609
2003	3423	4548
2004	1766	4477
2005	109	4267
2006	109	4028
2007	109	4050
2008	109	3801
2009	109	3496
2010	109	3309
2011	109	3138
2012	109	2974
2013	109	2752

Table 3A-4.31 Technology specific PCB emission factors for residential wood combustion.

Technology	dl-PCB emission factor, ng WHO- teq/GJ	$\Sigma$ dl-PCB emission factor, ng/GJ	Reference and assumptions
Old stove	53	7049	Hedman (2006), old boiler
New stove	53	7049	Hedman (2006), old boiler
Modern stove	7	931	Hedman (2006), modern boiler
Eco labelled stove	3.5	466	Hedman (2006), assumed 1/2 modern boiler
Other stove	53	7049	Hedman (2006), old boiler
Old boiler with acc. tank	53	7049	Hedman (2006), old boiler
Old boiler without acc. tank	53	7049	Hedman (2006), old boiler
New boiler with acc. tank	7	931	Hedman (2006), modern boiler
New boiler without acc. tank	7	931	Hedman (2006), modern boiler
Pellet boilers/stoves	3.5	466	Hedman (2006), assumed 1/2 modern boiler

Table 3A-4.32 Emission factors for NH<sub>3</sub>, 2013.

Fuel	NFR (SNAP)	Emission factor, g/GJ	Reference
Coal/BKB/COKE	1A4b	3.8	EEA (2009)
Wood	1A4b	40.9	EEA (2013), technology distribution based on DEPA (2013)
Waste	1A1a	0.29	Nielsen et al. (2010)
Straw	1A4b	3.8	EEA (2009)

Table 3A-4.33 Emission factor time series for NH<sub>3</sub>, residential wood combustion.

Year	NH <sub>3</sub> emission, g/GJ
1990	68.6
1991	68.5
1992	68.3
1993	68.1
1994	67.8
1995	67.3
1996	66.3
1997	65.5
1998	64.4
1999	61.0
2000	58.5
2001	53.7
2002	51.9
2003	51.8
2004	51.5
2005	49.9
2006	48.0
2007	49.2
2008	47.7
2009	45.5
2010	44.3
2011	43.4
2012	42.5
2013	40.9

Table 3A-4.34 BC fraction of PM<sub>2.5</sub>, 2013.

Fuel_id	Fuel	SNAP	BC_% Reference: EEA Guidebook 2013:
101A	Anodic carbon	032000	2.2% Energy Industries, Table 3-2
102A	Coal	010100	2.2% Energy Industries, Table 3-2
102A	Coal	010101	2.2% Energy Industries, Table 3-2
102A	Coal	010102	2.2% Energy Industries, Table 3-2
102A	Coal	010103	2.2% Energy Industries, Table 3-2
102A	Coal	010104	2.2% Energy Industries, Table 3-2
102A	Coal	010105	2.2% Energy Industries, Table 3-2
102A	Coal	010200	2.2% Energy Industries, Table 3-2
102A	Coal	010201	2.2% Energy Industries, Table 3-2
102A	Coal	010202	2.2% Energy Industries, Table 3-2
102A	Coal	010203	2.2% Energy Industries, Table 3-2
102A	Coal	020100	6.4% Small Combustion, Table 3-7
102A	Coal	020200	6.4% Small Combustion, Table 3-3
102A	Coal	020300	6.4% Small Combustion, Table 3-7
102A	Coal	020304	6.4% Small Combustion, Table 3-7
102A	Coal	030100	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030102	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030103	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030400	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030500	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030600	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030700	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030703	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030800	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030900	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030902	6.4% Manufacturing Industries, Table 3-2
102A	Coal	030903	6.4% Manufacturing Industries, Table 3-2
102A	Coal	031100	6.4% Manufacturing Industries, Table 3-2
102A	Coal	031102	6.4% Manufacturing Industries, Table 3-2
102A	Coal	031200	6.4% Manufacturing Industries, Table 3-2
102A	Coal	031300	6.4% Manufacturing Industries, Table 3-2
102A	Coal	031400	6.4% Manufacturing Industries, Table 3-2
102A	Coal	031600	6.4% Manufacturing Industries, Table 3-2
102A	Coal	032000	6.4% Manufacturing Industries, Table 3-2
103A	Fly ash fossil	010104	2.2% Assumed equal to coal. DCE assumption.
106A	Brown coal bri.	020100	6.4% Small Combustion, Table 3-7
106A	Brown coal bri.	020200	6.4% Small Combustion, Table 3-3
106A	Brown coal bri.	020300	6.4% Small Combustion, Table 3-7
106A	Brown coal bri.	030100	6.4% Manufacturing Industries, Table 3-2
106A	Brown coal bri.	030800	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	020200	6.4% Small Combustion, Table 3-3
107A	Coke oven coke	030100	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	030400	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	030700	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	030800	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	030900	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	030902	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	030903	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	031200	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	031300	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	031400	6.4% Manufacturing Industries, Table 3-2
107A	Coke oven coke	032000	6.4% Manufacturing Industries, Table 3-2
110A	Petroleum coke	010100	5.6% Energy Industries, table 3-5
110A	Petroleum coke	010102	5.6% Energy Industries, table 3-5
110A	Petroleum coke	020100	56.0% Small Combustion, Table 3-5
110A	Petroleum coke	020200	8.5% Small Combustion, Table 3-5
110A	Petroleum coke	020300	56.0% Small Combustion, Table 3-5
110A	Petroleum coke	030100	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	030400	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	030600	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	030700	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	030800	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	030900	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	031000	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	031100	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	031300	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	031400	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	031600	56.0% Manufacturing Industries, Table 3-4
110A	Petroleum coke	032000	56.0% Manufacturing Industries, Table 3-4
111A	Wood	010100	3.3% Energy Industries, Table 3-7
111A	Wood	010101	3.3% Energy Industries, Table 3-7
111A	Wood	010102	3.3% Energy Industries, Table 3-7



<b>Fuel_id</b>	<b>Fuel</b>	<b>SNAP</b>	<b>BC % Reference: EEA Guidebook 2013:</b>
117A	Straw	010202	3.3% Energy Industries, Table 3-7
117A	Straw	010203	3.3% Energy Industries, Table 3-7
117A	Straw	020103	28.0% Small Combustion, Table 3-10
117A	Straw	020200	10.0% Small Combustion, Table 3-6
117A	Straw	020300	28.0% Small Combustion, Table 3-10
117A	Straw	020302	28.0% Small Combustion, Table 3-10
117A	Straw	030100	28.0% Manufacturing Industries, Table 3-5
117A	Straw	030103	28.0% Manufacturing Industries, Table 3-5
117A	Straw	030105	28.0% Manufacturing Industries, Table 3-5
117A	Straw	030903	28.0% Manufacturing Industries, Table 3-5
117A	Straw	031305	28.0% Manufacturing Industries, Table 3-5
117A	Straw	032003	28.0% Manufacturing Industries, Table 3-5
203A	Residual oil	010100	5.6% Energy Industries, Table 3-5
203A	Residual oil	010101	5.6% Energy Industries, Table 3-5
203A	Residual oil	010102	5.6% Energy Industries, Table 3-5
203A	Residual oil	010103	5.6% Energy Industries, Table 3-5
203A	Residual oil	010104	5.6% Energy Industries, Table 3-5
203A	Residual oil	010105	5.6% Energy Industries, Table 3-5
203A	Residual oil	010200	5.6% Energy Industries, Table 3-5
203A	Residual oil	010202	5.6% Energy Industries, Table 3-5
203A	Residual oil	010203	5.6% Energy Industries, Table 3-5
203A	Residual oil	010306	5.6% Energy Industries, Table 4-4
203A	Residual oil	020100	56.0% Small Combustion, Table 3-9
203A	Residual oil	020103	56.0% Small Combustion, Table 3-9
203A	Residual oil	020200	8.5% Small Combustion, Table 3-5
203A	Residual oil	020300	56.0% Small Combustion, Table 3-9
203A	Residual oil	020302	56.0% Small Combustion, Table 3-9
203A	Residual oil	020304	56.0% Small Combustion, Table 3-9
203A	Residual oil	030100	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030102	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030103	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030104	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030105	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030400	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030403	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030500	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030600	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030603	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030700	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030800	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030900	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030902	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030903	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030904	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	030905	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031000	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031100	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031102	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031200	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031300	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031305	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031400	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031403	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031500	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031503	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031600	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	031603	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	032000	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	032002	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	032003	56.0% Manufacturing Industries, Table 3-4
203A	Residual oil	032005	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	010100	33.5% Energy Industries, Table 3-6
204A	Gas oil	010101	33.5% Energy Industries, Table 3-6
204A	Gas oil	010102	33.5% Energy Industries, Table 3-6
204A	Gas oil	010103	33.5% Energy Industries, Table 3-6
204A	Gas oil	010104	33.5% Energy Industries, Table 3-18
204A	Gas oil	010105	78.0% Energy Industries, Table 3-19
204A	Gas oil	010200	33.5% Energy Industries, Table 3-6
204A	Gas oil	010201	33.5% Energy Industries, Table 3-6
204A	Gas oil	010202	33.5% Energy Industries, Table 3-6
204A	Gas oil	010203	33.5% Energy Industries, Table 3-6
204A	Gas oil	010204	33.5% Energy Industries, Table 3-18
204A	Gas oil	010205	78.0% Energy Industries, Table 3-19

<b>Fuel_id</b>	<b>Fuel</b>	<b>SNAP</b>	<b>BC % Reference: EEA Guidebook 2013:</b>
204A	Gas oil	010306	33.5% Energy Industries, Table 4-5
204A	Gas oil	010504	33.5% Energy Industries, Table 3-18
204A	Gas oil	010505	78.0% Energy Industries, Table 3-19
204A	Gas oil	020100	56.0% Small Combustion, Table 3-9
204A	Gas oil	020102	56.0% Small Combustion, Table 3-9
204A	Gas oil	020103	56.0% Small Combustion, Table 3-9
204A	Gas oil	020105	78.0% Energy Industries, Table 3-37
204A	Gas oil	020200	3.9% Small Combustion, Table 3-21
204A	Gas oil	020204	78.0% Energy Industries, Table 3-19
204A	Gas oil	020300	56.0% Small Combustion, Table 3-9
204A	Gas oil	020302	56.0% Small Combustion, Table 3-9
204A	Gas oil	020304	78.0% Energy Industries, Table 3-37
204A	Gas oil	030100	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030102	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030103	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030104	33.5% Energy Industries, Table 3-18
204A	Gas oil	030105	78.0% Energy Industries, Table 3-19
204A	Gas oil	030400	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030402	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030403	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030500	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030600	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030602	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030604	33.5% Energy Industries, Table 3-18
204A	Gas oil	030700	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030703	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030800	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030900	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030902	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030903	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	030904	33.5% Energy Industries, Table 3-18
204A	Gas oil	031000	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031100	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031102	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031103	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031200	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031205	78.0% Energy Industries, Table 3-19
204A	Gas oil	031300	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031305	78.0% Energy Industries, Table 3-19
204A	Gas oil	031400	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031403	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031505	78.0% Energy Industries, Table 3-19
204A	Gas oil	031600	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	031603	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	032000	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	032002	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	032003	56.0% Manufacturing Industries, Table 3-4
204A	Gas oil	032005	78.0% Energy Industries, Table 3-19
206A	Kerosene	020100	56.0% Small Combustion, Table 3-9
206A	Kerosene	020200	8.5% Small Combustion, Table 3-5
206A	Kerosene	020300	56.0% Small Combustion, Table 3-9
206A	Kerosene	030100	56.0% Manufacturing Industries, Table 3-4
206A	Kerosene	031500	56.0% Manufacturing Industries, Table 3-4
206A	Kerosene	032000	56.0% Manufacturing Industries, Table 3-4
215A	Bio oil	010101	33.5% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	010102	33.5% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	010103	33.5% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	010105	78.0% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	010200	33.5% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	010202	33.5% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	010203	33.5% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	020105	78.0% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	020200	3.9% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	020304	78.0% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	030100	56.0% Manufacturing Industries, Table 3-4
215A	Bio oil	030103	56.0% Manufacturing Industries, Table 3-4
215A	Bio oil	030105	78.0% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	030605	56.0% Manufacturing Industries, Table 3-4
215A	Bio oil	030903	56.0% Manufacturing Industries, Table 3-4
215A	Bio oil	031305	78.0% Assumed equal to gas oil. DCE assumption.
215A	Bio oil	031600	56.0% Manufacturing Industries, Table 3-4
215A	Bio oil	032005	78.0% Assumed equal to gas oil. DCE assumption.
225A	Orimulsion	010101	2.2% Assumed equal to coal. DCE assumption.



<b>Fuel_id</b>	<b>Fuel</b>	<b>SNAP</b>	<b>BC % Reference: EEA Guidebook 2013:</b>
301A	Natural gas	010100	2.5% Energy Industries, Table 3-4
301A	Natural gas	010101	2.5% Energy Industries, Table 3-4
301A	Natural gas	010102	2.5% Energy Industries, Table 3-4
301A	Natural gas	010103	2.5% Energy Industries, Table 3-4
301A	Natural gas	010104	2.5% Energy Industries, Table 3-17
301A	Natural gas	010105	2.5% Energy Industries, Table 3-20
301A	Natural gas	010200	2.5% Energy Industries, Table 3-4
301A	Natural gas	010202	2.5% Energy Industries, Table 3-4
301A	Natural gas	010203	2.5% Energy Industries, Table 3-4
301A	Natural gas	010205	2.5% Energy Industries, Table 3-4
301A	Natural gas	010502	2.5% Energy Industries, Table 3-4
301A	Natural gas	010503	2.5% Energy Industries, Table 3-4
301A	Natural gas	010504	2.5% Energy Industries, Table 3-17
301A	Natural gas	010505	2.5% Energy Industries, Table 3-20
301A	Natural gas	020100	4.0% Small Combustion, Table 3-8
301A	Natural gas	020103	4.0% Small Combustion, Table 3-8
301A	Natural gas	020104	2.5% Small Combustion, Table 3-34
301A	Natural gas	020105	2.5% Energy Industries, Table 3-36
301A	Natural gas	020200	5.4% Small Combustion, Table 3-19
301A	Natural gas	020202	5.4% Small Combustion, Table 3-19
301A	Natural gas	020204	2.5% Energy Industries, Table 3-20
301A	Natural gas	020300	4.0% Small Combustion, Table 3-8
301A	Natural gas	020303	2.5% Energy Industries, Table 3-17
301A	Natural gas	020304	2.5% Energy Industries, Table 3-36
301A	Natural gas	030100	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030102	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030103	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030104	2.5% Energy Industries, Table 3-17
301A	Natural gas	030105	2.5% Energy Industries, Table 3-20
301A	Natural gas	030106	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030400	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030402	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030500	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030600	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030602	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030603	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030604	2.5% Energy Industries, Table 3-17
301A	Natural gas	030605	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030700	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030703	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030705	2.5% Energy Industries, Table 3-20
301A	Natural gas	030800	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030900	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030902	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030903	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	030904	2.5% Energy Industries, Table 3-17
301A	Natural gas	030905	2.5% Energy Industries, Table 3-20
301A	Natural gas	031000	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031005	2.5% Energy Industries, Table 3-20
301A	Natural gas	031100	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031102	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031103	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031104	2.5% Energy Industries, Table 3-17
301A	Natural gas	031200	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031205	2.5% Energy Industries, Table 3-20
301A	Natural gas	031300	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031305	2.5% Energy Industries, Table 3-20
301A	Natural gas	031400	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031405	2.5% Energy Industries, Table 3-20
301A	Natural gas	031500	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031503	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	031604	2.5% Energy Industries, Table 3-17
301A	Natural gas	031605	2.5% Energy Industries, Table 3-20
301A	Natural gas	032000	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	032003	4.0% Manufacturing Industries, Table 3-3
301A	Natural gas	032004	2.5% Energy Industries, Table 3-17
301A	Natural gas	032005	2.5% Energy Industries, Table 3-20
303A	LPG	010100	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010101	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010102	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010103	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010104	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010200	2.5% Assumed equal to natural gas. DCE assumption.

Fuel_id	Fuel	SNAP	BC % Reference: EEA Guidebook 2013:
303A	LPG	010202	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010203	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	010306	2.5% Assumed equal to natural gas. DCE assumption.
303A	LPG	020100	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	020103	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	020105	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	020200	5.4% Assumed equal to natural gas. DCE assumption.
303A	LPG	020300	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030100	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030400	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030402	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030500	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030600	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030602	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030700	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030800	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	030900	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	031000	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	031100	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	031200	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	031300	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	031400	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	031500	4.0% Assumed equal to natural gas. DCE assumption.
303A	LPG	032000	4.0% Assumed equal to natural gas. DCE assumption.
308A	Refinery gas	010101	18.4% Energy Industries, Table 4-2
308A	Refinery gas	010203	18.4% Energy Industries, Table 4-2
308A	Refinery gas	010300	18.4% Energy Industries, Table 4-2
308A	Refinery gas	010304	18.4% Energy Industries, Table 4-2
308A	Refinery gas	010306	18.4% Energy Industries, Table 4-2
308A	Refinery gas	032000	18.4% Energy Industries, Table 4-2
309A	Biogas	010100	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010101	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010102	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010103	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010104	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010105	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010200	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010203	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010205	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	010505	3.3% Assumed % equal to wood. DCE assumption
309A	Biogas	020100	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	020103	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	020104	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	020105	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	020300	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	020304	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030100	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030102	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030103	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030104	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030105	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030400	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030900	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030902	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030903	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	030905	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	031300	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	032000	28.0% Assumed % equal to wood. DCE assumption
309A	Biogas	032005	28.0% Assumed % equal to wood. DCE assumption
310A	Bio gasif. gas	010105	3.3% Assumed % equal to wood. DCE assumption
310A	Bio gasif. gas	020105	3.3% Assumed % equal to wood. DCE assumption
310A	Bio gasif. gas	020304	28.0% Assumed % equal to wood. DCE assumption
310A	Bio gasif. gas	030105	28.0% Assumed % equal to wood. DCE assumption
310A	Bio gasif. gas	031305	28.0% Assumed % equal to wood. DCE assumption

Table 3A-4.35 Residential wood combustion, BC fraction of PM<sub>2.5</sub>, time series.

<b>Year</b>	<b>% of PM<sub>2.5</sub></b>
1990	12.30
1991	12.29
1992	12.29
1993	12.28
1994	12.27
1995	12.27
1996	12.26
1997	12.26
1998	12.25
1999	12.29
2000	12.37
2001	12.46
2002	12.54
2003	12.62
2004	12.70
2005	12.78
2006	12.89
2007	13.08
2008	13.31
2009	13.52
2010	13.71
2011	13.92
2012	14.15
2013	14.40

## Annex 3A-5 Implied emission factors for waste incineration plants and power plants combustion coal

Table 3A-5.1 Implied emission factors for municipal waste incineration plants 2013.

Pollutant	Implied Emission factor Unit
SO <sub>2</sub>	6.0 g pr GJ
NO <sub>x</sub>	93 g pr GJ
TSP	0.80 g pr GJ
PM <sub>10</sub>	0.64 g pr GJ
PM <sub>2.5</sub>	0.50 g pr GJ
As	0.54 mg pr GJ
Cd	0.36 mg pr GJ
Cr	1.42 mg pr GJ
Cu	1.32 mg pr GJ
Hg	1.58 mg pr GJ
Ni	2.45 mg pr GJ
Pb	4.58 mg pr GJ
Se	1.16 mg pr GJ
Zn	2.85 mg pr GJ

Table 3A-5.2 Implied emission factors for power plants combusting coal, 2013.

Pollutant	Implied Emission factor	Unit
SO <sub>2</sub>	12	g pr GJ
NO <sub>x</sub>	29	g pr GJ
TSP	3.7	g pr GJ
PM <sub>10</sub>	3.2	g pr GJ
PM <sub>2.5</sub>	2.6	g pr GJ
As	0.69	mg pr GJ
Cd	0.07	mg pr GJ
Cr	0.79	mg pr GJ
Cu	0.64	mg pr GJ
Hg	0.73	mg pr GJ
Ni	1.1	mg pr GJ
Pb	0.75	mg pr GJ
Se	4.8	mg pr GJ
Zn	2.2	mg pr GJ

## Annex 3A-6 Large point sources

Table 3A-6.1 Large point sources, 2013.

---

Large point sources
AffaldPlus+, Naestved Forbraendingsanlaeg
AffaldPlus+, Naestved Kraftvarmevaerk
Affaldplus+, Slagelse Forbr. and DONG Slagelse KVV
Affaldscenter aarhus - Forbraendsanlaegget
Affaldsforbraendingsanlaeg I/S REFA
Amagerforbraending
Amagervaerket
Ardagh Glass Holmegaard A/S
Asnaesvaerket
Avedoerevaerket
AVV Forbraendingsanlaeg
Bofa I/S
Centralkommunernes Transmissionsselskab F_berg
Cheminova
Dalum Papir
Danisco Grindsted
DanSteel
DTU
Energi Randers Produktion
Enstedvaerket
Esbjergvaerket
Faxe Kalk
Fjernvarme Fyn, Centrum Varmecentral
Frederikshavn Affaldskraftvarmevaerk
Frederikshavn Kraftvarmevaerk
Fynsvaerket
Grenaa Forbraending
Grenaa Kraftvarmevaerk
H.C.Oerstedsvaerket
Haderslev Kraftvarmevaerk
Haldor Topsoee
Hammel Fjernvarmeselskab
Helsingoer Kraftvarmevaerk
Herningvaerket
Hilleroed Kraftvarmevaerk
Hjoerring Varmeforsyning
Horsens Kraftvarmevaerk
I/S Faelles Forbraending
I/S Kara Affaldsforbraendingsanlaeg
I/S Kraftvarmevaerk Thisted
I/S Nordforbraending
I/S Reno Nord
I/S Reno Syd
I/S Vestforbraending
Koege Kraftvarmevaerk
Kolding Forbraendingsanlaeg TAS
Kommunekemi
Koppers
Kyndbyvaerket
L90 Affaldsforbraending
Maricogen
Masnedoevaerket
Maabjergvaerket
Nordic Sugar Nakskov
Nordic Sugar Nykoebing
Nordjyllandsvaerket
Nybro Gasbehandlingsanlaeg
Odense Kraftvarmevaerk
Oestkraft
Rensningsanlaegget Lynetten
Rockwool A/S Doense
Rockwool A/S Vamdrup
Saint-Gobain Isover A/S
Shell Raffinaderi
Silkeborg Kraftvarmevaerk
Skaerbaekvaerket
Skagen Forbraending
Soenderborg Kraftvarmevaerk
Special Waste System

---

---

Statoil Raffinaderi
Stigsnaesvaerket
Studstrupvaerket
Svanemoellevaerket
Svendborg Kraftvarmevaerk
Viborg Kraftvarme
Vordingborg Kraftvarme
Aalborg Portland
AarhusKarlskamm Denmark A/S

---

Table 3A-6.2 Large point sources, aggregated fuel consumption in 2013.

Year		2013	
nfr_id_EA	fuel_id	fuel_gr_abbr	Sum of Fuel_TJ
1A1a	102A	COAL	130035
	103A	SUB-BITUMINOUS	83
	107A	COKE OVEN COKE	0
	111A	WOOD	30300
	114A	WASTE	34706
	117A	STRAW	9324
	203A	RESIDUAL OIL	1292
	204A	GAS OIL	1328
	215A	BIO OIL	46
	301A	NATURAL GAS	23102
	303A	LPG	22
	309A	BIOGAS	142
1A1a Total			230381
1A1b	203A	RESIDUAL OIL	656
	204A	GAS OIL	1
	303A	LPG	0
	308A	REFINERY GAS	14751
1A1b Total			15408
1A1c	204A	GAS OIL	0
	301A	NATURAL GAS	139
1A1c Total			140
1A2a	204A	GAS OIL	0
	301A	NATURAL GAS	1222
	303A	LPG	4
1A2a Total			1226
1A2c	203A	RESIDUAL OIL	191
	204A	GAS OIL	2
	301A	NATURAL GAS	1499
	303A	LPG	1
1A2c Total			1693
1A2d	111A	WOOD	454
	204A	GAS OIL	0
	301A	NATURAL GAS	2
1A2d Total			455
1A2e	102A	COAL	1166
	107A	COKE OVEN COKE	121
	111A	WOOD	12
	203A	RESIDUAL OIL	2586
	204A	GAS OIL	18
	215A	BIO OIL	51
	301A	NATURAL GAS	55
	309A	BIOGAS	90
1A2e Total			4099
1A2f	102A	COAL	1418
	110A	PETROLEUM COKE	6014
	115A	INDUSTR. WASTES	1781
	203A	RESIDUAL OIL	186
	204A	GAS OIL	4
	215A	BIO OIL	0
	301A	NATURAL GAS	6
1A2f Total			9409
1A2g viii	101A	ANODIC CARBON	22
	102A	COAL	93
	107A	COKE OVEN COKE	400
	204A	GAS OIL	1
	301A	NATURAL GAS	1171
	303A	LPG	0
1A2g viii Total			1687
1A4a i	114A	WASTE	116
	204A	GAS OIL	0
	309A	BIOGAS	0
1A4a i Total			116
Grand Total			264613

Table 3A-6.3 Large point sources, plant specific emissions<sup>1)</sup>.

Year	2013		SO <sub>2</sub>	NO <sub>x</sub>	NMVO C	CO	NH <sub>3</sub>	TSP	PM <sub>10</sub> <sup>(2)</sup>	PM <sub>2.5</sub> <sup>(2)</sup>	As	Cd	Cr	Cu	Hg	Ni	Pb	Se	Zn	PCDD/ F	
nfr_id	EA	lps_name																			
1A1a		AffaldPlus+, Naestved Forbraendingsanlaeg	x	x	x	x		x	x	x					x						
1A1a		AffaldPlus+, Naestved Kraftvarmevaerk		x		x															
1A1a		Affaldplus+, Slagelse Forbr. and DONG Slagelse KVV	x	x		x															x
1A1a		Affaldscenter aarhus - Forbraendsanlaegget	x	x	x			x	x	x					x						x
1A1a		Affaldsforbraendingsanlaeg I/S REFA	x	x		x					x	x	x	x	x	x	x				
1A1a		Amagerforbraending	x	x		x	x	x	x						x						x
1A1a		Amagervaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Asnaesvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Avedoerevaerket	x	x		x		x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		AVV Forbraendingsanlaeg	x	x	x	x									x						x
1A1a		Bofa I/S	x	x		x					x	x	x	x	x	x	x				x
1A1a		Centralkommunernes Transmissionsselskab F_berg	x	x																	
1A1a		DTU		x		x															
1A1a		Energi Randers Produktion	x	x				x	x	x											
1A1a		Enstedvaerket									x	x	x	x	x	x	x	x	x		
1A1a		Esbjergvaerket	x	x		x		x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Fjernvarme Fyn, Centrum Varmecentral		x																	
1A1a		Frederikshavn Affaldskraftvarmevaerk	x	x	x	x		x	x	x	x	x	x	x	x	x	x				x
1A1a		Frederikshavn Kraftvarmevaerk	x	x				x	x	x											
1A1a		Fynsvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Grenaa Kraftvarmevaerk	x	x		x		x	x	x											
1A1a		H.C.Oerstedsvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Haderslev Kraftvarmevaerk	x	x		x		x	x	x											
1A1a		Helsingoer Kraftvarmevaerk		x																	
1A1a		Herningvaerket	x	x		x		x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Hilleroed Kraftvarmevaerk		x																	
1A1a		Hjoerring Varmeforsyning		x		x															
1A1a		Horsens Kraftvarmevaerk	x	x				x	x	x											
1A1a		I/S Faelles Forbraending	x	x		x		x	x	x					x						x
1A1a		I/S Kara Affaldsforbraendingsanlaeg	x	x		x		x	x	x					x						x
1A1a		I/S Nordforbraending	x	x		x	x	x	x	x					x						x
1A1a		I/S Reno Nord	x	x	x	x		x	x	x											
1A1a		I/S Reno Syd	x	x	x	x		x	x	x					x						x
1A1a		I/S Vestforbraending	x	x	x	x		x	x	x	x	x	x	x	x	x	x				x
1A1a		Koege Kraftvarmevaerk	x	x		x		x	x	x											
1A1a		Kolding Forbraendingsanlaeg TAS	x	x	x	x	x	x	x	x					x						x
1A1a		Kommunekemi	x	x	x	x		x	x	x											
1A1a		Kyndbyvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		L90 Affaldsforbraending	x	x		x		x	x	x	x	x	x	x	x	x	x				x
1A1a		Masnedeovaerket	x	x																	
1A1a		Maabjergvaerket	x	x		x		x	x	x	x	x	x	x	x	x	x				
1A1a		Nordjyllandsvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x		
1A1a		Odense Kraftvarmevaerk	x	x	x	x		x	x	x	x	x	x	x	x	x	x				x
1A1a		Oestkraft	x	x				x													
1A1a		Silkeborg Kraftvarmevaerk		x																	



1A1a	Skaerbaekvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x		
1A1a	Skagen Forbraending	x	x	x												x			x
1A1a	Soenderborg Kraftvarmevaerk	x	x	x	x		x	x	x	x					x				x
1A1a	Special Waste System	x	x	x	x										x				x
1A1a	Studstrupvaerket	x	x				x	x	x	x	x	x	x	x	x	x	x	x	
1A1a	Svanemoellevaerket				x														
1A1a	Svendborg Kraftvarmevaerk	x	x	x	x		x	x	x							x			x
1A1a	Viborg Kraftvarme																		
1A1a	Vordingborg Kraftvarme	x	x																
1A1b	Shell Raffinaderi	x	x																
1A1b	Statoil Raffinaderi	x	x																
1A1c	Nybro Gasbehandlingsanlaeg																		
1A2a	DanSteel																		
1A2c	Cheminova																		
1A2c	Haldor Topsoee																		
1A2c	Koppers	x	x	x															
1A2e	Danisco Grindsted																		
1A2e	Maricogen																		
1A2e	Nordic Sugar Nakskov	x	x																
1A2e	Nordic Sugar Nykoebing	x	x				x	x	x										
1A2e	AarhusKarlshamn Denmark A/S	x	x				x	x	x										
1A2f	Faxe Kalk	x	x																
1A2f	Aalborg Portland	x	x		x		x	x	x					x					
1A2g viii	Ardagh Glass Holmegaard A/S					x													
1A2g viii	Rockwool A/S Doense	x	x																
1A2g viii	Rockwool A/S Vamdrup	x	x																
1A2g viii	Saint-Gobain Isover A/S																		
1A4a i	Rensningsanlaegget Lynetten	x	x		x		x	x	x		x			x		x			x
Grand Total		4319	13040	7.2	2918	7.2	698	592	453	94	14	122	101	180	158	148	667	375	189
		8612	30986	15748	11509														1856
Total emission from stationary combustion					2	1303	16511	15456	14849	281	494	1083	610	303	2115	2138	975	4	21775
Share of total emission from stationary combustion based on plant specific data, %.		50%	42%	0.05%	3%	1%	4%	4%	3%	33%	3%	11%	17%	59%	7%	7%	68%	2%	0.9%

1) Emissions of the pollutants marked with "x" are plant specific. Emission of other pollutants is estimated based on emission factors. The total shown *in this table* only includes plant specific data.

2) Based on particle size distribution.

### Annex 3A-7 Uncertainty estimates, 2013

Table 3A-7.1 Uncertainty estimation.

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Mg SO <sub>2</sub>	Input data Mg SO <sub>2</sub>	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	SO <sub>2</sub>	126144.202	3305.632	1.000	10.000	10.050	3.857	-0.024	0.021	-0.240	0.030	0.242
1A1b Petroleum refining	SO <sub>2</sub>	1058.722	301.760	1.000	10.000	10.050	0.352	0.002	0.002	0.016	0.003	0.016
1A1c_ii Oil and gas extraction	SO <sub>2</sub>	4.077	10.074	1.000	10.000	10.050	0.012	0.000	0.000	0.001	0.000	0.001
1A2 Manufacturing industries and construction	SO <sub>2</sub>	15764.126	2821.617	2.000	10.000	10.198	3.341	0.013	0.018	0.126	0.052	0.136
1A4a_i Commercial / institutional	SO <sub>2</sub>	1876.067	115.838	3.000	20.000	20.224	0.272	0.000	0.001	0.001	0.003	0.003
1A4b_i Residential (excluding wood)	SO <sub>2</sub>	6192.989	552.856	3.000	20.000	20.224	1.298	0.001	0.004	0.027	0.015	0.031
1A4b_i Residential wood	SO <sub>2</sub>	98.499	343.126	20.000	20.000	28.284	1.127	0.002	0.002	0.044	0.063	0.077
1A4c_i Agriculture / forestry / fishing	SO <sub>2</sub>	3188.681	1161.404	3.000	20.000	20.224	2.727	0.006	0.008	0.127	0.032	0.131
Total	SO <sub>2</sub>	154327.363	8612.306				36.634					0.101
Total uncertainties					Overall uncertainty in the year (%):	6.053		Trend uncertainty (%):	0.318			

Continued

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data	Input data	Input data	Input data							
		Mg NO <sub>x</sub>	Mg NO <sub>x</sub>	%	%	%	%	%	%	%	%	%
1A1a Public electricity and heat production	NO <sub>x</sub>	90733.692	13800.006	1.000	20.000	20.025	8.918	-0.096	0.122	-1.916	0.172	1.923
1A1b Petroleum refining	NO <sub>x</sub>	1461.869	1300.623	1.000	20.000	20.025	0.841	0.008	0.011	0.159	0.016	0.160
1A1c_ii Oil and gas extraction	NO <sub>x</sub>	2370.571	5823.990	1.000	20.000	20.025	3.764	0.046	0.051	0.912	0.073	0.915
1A2 Manufacturing industries and construction	NO <sub>x</sub>	11985.765	5000.749	2.000	20.000	20.100	3.244	0.015	0.044	0.304	0.125	0.329
1A4a_i Commercial / institutional	NO <sub>x</sub>	1333.804	720.994	3.000	50.000	50.090	1.166	0.003	0.006	0.157	0.027	0.159
1A4b_i Residential (excluding wood)	NO <sub>x</sub>	3949.735	1420.625	3.000	50.000	50.090	2.297	0.003	0.013	0.151	0.053	0.160
1A4b_i Residential wood	NO <sub>x</sub>	538.016	2337.954	20.000	50.000	53.852	4.063	0.019	0.021	0.965	0.582	1.127
1A4c_i Agriculture / forestry / fishing	NO <sub>x</sub>	1165.704	580.667	3.000	50.000	50.090	0.939	0.002	0.005	0.116	0.022	0.118
Total	NO <sub>x</sub>	113539.157	30985.608				128.959					6.004
Total uncertainties						Overall uncertainty i the year (%):	11.356			Trend uncertainty (%):		2.450

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data	Input data	Input data	Input data							
		Mg NMVOC	Mg NMVOC	%	%	%	%	%	%	%	%	%
1A1a Public electricity and heat production	NMVOC	449.923	1305.581	1.000	50.000	50.010	4.146	0.054	0.082	2.701	0.116	2.703
1A1b Petroleum refining	NMVOC	22.855	23.106	1.000	50.000	50.010	0.073	0.000	0.001	0.002	0.002	0.003
1A1c_ii Oil and gas extraction	NMVOC	13.275	37.467	1.000	50.000	50.010	0.119	0.002	0.002	0.076	0.003	0.076
1A2 Manufacturing industries and construction	NMVOC	1099.177	260.516	2.000	50.000	50.040	0.828	-0.052	0.016	-2.590	0.046	2.590
1A4a_i Commercial / institutional	NMVOC	130.957	198.464	3.000	50.000	50.090	0.631	0.004	0.012	0.217	0.053	0.223
1A4b_i Residential (excluding wood)	NMVOC	4627.179	1994.950	3.000	50.000	50.090	6.345	-0.161	0.125	-8.067	0.531	8.085
1A4b_i Residential wood	NMVOC	7224.946	10551.110	20.000	50.000	53.852	36.081	0.213	0.662	10.654	18.729	21.547
1A4c_i Agriculture / forestry / fishing	NMVOC	2365.420	1376.640	3.000	50.000	50.090	4.379	-0.060	0.086	-3.012	0.367	3.034
Total	NMVOC	15933.732	15747.834				1379.557					552.931
Total uncertainties						Overall uncertainty i the year (%):	37.142			Trend uncertainty (%):		23.514

Continued

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the total national emissions
		Input data	Input data	Input data	Input data							
		Mg CO	Mg CO	%	%	%	%	%	%	%	%	%
1A1a Public electricity and heat production	CO	7920.045	10147.839	1.000	20.000	20.025	1.766	0.027	0.070	0.534	0.099	0.543
1A1b Petroleum refining	CO	125.940	115.426	1.000	20.000	20.025	0.020	0.000	0.001	0.002	0.001	0.002
1A1c_ii Oil and gas extraction	CO	58.790	112.403	1.000	20.000	20.025	0.020	0.000	0.001	0.009	0.001	0.009
1A2 Manufacturing industries and construction	CO	4654.926	4386.505	2.000	20.000	20.100	0.766	0.005	0.030	0.097	0.085	0.129
1A4a_i Commercial / institutional	CO	948.271	767.500	3.000	50.000	50.090	0.334	0.000	0.005	0.006	0.022	0.023
1A4b_i Residential (excluding wood)	CO	47956.385	12470.291	3.000	50.000	50.090	5.427	-0.174	0.086	-8.697	0.363	8.704
1A4b_i Residential wood	CO	51882.435	77690.708	20.000	50.000	53.852	36.352	0.251	0.533	12.552	15.085	19.624
1A4c_i Agriculture / forestry / fishing	CO	32121.056	9401.442	3.000	50.000	50.090	4.092	-0.109	0.065	-5.472	0.274	5.479
Total	CO	145667.848	115092.113				1371.446					491.209
Total uncertainties						Overall uncertainty in the year (%):	37.033			Trend uncertainty (%):		22.163

SNAP	Gas	Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the total national emissions
		Input data	Input data	Input data	Input data							
		Mg NH <sub>3</sub>	Mg NH <sub>3</sub>	%	%	%	%	%	%	%	%	%
1A1a Public electricity and heat production	NH <sub>3</sub>	0.287	16.162	1.000	1000.000	1000.000	12.404	0.024	0.025	24.454	0.036	24.454
1A1b Petroleum refining	NH <sub>3</sub>			1.000	1000.000	1000.000						
1A1c_ii Oil and gas extraction	NH <sub>3</sub>			1.000	1000.000	1000.000						
1A2 Manufacturing industries and construction	NH <sub>3</sub>			2.000	1000.000	1000.002						
1A4a_i Commercial / institutional	NH <sub>3</sub>			3.000	1000.000	1000.004						
1A4b_i Residential (excluding wood)	NH <sub>3</sub>	22.166	11.064	3.000	1000.000	1000.004	8.491	-0.054	0.017	-53.813	0.074	53.813
1A4b_i Residential wood	NH <sub>3</sub>	614.423	1275.725	20.000	1000.000	1000.200	979.301	0.029	2.003	29.097	56.656	63.691
1A4c_i Agriculture / forestry / fishing	NH <sub>3</sub>			3.000	1000.000	1000.004						
Total	NH <sub>3</sub>	636.876	1302.951				959255.494					7550.412
Total uncertainties						Overall uncertainty in the year (%):	979.416			Trend uncertainty (%):		86.893

Continued

SNAP	Gas	Base year emission (year 2000)	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Mg TSP	Input data Mg TSP	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	TSP	1020.955	930.200	1.000	50.000	50.010	2.817	-0.013	0.062	-0.668	0.088	0.673
1A1b Petroleum refining	TSP	143.931	110.100	1.000	50.000	50.010	0.333	-0.003	0.007	-0.165	0.010	0.165
1A1c_ii Oil and gas extraction	TSP	2.543	2.343	1.000	50.000	50.010	0.007	-0.000	0.000	-0.002	0.000	0.002
1A2 Manufacturing industries and construction	TSP	755.785	350.809	2.000	50.000	50.040	1.063	-0.033	0.024	-1.625	0.066	1.627
1A4a_i Commercial / institutional	TSP	164.546	147.203	3.000	50.000	50.090	0.447	-0.002	0.010	-0.117	0.042	0.124
1A4b_i Residential (excluding wood)	TSP	886.344	711.685	3.000	50.000	50.090	2.159	-0.018	0.048	-0.901	0.202	0.923
1A4b_i Residential wood	TSP	11381.787	13748.52	20.000	500.000	500.400	416.666	0.077	0.921	38.429	26.057	46.431
			2									
1A4c_i Agriculture / forestry / fishing	TSP	567.751	510.589	3.000	50.000	50.090	1.549	-0.008	0.034	-0.394	0.145	0.420
Total	TSP	14923.644	16511.45	1			173626.883					2159.969
Total uncertainties							Overall uncertainty i the year (%):				Trend uncertainty (%):	46.475

SNAP	Gas	Base year emission (year 2000)	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Mg PM <sub>10</sub>	Input data Mg PM <sub>10</sub>	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	PM <sub>10</sub>	829.352	731.255	1.000	50.000	50.010	2.366	-0.014	0.053	-0.700	0.075	0.704
1A1b Petroleum refining	PM <sub>10</sub>	130.701	103.500	1.000	50.000	50.010	0.335	-0.003	0.007	-0.153	0.011	0.153
1A1c_ii Oil and gas extraction	PM <sub>10</sub>	1.551	1.429	1.000	50.000	50.010	0.005	-0.000	0.000	-0.001	0.000	0.001
1A2 Manufacturing industries and construction	PM <sub>10</sub>	546.144	263.411	2.000	50.000	50.040	0.853	-0.025	0.019	-1.248	0.054	1.249
1A4a_i Commercial / institutional	PM <sub>10</sub>	157.542	145.368	3.000	50.000	50.090	0.471	-0.002	0.010	-0.110	0.045	0.118
1A4b_i Residential (excluding wood)	PM <sub>10</sub>	847.425	676.470	3.000	50.000	50.090	2.192	-0.019	0.049	-0.970	0.207	0.992
1A4b_i Residential wood	PM <sub>10</sub>	10811.748	13056.96	20.000	500.000	500.400	422.718	0.071	0.943	35.600	26.658	44.475
			2									
1A4c_i Agriculture / forestry / fishing	PM <sub>10</sub>	528.827	478.014	3.000	50.000	50.090	1.549	-0.008	0.035	-0.404	0.146	0.430
Total	PM <sub>10</sub>	13853.289	15456.41	0			178704.316					1981.326
Total uncertainties							Overall uncertainty i the year (%):				Trend uncertainty (%):	44.512

Continued

SNAP	Gas	Base year emission (year 2000)	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data Mg PM <sub>2.5</sub>	Input data Mg PM <sub>2.5</sub>	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	PM <sub>2.5</sub>	693.073	579.918	1.000	50.000	50.010	1.953	-0.016	0.044	-0.798	0.063	0.801
1A1b Petroleum refining	PM <sub>2.5</sub>	124.086	100.200	1.000	50.000	50.010	0.337	-0.003	0.008	-0.157	0.011	0.157
1A1c_ii Oil and gas extraction	PM <sub>2.5</sub>	1.297	1.195	1.000	50.000	50.010	0.004	-0.000	0.000	-0.001	0.000	0.001
1A2 Manufacturing industries and construction	PM <sub>2.5</sub>	305.463	181.828	2.000	50.000	50.040	0.613	-0.013	0.014	-0.635	0.039	0.636
1A4a_i Commercial / institutional	PM <sub>2.5</sub>	146.728	137.465	3.000	50.000	50.090	0.464	-0.002	0.011	-0.113	0.045	0.121
1A4b_i Residential (excluding wood)	PM <sub>2.5</sub>	812.485	644.389	3.000	50.000	50.090	2.174	-0.021	0.049	-1.072	0.209	1.092
1A4b_i Residential wood	PM <sub>2.5</sub>	10476.618	12756.911	20.000	500.000	500.400	429.898	0.064	0.977	31.856	27.644	42.178
1A4c_i Agriculture / forestry / fishing	PM <sub>2.5</sub>	492.682	447.094	3.000	50.000	50.090	1.508	-0.009	0.034	-0.434	0.145	0.458
Total	PM <sub>2.5</sub>	13052.432	14849.000				184823.859					1781.463
Total uncertainties						Overall uncertainty i the year (%):	429.911			Trend uncertainty (%):		42.207

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data kg As	Input data kg As	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	As	926.933	149.193	1.000	100.000	100.005	53.090	-0.063	0.128	-6.288	0.181	6.291
1A1b Petroleum refining	As	30.790	30.593	1.000	100.000	100.005	10.886	0.020	0.026	1.985	0.037	1.986
1A1c_ii Oil and gas extraction	As	1.128	2.787	1.000	100.000	100.005	0.992	0.002	0.002	0.215	0.003	0.215
1A2 Manufacturing industries and construction	As	156.335	74.281	2.000	100.000	100.020	26.437	0.031	0.064	3.135	0.180	3.140
1A4a_i Commercial / institutional	As	13.428	2.897	3.000	1000.000	1000.004	10.307	-0.000	0.002	-0.288	0.011	0.288
1A4b_i Residential (excluding wood)	As	14.210	6.448	3.000	1000.000	1000.004	22.942	0.003	0.006	2.592	0.023	2.592
1A4b_i Residential wood	As	1.701	5.927	20.000	1000.000	1000.200	21.093	0.005	0.005	4.727	0.144	4.729
1A4c_i Agriculture / forestry / fishing	As	22.705	8.909	3.000	1000.000	1000.004	31.699	0.003	0.008	2.948	0.032	2.948
Total	As	1167.230	281.033				5719.317					91.285
Total uncertainties						Overall uncertainty i the year (%):	75.626			Trend uncertainty (%):		9.554

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Cd	Input data kg Cd	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Cd	718.812	31.714	1.000	100.000	100.005	6.415	-0.348	0.033	-34.825	0.047	34.825
1A1b	Petroleum refining	Cd	21.271	22.445	1.000	100.000	100.005	4.540	0.012	0.023	1.197	0.033	1.197
1A1c_ii	Oil and gas extraction	Cd	0.002	0.006	1.000	100.000	100.005	0.001	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	Cd	53.971	24.210	2.000	100.000	100.020	4.898	-0.004	0.025	-0.366	0.071	0.372
1A4a_i	Commercial / institutional	Cd	30.035	0.733	3.000	1000.000	1000.004	1.483	-0.015	0.001	-15.271	0.003	15.271
1A4b_i	Residential (excluding wood)	Cd	9.745	4.150	3.000	1000.000	1000.004	8.393	-0.001	0.004	-0.891	0.018	0.891
1A4b_i	Residential wood	Cd	116.408	405.512	20.000	1000.000	1000.200	820.408	0.359	0.421	358.840	11.920	359.038
1A4c_i	Agriculture / forestry / fishing	Cd	11.984	5.610	3.000	1000.000	1000.004	11.347	-0.001	0.006	-0.569	0.025	0.569
Total		Cd	962.229	494.380				673356.462					130356.632
Total uncertainties							Overall uncertainty i the year (%):	820.583			Trend uncertainty (%):		361.049

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Cr	Input data kg Cr	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Cr	4639.758	218.432	1.000	100.000	100.005	20.175	-0.134	0.041	-13.354	0.058	13.354
1A1b	Petroleum refining	Cr	23.104	23.369	1.000	100.000	100.005	2.159	0.003	0.004	0.349	0.006	0.349
1A1c_ii	Oil and gas extraction	Cr	0.007	0.018	1.000	100.000	100.005	0.002	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	Cr	216.498	83.143	2.000	100.000	100.020	7.681	0.007	0.016	0.735	0.044	0.736
1A4a_i	Commercial / institutional	Cr	177.268	3.572	3.000	1000.000	1000.004	3.299	-0.006	0.001	-6.036	0.003	6.036
1A4b_i	Residential (excluding wood)	Cr	37.489	10.113	3.000	1000.000	1000.004	9.340	0.000	0.002	0.472	0.008	0.472
1A4b_i	Residential wood	Cr	205.952	717.445	20.000	1000.000	1000.200	662.764	0.126	0.134	126.267	3.793	126.324
1A4c_i	Agriculture / forestry / fishing	Cr	49.737	26.629	3.000	1000.000	1000.004	24.595	0.003	0.005	3.096	0.021	3.096
Total		Cr	5349.813	1082.720				440429.688					16182.880
Total uncertainties							Overall uncertainty i the year (%):	663.649			Trend uncertainty (%):		127.212

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Cu	Input data kg Cu	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Cu	2870.506	196.307	1.000	100.000	100.005	32.207	-0.083	0.055	-8.261	0.078	8.262
1A1b	Petroleum refining	Cu	45.203	45.395	1.000	100.000	100.005	7.448	0.011	0.013	1.059	0.018	1.059
1A1c_ii	Oil and gas extraction	Cu	0.001	0.002	1.000	100.000	100.005	0.000	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	Cu	306.940	106.285	2.000	100.000	100.020	17.440	0.015	0.030	1.508	0.085	1.510
1A4a_i	Commercial / institutional	Cu	122.640	3.838	3.000	1000.000	1000.004	6.296	-0.005	0.001	-4.836	0.005	4.836
1A4b_i	Residential (excluding wood)	Cu	69.877	26.427	3.000	1000.000	1000.004	43.355	0.004	0.007	4.063	0.032	4.063
1A4b_i	Residential wood	Cu	53.727	187.159	20.000	1000.000	1000.200	307.107	0.050	0.053	50.059	1.489	50.081
1A4c_i	Agriculture / forestry / fishing	Cu	85.290	44.136	3.000	1000.000	1000.004	72.408	0.008	0.012	8.301	0.053	8.301
Total		Cu	3554.183	609.549				102873.877					2688.564
Total uncertainties							Overall uncertainty i the year (%):	320.740			Trend uncertainty (%):		51.851

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Hg	Input data kg Hg	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Hg	2448.434	170.976	1.000	100.000	100.005	56.444	-0.032	0.060	-3.193	0.085	3.194
1A1b	Petroleum refining	Hg	20.093	21.851	1.000	100.000	100.005	7.214	0.007	0.008	0.696	0.011	0.696
1A1c_ii	Oil and gas extraction	Hg	0.948	2.342	1.000	100.000	100.005	0.773	0.001	0.001	0.079	0.001	0.079
1A2	Manufacturing industries and construction	Hg	192.046	69.725	2.000	100.000	100.020	23.022	0.017	0.025	1.736	0.070	1.738
1A4a_i	Commercial / institutional	Hg	124.038	2.284	3.000	1000.000	1000.004	7.541	-0.004	0.001	-3.884	0.003	3.884
1A4b_i	Residential (excluding wood)	Hg	16.349	5.177	3.000	1000.000	1000.004	17.089	0.001	0.002	1.211	0.008	1.211
1A4b_i	Residential wood	Hg	5.014	17.468	20.000	1000.000	1000.200	57.676	0.006	0.006	5.983	0.175	5.986
1A4c_i	Agriculture / forestry / fishing	Hg	22.750	13.106	3.000	1000.000	1000.004	43.263	0.004	0.005	3.771	0.020	3.771
Total		Hg	2829.672	302.929				9315.643					80.313
Total uncertainties							Overall uncertainty i the year (%):	96.518			Trend uncertainty (%):		8.962



SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the total national emissions	
		Input data kg Ni	Input data kg Ni	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Ni	8038.635	375.149	1.000	100.000	100.005	17.737	-0.039	0.023	-3.896	0.032	3.896
1A1b	Petroleum refining	Ni	493.501	260.490	1.000	100.000	100.005	12.316	0.012	0.016	1.190	0.022	1.191
1A1c_ii	Oil and gas extraction	Ni	0.005	0.012	1.000	100.000	100.005	0.001	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	Ni	6274.997	1292.041	2.000	100.000	100.020	61.098	0.030	0.078	2.955	0.220	2.963
1A4a_i	Commercial / institutional	Ni	587.203	25.334	3.000	1000.000	1000.004	11.977	-0.003	0.002	-2.983	0.006	2.983
1A4b_i	Residential (excluding wood)	Ni	386.269	28.786	3.000	1000.000	1000.004	13.610	-0.001	0.002	-1.232	0.007	1.232
1A4b_i	Residential wood	Ni	17.909	62.386	20.000	1000.000	1000.200	29.502	0.004	0.004	3.622	0.106	3.624
1A4c_i	Agriculture / forestry / fishing	Ni	793.891	70.912	3.000	1000.000	1000.004	33.527	-0.002	0.004	-1.825	0.018	1.825
Total		Ni	16592.410	2115.111				6522.391					52.259
Total uncertainties							Overall uncertainty i the year (%):	80.761			Trend uncertainty (%):		7.229

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the total national emissions	
		Input data kg Pb	Input data kg Pb	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Pb	12005.915	390.257	1.000	100.000	100.005	18.255	-0.084	0.026	-8.391	0.036	8.391
1A1b	Petroleum refining	Pb	63.725	66.454	1.000	100.000	100.005	3.108	0.004	0.004	0.377	0.006	0.377
1A1c_ii	Oil and gas extraction	Pb	0.014	0.035	1.000	100.000	100.005	0.002	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	Pb	1483.940	437.215	2.000	100.000	100.020	20.454	0.015	0.029	1.501	0.081	1.503
1A4a_i	Commercial / institutional	Pb	679.141	5.209	3.000	1000.000	1000.004	2.436	-0.006	0.000	-5.886	0.001	5.886
1A4b_i	Residential (excluding wood)	Pb	308.283	118.469	3.000	1000.000	1000.004	55.413	0.005	0.008	4.931	0.033	4.931
1A4b_i	Residential wood	Pb	241.770	842.217	20.000	1000.000	1000.200	394.017	0.053	0.055	52.942	1.560	52.965
1A4c_i	Agriculture / forestry / fishing	Pb	483.468	278.085	3.000	1000.000	1000.004	130.072	0.014	0.018	13.776	0.077	13.776
Total		Pb	15266.256	2137.941				176006.161					3126.897
Total uncertainties							Overall uncertainty i the year (%):	419.531			Trend uncertainty (%):		55.919

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Se	Input data kg Se	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Se	3453.494	730.769	1.000	100.000	100.005	74.989	-0.031	0.185	-3.054	0.262	3.065
1A1b	Petroleum refining	Se	97.931	106.305	1.000	100.000	100.005	10.909	0.021	0.027	2.080	0.038	2.080
1A1c_ii	Oil and gas extraction	Se	0.095	0.234	1.000	100.000	100.005	0.024	0.000	0.000	0.005	0.000	0.005
1A2	Manufacturing industries and construction	Se	276.929	79.923	2.000	100.000	100.020	8.203	0.003	0.020	0.293	0.057	0.298
1A4a_i	Commercial / institutional	Se	27.765	1.291	3.000	1000.000	1000.004	1.324	-0.001	0.000	-1.409	0.001	1.409
1A4b_i	Residential (excluding wood)	Se	19.855	1.855	3.000	1000.000	1000.004	1.903	-0.001	0.000	-0.771	0.002	0.771
1A4b_i	Residential wood	Se	4.477	15.597	20.000	1000.000	1000.200	16.007	0.004	0.004	3.670	0.112	3.672
1A4c_i	Agriculture / forestry / fishing	Se	67.852	38.581	3.000	1000.000	1000.004	39.588	0.006	0.010	5.529	0.041	5.529
Total		Se	3948.398	974.553				7638.435					60.441
Total uncertainties							Overall uncertainty i the year (%):	87.398			Trend uncertainty (%):		7.774

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Zn	Input data kg Zn	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Zn	16298.113	552.277	1.000	100.000	100.005	2.975	-0.373	0.020	-37.302	0.028	37.302
1A1b	Petroleum refining	Zn	148.226	89.791	1.000	100.000	100.005	0.484	-0.000	0.003	-0.035	0.005	0.035
1A1c_ii	Oil and gas extraction	Zn	0.014	0.035	1.000	100.000	100.005	0.000	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	Zn	3706.273	982.569	2.000	100.000	100.020	5.294	-0.054	0.036	-5.428	0.100	5.429
1A4a_i	Commercial / institutional	Zn	866.574	14.770	3.000	1000.000	1000.004	0.796	-0.020	0.001	-20.471	0.002	20.471
1A4b_i	Residential (excluding wood)	Zn	927.407	388.416	3.000	1000.000	1000.004	20.923	-0.008	0.014	-8.447	0.060	8.447
1A4b_i	Residential wood	Zn	4584.669	15970.939	20.000	1000.000	1000.200	860.473	0.465	0.577	465.254	16.325	465.541
1A4c_i	Agriculture / forestry / fishing	Zn	1138.940	565.558	3.000	1000.000	1000.004	30.465	-0.007	0.020	-7.173	0.087	7.174
Total		Zn	27670.217	18564.356				741817.967					218690.871
Total uncertainties							Overall uncertainty i the year (%):	861.289			Trend uncertainty (%):		467.644

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data g Dioxin	Input data g Dioxin	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Dioxin	30.911	1.163	1.000	500.000	500.001	37.881	-0.189	0.025	-94.628	0.035	94.628
1A1b	Petroleum refining	Dioxin	0.002	0.001	1.000	1000.000	1000.000	0.063	0.000	0.000	0.010	0.000	0.010
1A1c_ii	Oil and gas extraction	Dioxin	0.000	0.001	1.000	1000.000	1000.000	0.038	0.000	0.000	0.011	0.000	0.011
1A2	Manufacturing industries and construction	Dioxin	0.904	0.055	2.000	1000.000	1000.002	3.572	-0.005	0.001	-5.126	0.003	5.126
1A4a_i	Commercial / institutional	Dioxin	2.189	0.411	3.000	1000.000	1000.004	26.734	-0.007	0.009	-6.501	0.037	6.501
1A4b_i	Residential (excluding wood)	Dioxin	4.297	1.580	3.000	1000.000	1000.004	102.872	0.004	0.034	3.713	0.143	3.716
1A4b_i	Residential wood	Dioxin	6.234	10.832	20.000	1000.000	1000.200	705.583	0.187	0.231	187.029	6.525	187.143
1A4c_i	Agriculture / forestry / fishing	Dioxin	2.419	1.313	3.000	1000.000	1000.004	85.518	0.011	0.028	11.112	0.119	11.113
Total		Dioxin	46.955	15.355				517905.546					44182.926
Total uncertainties						Overall uncertainty i the year (%):		719.657			Trend uncertainty (%):		210.197

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg Benzo(b)	Input data kg Benzo(b)	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	Benzo(b)	22.623	35.777	1.000	100.000	100.005	1.670	0.001	0.025	0.117	0.036	0.123
1A1b	Petroleum refining	Benzo(b)	0.622	0.314	1.000	100.000	100.005	0.015	-0.000	0.000	-0.044	0.000	0.044
1A1c_ii	Oil and gas extraction	Benzo(b)	0.009	0.023	1.000	100.000	100.005	0.001	0.000	0.000	0.001	0.000	0.001
1A2	Manufacturing industries and construction	Benzo(b)	48.987	71.757	2.000	100.000	100.020	3.350	-0.001	0.051	-0.148	0.143	0.206
1A4a_i	Commercial / institutional	Benzo(b)	52.109	202.299	3.000	1000.000	1000.004	94.434	0.087	0.142	87.052	0.604	87.054
1A4b_i	Residential (excluding wood)	Benzo(b)	121.877	38.542	3.000	1000.000	1000.004	17.992	-0.102	0.027	-102.109	0.115	102.109
1A4b_i	Residential wood	Benzo(b)	943.889	1628.037	20.000	1000.000	1000.200	760.125	0.143	1.146	143.293	32.408	146.912
1A4c_i	Agriculture / forestry / fishing	Benzo(b)	230.761	165.479	3.000	1000.000	1000.004	77.247	-0.128	0.116	-128.188	0.494	128.189
Total		Benzo(b)	1420.877	2142.229				593013.265					56020.363
Total uncertainties						Overall uncertainty i the year (%):		770.074			Trend uncertainty (%):		236.686

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data kg Benzo(k)	Input data kg Benzo(k)	Input data %	Input data %	%	%	%	%	%	%	%
1A1a	Public electricity and heat production	Benzo(k) 10.406	23.119	1.000	100.000	100.005	3.076	0.014	0.048	1.422	0.068	1.424
1A1b	Petroleum refining	Benzo(k) 0.122	0.062	1.000	100.000	100.005	0.008	-0.000	0.000	-0.027	0.000	0.027
1A1c_ii	Oil and gas extraction	Benzo(k) 0.019	0.047	1.000	100.000	100.005	0.006	0.000	0.000	0.004	0.000	0.004
1A2	Manufacturing industries and construction	Benzo(k) 22.692	13.878	2.000	100.000	100.020	1.846	-0.045	0.029	-4.502	0.082	4.503
1A4a_i	Commercial / institutional	Benzo(k) 16.050	67.142	3.000	1000.000	1000.004	89.317	0.087	0.140	87.459	0.593	87.461
1A4b_i	Residential (excluding wood)	Benzo(k) 39.727	20.439	3.000	1000.000	1000.004	27.190	-0.087	0.043	-86.814	0.181	86.814
1A4b_i	Residential wood	Benzo(k) 356.368	595.293	20.000	1000.000	1000.200	792.053	0.078	1.239	77.633	35.054	85.180
1A4c_i	Agriculture / forestry / fishing	Benzo(k) 34.947	31.753	3.000	1000.000	1000.004	42.240	-0.048	0.066	-47.725	0.280	47.726
Total		Benzo(k) 480.331	751.733				637861.865					24741.790
Total uncertainties						Overall uncertainty i the year (%):	798.663			Trend uncertainty (%):		157.295

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data kg Benzo(a)	Input data kg Benzo(a)	Input data %	Input data %	%	%	%	%	%	%	%
1A1a	Public electricity and heat production	Benzo(a) 6.882	8.884	1.000	100.000	100.005	0.444	-0.001	0.006	-0.062	0.009	0.063
1A1b	Petroleum refining	Benzo(a) 0.143	0.072	1.000	100.000	100.005	0.004	-0.000	0.000	-0.009	0.000	0.009
1A1c_ii	Oil and gas extraction	Benzo(a) 0.009	0.023	1.000	100.000	100.005	0.001	0.000	0.000	0.001	0.000	0.001
1A2	Manufacturing industries and construction	Benzo(a) 10.801	19.744	2.000	100.000	100.020	0.988	0.003	0.014	0.314	0.040	0.317
1A4a_i	Commercial / institutional	Benzo(a) 40.998	154.008	3.000	1000.000	1000.004	77.043	0.068	0.109	67.982	0.463	67.984
1A4b_i	Residential (excluding wood)	Benzo(a) 116.485	39.069	3.000	1000.000	1000.004	19.544	-0.089	0.028	-89.334	0.118	89.334
1A4b_i	Residential wood	Benzo(a) 1023.422	1628.422	20.000	1000.000	1000.200	814.785	0.125	1.155	125.003	32.665	129.201
1A4c_i	Agriculture / forestry / fishing	Benzo(a) 211.305	148.769	3.000	1000.000	1000.004	74.422	-0.107	0.106	-106.782	0.448	106.783
Total		Benzo(a) 1410.046	1998.991				675731.957					40697.920
Total uncertainties						Overall uncertainty i the year (%):	822.029			Trend uncertainty (%):		201.737

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data kg Indeno	Input data kg Indeno	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	Indeno	6.060	6.389	1.000	100.000	100.005	0.506	-0.001	0.006	-0.053	0.008	0.053
1A1b Petroleum refining	Indeno	0.232	0.117	1.000	100.000	100.005	0.009	-0.000	0.000	-0.014	0.000	0.014
1A1c_ii Oil and gas extraction	Indeno	0.028	0.070	1.000	100.000	100.005	0.006	0.000	0.000	0.003	0.000	0.003
1A2 Manufacturing industries and construction	Indeno	13.576	3.445	2.000	100.000	100.020	0.273	-0.011	0.003	-1.107	0.009	1.107
1A4a_i Commercial / institutional	Indeno	37.086	109.541	3.000	1000.000	1000.004	86.690	0.061	0.100	60.859	0.423	60.861
1A4b_i Residential (excluding wood)	Indeno	117.831	15.385	3.000	1000.000	1000.004	12.176	-0.109	0.014	-109.220	0.059	109.220
1A4b_i Residential wood	Indeno	599.139	918.804	20.000	1000.000	1000.200	727.280	0.208	0.836	207.973	23.653	209.314
1A4c_i Agriculture / forestry / fishing	Indeno	324.751	209.845	3.000	1000.000	1000.004	166.070	-0.149	0.191	-148.504	0.810	148.507
Total	Indeno	1098.704	1263.596				564178.886					81500.733
Total uncertainties						Overall uncertainty i the year (%):	751.118			Trend uncertainty (%):		285.483

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncertainty	Uncertainty in trend in national emissions introduced by activity data uncertainty	Uncertainty introduced into the trend in total national emissions
		Input data kg HCB	Input data kg HCB	Input data %	Input data %	%	%	%	%	%	%	%
1A1a Public electricity and heat production	HCB	4.369	1.246	1.000	1000.000	1000.000	795.347	0.012	0.214	11.721	0.303	11.725
1A1b Petroleum refining	HCB	0.000	0.000	1.000	1000.000	1000.000	0.093	0.000	0.000	0.012	0.000	0.012
1A1c_ii Oil and gas extraction	HCB		0.000	1.000	1000.000	1000.000	0.000	0.000	0.000	0.000	0.000	0.000
1A2 Manufacturing industries and construction	HCB	0.140	0.075	2.000	1000.000	1000.002	47.816	0.006	0.013	6.403	0.036	6.403
1A4a_i Commercial / institutional	HCB	0.181	0.007	3.000	1000.000	1000.004	4.307	-0.007	0.001	-7.225	0.005	7.225
1A4b_i Residential (excluding wood)	HCB	0.932	0.038	3.000	1000.000	1000.004	24.166	-0.037	0.007	-36.652	0.028	36.652
1A4b_i Residential wood	HCB	0.045	0.156	20.000	1000.000	1000.200	99.542	0.025	0.027	24.748	0.759	24.760
1A4c_i Agriculture / forestry / fishing	HCB	0.146	0.045	3.000	1000.000	1000.004	28.750	0.001	0.008	0.962	0.033	0.963
Total	HCB	5.814	1.567				646200.855					2188.043
Total uncertainties						Overall uncertainty i the year (%):	803.866			Trend uncertainty (%):		46.777

Continued

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data kg PCB	Input data kg PCB	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	PCB	0.893	0.283	1.000	1000.000	1000.000	625.659	-0.051	0.240	-50.515	0.340	50.516
1A1b	Petroleum refining	PCB	0.001	0.001	1.000	1000.000	1000.000	1.225	0.000	0.000	0.112	0.001	0.112
1A1c_ii	Oil and gas extraction	PCB		0.000	1.000	1000.000	1000.000	0.000	0.000	0.000	0.000	0.000	0.000
1A2	Manufacturing industries and construction	PCB	0.121	0.052	2.000	1000.000	1000.002	115.888	0.005	0.044	5.057	0.126	5.058
1A4a_i	Commercial / institutional	PCB	0.045	0.003	3.000	1000.000	1000.004	6.526	-0.012	0.003	-12.181	0.011	12.181
1A4b_i	Residential (excluding wood)	PCB	0.027	0.010	3.000	1000.000	1000.004	21.533	-0.000	0.008	-0.497	0.035	0.498
1A4b_i	Residential wood	PCB	0.060	0.086	20.000	1000.000	1000.200	189.885	0.053	0.073	53.450	2.061	53.490
1A4c_i	Agriculture / forestry / fishing	PCB	0.031	0.018	3.000	1000.000	1000.004	39.322	0.005	0.015	4.928	0.064	4.928
Total		PCB	1.178	0.452				442990.128					5611.538
Total uncertainties							Overall uncertainty i the year (%):	665.575			Trend uncertainty (%):		74.910

SNAP		Base year emission	Year t emission	Activity data uncertainty	Emission factor uncertainty	Combined uncertainty	Combined uncertainty as % of total national emissions in year t	Type A sensitivity	Type B sensitivity	Uncertainty in trend in national emissions introduced by emission factor uncer- tainty	Uncertainty in trend in national emissions introduced by activity data uncer- tainty	Uncertainty introduced into the trend in total national emissions	
		Input data Gg BC	Input data Gg BC	Input data %	Input data %	%	%	%	%	%	%	%	
1A1a	Public electricity and heat production	BC	0.570	0.283	1.000	1000.000	1000.000	149.606	-0.117	0.169	-116.800	0.239	116.801
1A1b	Petroleum refining	BC	0.020	0.001	1.000	1000.000	1000.000	7.874	-0.001	0.009	-1.146	0.013	1.146
1A1c_ii	Oil and gas extraction	BC		0.000	1.000	1000.000	1000.000						
1A2	Manufacturing industries and construction	BC	0.090	0.052	2.000	1000.000	1000.002	31.496	-0.010	0.036	-9.596	0.101	9.597
1A4a_i	Commercial / institutional	BC	0.060	0.003	3.000	1000.000	1000.004	15.748	-0.012	0.018	-12.323	0.075	12.323
1A4b_i	Residential (excluding wood)	BC	0.070	0.010	3.000	1000.000	1000.004	23.622	-0.008	0.027	-8.452	0.113	8.452
1A4b_i	Residential wood	BC	1.300	0.086	20.000	1000.000	1000.200	724.554	0.165	0.818	164.580	23.130	166.197
1A4c_i	Agriculture / forestry / fishing	BC	0.140	0.018	3.000	1000.000	1000.004	47.244	-0.017	0.053	-16.898	0.226	16.900
Total		BC	2.250	2.540				551453.064					41866.266
Total uncertainties							Overall uncertainty i the year (%):	742.599			Trend uncertainty (%):		204.612

## Annex 3A-8 Emission inventory 2013 based on SNAP sectors

Table 3A-8.1 Emission inventory 2013 based on SNAP sectors.

SNAP	SO2 [Mg]	NOX [Mg]	NMVOG [Mg]	CO [Mg]	NH3 [Mg]	TSP [Mg]	PM10 [Mg]	PM2.5 [Mg]	As [kg]	Cd [kg]	Cr [kg]	Cu [kg]	Hg [kg]	Ni [kg]	Pb [kg]	Se [kg]	Zn [kg]	HCb [kg]	DIOX [g]	Benzo(b) [kg]	Benzo(k) [kg]	Benzo(a) [kg]	Indeno [kg]	PCBs [kg]
10100	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10101	1564.74	4270.36	203.03	2050.95	0	483.08	418.08	337.13	89.56	10.28	107.9	84.98	95.41	147.29	99.66	625.75	324.01	0.89	0.31	1	0.24	0.28	0.37	0.13
10102	510.67	2797.94	67.11	996.08	4.85	49.36	43.3	32.29	16.44	7.19	34.31	35.01	26.25	112.45	91.6	33.38	92.14	0.14	0.29	0.36	0.1	0.16	0.1	0.04
10103	159.85	1357.07	11.41	272.25	7.76	21.92	17.19	12.29	11.52	5.79	24.01	24.48	26.67	49.18	86.46	13.93	30.26	0.05	0.14	0.1	0.03	0.04	0.03	0.01
10104	68.34	975.24	94.24	505.83	0.94	15.1	11.39	7.81	3.93	2.12	10.18	7.2	8.2	10.34	21.11	44.31	30.76	0.09	0.26	0.29	0.11	0.18	0.07	0.05
10105	55.31	1728.26	777.21	1479.21	0	14.08	3.23	2.29	0.61	0.05	0.93	0.93	1.34	9.47	0.42	0.71	37.26	0	0.01	0.09	0.02	0.02	0.02	0
10200	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10201	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10202	4.5	12.47	0.21	6.65	0	1.01	1.01	1.01	0	0	0.04	0.03	0.03	0	0	0	0.08	0	0	0.1	0.02	0.02	0.04	0
10203	942.21	2658.67	152.36	4836.86	2.61	345.66	237.06	187.11	27.14	6.29	41.06	43.68	13.06	46.43	91.01	12.69	37.76	0.07	0.15	33.85	22.6	8.17	5.75	0.05
10204	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10205	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10300	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10301	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10302	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10303	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10304	6.87	282.95	2.49	11.02	0	8.89	8.89	8.89	3.2	2.49	2.49	4.8	2.49	2.49	7.29	12.09	3.2	0	0	0	0	0	0	0
10305	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10306	294.89	1017.67	20.62	104.4	0	101.21	94.61	91.31	27.39	19.96	20.88	40.6	19.36	25.8	59.17	94.22	86.59	0	0.31	0.06	0.07	0.12	0	
10400	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10401	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10402	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10403	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10404	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10405	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10406	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10407	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10500	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10501	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10502	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10503	0.06	4.66	0.22	0.67	0	0.01	0.01	0.01	0.02	0	0	0	0.01	0	0	0	0	0	0	0	0	0	0	0
10504	10.01	5819.33	37.24	111.73	0	2.33	1.42	1.19	2.77	0.01	0.02	0	2.33	0.01	0.03	0.23	0.03	0	0	0.02	0.05	0.02	0.07	0
10505	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10506	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20100	102.52	556.36	166.1	572.24	0	145.31	144.87	137.12	2.79	0.43	3.26	3.5	2.08	24.95	4.89	1.04	11.36	0.01	0.41	202.29	67.14	154	109.54	0
20101	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20102	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20103	2.31	10.29	0.47	2.82	0	0.18	0.18	0.18	0.07	0.3	0.19	0.16	0.11	0.24	0.3	0.13	0.34	0	0	0	0	0	0	0
20104	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20105	11.01	154.34	31.89	192.44	0	1.71	0.31	0.17	0.04	0	0.12	0.18	0.1	0.14	0.01	0.12	3.07	0	0	0	0	0	0	0
20106	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20200	895.64	3693.81	12503.37	90122.52	1286.64	14458.21	13731.77	13399.68	12.33	409.61	727.45	213.56	22.58	91.14	960.57	17.44	16356.16	0.19	12.41	1666.38	615.66	1667.3	934.08	0.1
20201	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20202	0.12	5	1.96	12.8	0.15	1.66	1.58	1.54	0.02	0.05	0.09	0.02	0.02	0.01	0.1	0	1.91	0	0	0.19	0.07	0.19	0.11	0
20203	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20204	0.22	59.76	40.73	25.68	0	0.34	0.08	0.07	0.02	0	0.02	0	0.04	0.02	0.02	0	1.28	0	0	0	0	0	0	0
20205	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20300	1148.05	398.41	1322.88	9195.11	0	507.91	476.94	446.25	8.84	5.58	26.44	43.79	12.98	70.67	277.27	38.45	559.34	0.04	1.3	164.55	31.39	148.01	209.4	0.02
20301	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20302	2.62	2.11	0.64	7.14	0	0.85	0.73	0.65	0.02	0.03	0.06	0.18	0.01	0.09	0.8	0.01	2.56	0	0.01	0.92	0.36	0.76	0.44	0
20303	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20304	10.73	180.15	53.12	199.19	0	1.83	0.34	0.2	0.05	0	0.12	0.17	0.12	0.15	0.02	0.12	3.66	0	0	0.01	0	0	0	0
20305	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30100	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30101	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30102	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30103	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30104	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30105	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

SNAP	SO2 [Mg]	NOX [Mg]	NMVOG [Mg]	CO [Mg]	NH3 [Mg]	TSP [Mg]	PM10 [Mg]	PM2.5 [Mg]	As [kg]	Cd [kg]	Cr [kg]	Cu [kg]	Hg [kg]	Ni [kg]	Pb [kg]	Se [kg]	Zn [kg]	HCB [kg]	DIOX [g]	Benzo(b) [kg]	Benzo(k) [kg]	Benzo(a) [kg]	Indeno [kg]	PCBs [kg]
30106	0.01	0.85	0.04	0.57	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30200	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30203	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30204	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30205	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30400	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30401	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30402	0.53	150.25	2.46	34.31	0	0.12	0.12	0.12	0.15	0	0	0	0.12	0	0	0.01	0	0	0	0	0	0	0	0
30403	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30404	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30405	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30406	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30500	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30501	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30502	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30503	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30504	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30505	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30506	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30600	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30601	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30602	0.32	4.75	1.35	18.74	0	0	0	0	0.08	0	0	0	0.07	0	0	0.01	0	0	0	0	0	0	0	0
30603	9	32.4	1.01	1.54	0	1.82	1.36	0.92	0.83	0.25	0.52	1.09	0.08	69.12	0.94	0.42	17.95	0	0	0.01	0.01	0.02	0.03	0
30604	0	100.2	1.27	3.82	0	0.08	0.05	0.04	0.09	0	0	0	0.08	0	0	0.01	0	0	0	0	0	0	0	0
30605	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30606	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30700	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30701	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30702	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30703	0.06	60.59	2.44	2.59	0	4.13	2.92	1.7	0.97	0.44	3.28	4.25	1.92	3.16	32.57	6.08	48.61	0	0	0.23	0.23	0.01	0.17	0
30704	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30705	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30706	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30800	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30801	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30802	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30803	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30804	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30805	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30806	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30900	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30901	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30902	703.26	343.4	6.88	21.38	0	34.69	27.53	18.97	10.08	3.62	17.02	25.21	8.25	532.8	136.96	26.32	329.17	0.01	0	0.98	1.02	0.14	0.92	0.01
30903	296.74	332	4.18	7.8	0	61.42	57.47	45.21	6.21	2.07	7.4	12.13	2.98	419.97	48.26	8.77	171.77	0	0	0.34	0.37	0.1	0.41	0
30904	0.01	0.11	0.04	0.13	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30905	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30906	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31001	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31002	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31003	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31004	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31100	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31101	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31102	4.99	40.9	4.54	108.94	0	8.62	5.9	4.54	0.64	0.12	1.06	1.18	0.18	1.06	1.64	0.23	1.04	0	0	0.59	0.59	0	0.01	0
31103	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31104	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31105	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31106	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31200	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31201	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0



SNAP	SO2 [Mg]	NOX [Mg]	NMVOG [Mg]	CO [Mg]	NH3 [Mg]	TSP [Mg]	PM10 [Mg]	PM2.5 [Mg]	As [kg]	Cd [kg]	Cr [kg]	Cu [kg]	Hg [kg]	Ni [kg]	Pb [kg]	Se [kg]	Zn [kg]	HCb [kg]	DIOX [g]	Benzo(b) [kg]	Benzo(k) [kg]	Benzo(a) [kg]	Indeno [kg]	PCBs [kg]	
31202	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
31203	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31204	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31205	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31206	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31300	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31301	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31302	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31303	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31304	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31305	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31306	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31400	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31401	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31402	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31403	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31404	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31405	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31406	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31500	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31501	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31502	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31503	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31504	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31505	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31506	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31600	587	1401	75.65	1678	0	81	57.85	26.91	36.5	12.78	18.25	18.25	40	36.5	18.25	12.78	91.26	0.02	0.04	58.56	1.11	19.19	0.85	0.01	
31601	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31602	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31603	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31604	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31605	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31606	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32000	0.44	225.08	1.69	2.81	0	0	0	0	0.02	0	0	0	0.1	0	0	0	0	0	0	0	0	0	0	0	0
32001	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32002	489.35	135	5.53	0	0	0	0	0	0	0.93	6.95	9.01	4.09	6.7	69.01	3.08	103	0	0	0.48	0.48	0.01	0.36	0	
32003	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32004	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32005	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32006	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

## **Annex 3A-9 Description of the Danish energy statistics**

This description of the Danish energy statistics has been prepared by Denmark's National Environmental Research Institute, NERI (now DCE) in cooperation with the Danish Energy Agency (DEA) as background information to the Danish National Inventory Report (NIR).

### **The Danish energy statistics system**

DEA is responsible for the Danish energy balance. Main contributors to the energy statistics outside DEA are Statistics Denmark and Danish Energy Association (before Association of Danish Energy Companies). The statistics is performed using an integrated statistical system building on an Access database and Excel spreadsheets.

The DEA follows the recommendations of the International Energy Agency as well as Eurostat.

The national energy statistics is updated annually and all revisions are immediately included in the published statistics, which can be found on the DEA homepage. It is an easy task to check for breaks in a series because the statistics is 100 % time-series oriented.

The national energy statistics does not include Greenland and Faroe Islands.

For historical reasons, DEA receive monthly information from the Danish oil companies regarding Danish deliveries of oil products to Greenland and Faroe Islands. However, the monthly (MOS) and annual (AOS) reporting of oil statistics to Eurostat and IEA exclude Greenland and Faroe Islands. For all other energy products, the Danish figures are also excluding Greenland and Faroe Islands.

### **Reporting to the Danish Energy Agency**

The Danish Energy Agency receives monthly statistics for the following fuel groups:

- Crude oil and oil products.
  - Monthly data from 46 oil companies, the main purpose is monitoring oil stocks according to the oil preparedness system.
- Natural gas.
  - Fuel/flare from platforms in the North Sea.
  - Natural gas balance from the regulator Energinet.dk (National monopoly).
- Coal and coke.
  - Power plants (94 %).
  - Industry companies (4 %).
  - Coal and coke traders (2 %).
- Electricity.
  - Monthly reporting by e-mail from the regulator Energinet.dk (National monopoly).
  - The statistics covers:
    - Production by type of producer.
    - Own use of electricity.
    - Import and export by country.
    - Domestic supply (consumption + distribution loss).
- Town gas (quarterly) from two town gas producers.

- The large central power plants also report monthly consumption of biomass.

Annual data includes renewable energy including waste. The DEA conducts a biannual survey on wood pellets and wood fuel. Statistics Denmark conducts biannual surveys on the energy consumption in the service and industrial sectors. Statistics Denmark prepares annual surveys on forest (wood fuel) & straw.

Other annual data sources include:

- DEA:
  - Survey on production of electricity and heat and fuels used.
  - Survey on end use of oil.
  - Survey on end use of natural gas.
  - Survey on end use of coal and coke.
- DCE (former NERI), Aarhus University.
  - Energy consumption for domestic air transport.
- Danish Energy Association (Association of Danish Energy companies).
  - Survey on electricity consumption.
- Ministry of Taxation.
  - Border trade.
- Centre for Biomass Technology.
  - Annual estimates of final consumption of straw and wood chips.

### **Annual revisions**

In general, DEA follows the same procedures as in the Danish national account. This means that normally only figures for the last two years are revised.

### **Aggregating the energy statistics on SNAP level**

The sectors used in the official energy statistics have been mapped to SNAP categories, used in the Danish emission database. DCE aggregates the official energy statistics to SNAP level based on a source correspondence table.

In cooperation between DEA and DCE, a fuel correspondence table has been developed mapping the fuels used by the DEA in the official energy statistics with the fuel codes used in the Danish national emission database. The fuel correspondence table between fuel categories used by the DEA, DCE and NFR is presented in Annex 3A-3.

The mapping between the energy statistics and the SNAP and fuel codes used by DCE can be seen in the table below.

Table 3A-9.1 Correspondence between the Danish national energy statistics and the SNAP nomenclature (only stationary combustion part shown).

Unit: TJ	End-use		Transformation 1980-1993	
	SNAP	Fuel ( <i>in Danish</i> )	SNAP	Fuel-code
<b>Foreign Trade</b>				
- <i>Border Trade</i>				
	0202	Petrokoks		110A
<b>Vessels in Foreign Trade</b>				
- <i>International Marine Bunkers</i>				
<b>Energy Sector</b>				
<b>Extraction and Gasification</b>				
- <i>Extraction</i>				
	010504	Naturgas		301A
- <i>Gasification</i>				
	091006	Biogas		309A
	091006	Biogas		309A
<b>Refineries</b>				
- Own Use				
	010306	Raffinaderigas		308A
	010306	LPG		303A
	010306	Gas & Dieselolie		204A
	010306	Fuelolie & Spildolie		203A
<b>Transformation Sector</b>				
<b>Large-scale Power Units</b>				
- <i>Fuels Used for Power Production</i>				
			0101	204A
			0101	203A
			0101	102A
			0101	117A
<b>Large-Scale CHP Units</b>				
- <i>Fuels Used for Power Production</i>				
			0103	308A
			0101	303A
			0101	210A
			0101	204A
			0101	203A
			0101	110A
			0101	225A
			0101	301A
			0101	102A
			0101	117A
			0101	111A
			0101	111A
			0101	111A
			0101	309A
			0101	309A
			0101	114A
			0101	114A
- <i>Fuels Used for Heat Production</i>				
			0103	308A
			0101	303A
			0101	210A
			0101	204A
			0101	203A
			0101	110A
			0101	225A
			0101	301A
			0101	102A
			0101	117A
			0101	111A
			0101	111A
			0101	111A
			0101	309A
			0101	309A
			0101	114A
			0101	114A
<b>Small-Scale CHP Units</b>				

Unit: TJ	End-use		Transformation 1980-1993		
	SNAP	Fuel (in Danish)	Fuel-code	SNAP	Fuel-code
<i>- Fuels Used for Power Production</i>					
- - Gas-/Diesel Oil				0101	204A
- - Fuel Oil				0101	203A
- - Natural Gas				0101	301A
- - Hard Coal				0101	102A
- - Straw				0101	117A
- - Wood Chips				0101	111A
- - Wood Pellets				0101	111A
- - Wood Waste				0101	111A
- - Biogas, Landfill				0101	309A
- - Biogas, Other				0101	309A
- - Waste, Non-renewable				0101	114A
- - Wastes, Renewable				0101	114A
<i>- Fuels Used for Heat Production</i>					
- - Gas-/Diesel Oil				0101	204A
- - Fuel Oil				0101	203A
- - Natural Gas				0101	301A
- - Coal				0101	102A
- - Straw				0101	117A
- - Wood Chips				0101	111A
- - Wood Pellets				0101	111A
- - Wood Waste				0101	111A
- - Biogas, Landfill				0101	309A
- - Biogas, Other				0101	309A
- - Waste, Non-renewable				0101	114A
- - Wastes, Renewable				0101	114A
<b>District Heating Units</b>					
<i>- Fuels Used for Heat Production</i>					
- - Refinery Gas				0103	308A
- - LPG				0102	303A
- - Gas-/Diesel Oil				0102	204A
- - Fuel Oil				0102	203A
- - Waste Oil				0102	203A
- - Petroleum Coke				0102	110A
- - Natural Gas				0102	301A
- - Electricity Plant Coal				0102	102A
- - Coal				0102	102A
- - Straw				0102	117A
- - Wood Chips				0102	111A
- - Wood Pellets				0102	111A
- - Wood Waste				0102	111A
- - Biogas, Landfill				0102	309A
- - Biogas, Sludge				0102	309A
- - Biogas, Other				0102	309A
- - Waste, Non-renewable				0102	114A
- - Wastes, Renewable				0102	114A
- - Fish Oil				0102	215A
<b>Autoproducers, Electricity Only</b>					
<i>- Fuels Used for Power Production</i>					
- - Natural Gas				0320	301A
- - Biogas, Landfill				0320	309A
- - Biogas, Sewage Sludge				0320	309A
- - Biogas, Other				0320	309A
<b>Autoproducers, CHP Units</b>					
<i>- Fuels Used for Power Production</i>					
- - Refinery Gas				0103	308A
- - Gas-/Diesel Oil				0320	204A
- - Fuel Oil				0320	203A
- - Waste Oil				0320	203A
- - Natural Gas				0320	301A
- - Coal				0320	102A
- - Straw				0320	117A
- - Wood Chips				0320	111A
- - Wood Pellets				0320	111A
- - Wood Waste				0320	111A
- - Biogas, Landfill				0320	309A
- - Biogas, Sludge				0320	309A
- - Biogas, Other				0320	309A
- - Fish Oil				0320	215A
- - Waste, Non-renewable				0320	114A




Unit: TJ	End-use			Transformation 1980-1993	
	SNAP	Fuel ( <i>in Danish</i> )	Fuel-code	SNAP	Fuel-code
- - Wastes, Renewable				0320	114A
- <i>Fuels Used for Heat Production</i>					
- - Refinery Gas				0103	308A
- - Gas-/Diesel Oil				0320	204A
- - Fuel Oil				0320	203A
- - Waste Oil				0320	203A
- - Natural Gas				0320	301A
- - Coal				0320	102A
- - Wood Chips				0320	111A
- - Wood Waste				0320	111A
- - Biogas, Landfill				0320	309A
- - Biogas, Sludge				0320	309A
- - Biogas, Other				0320	309A
- - Waste, Non-renewable				0320	114A
- - Wastes, Renewable				0320	114A
<b>Autoproducers, Heat Only</b>					
- <i>Fuels Used for Heat Production</i>					
- - Gas-/Diesel Oil				0320	204A
- - Fuel Oil				0320	203A
- - Waste Oil				0320	203A
- - Natural Gas				0320	301A
- - Straw				0320	117A
- - Wood Chips				0320	111A
- - Wood Chips				0320	111A
- - Wood Waste				0320	111A
- - Biogas, Landfill				0320	309A
- - Biogas, Sludge				0320	309A
- - Biogas, Other				0320	309A
- - Waste, Non-renewable				0102	114A
- - Wastes, Renewable				0102	114A
Town Gas Units	030106	Naturgas	301A		
- Fuels Used for Production of District Heating	030106	Kul (-83) / Gasolie (84-)	102A / 204A		
<b>Transport sector</b>					
<b>Military Transport</b>					
- Aviation Gasoline					
- Motor Gasoline					
- JP4					
- JP1					
- Gas-/Diesel Oil					
<b>Road</b>					
- LPG					
- Motor Gasoline					
- Other Kerosene	0202	Petroleum	206A		
- Gas-/Diesel Oil					
- Fuel Oil					
<b>Rail</b>					
- Motor Gasoline					
- Other Kerosene					
- Gas-/Diesel Oil					
- Electricity					
<b>Domestic Sea Transport</b>					
- LPG					
- Other Kerosene					
- Gas-/Diesel Oil					
- Fuel Oil					
<b>Air Transport, Domestic</b>					
- LPG					
- Aviation Gasoline					
- Motor Gasoline					
- Other Kerosene	0201	Petroleum	206A		
- JP1					
<b>Air Transport, International</b>					
- Aviation Gasoline					
- JP1					
<b>Agriculture and Forestry</b>					
- LPG					
- Motor Gasoline					
- Other Kerosene	0203	Petroleum	206A		
- Gas-/Diesel Oil					

Unit: TJ	End-use		Transformation 1980-1993		
	SNAP	Fuel ( <i>in Danish</i> )	Fuel-code	SNAP	Fuel-code
- Fuel Oil	0203	Fuelolie & Spildolie	203A		
- Petroleum Coke	0203	Petrokoks	110A		
- Natural Gas	0203	Naturgas	301A		
- Coal	0203	Kul	102A		
- Brown Coal Briquettes	0203	Brunkul	106A		
- Straw	0203	Halm	117A		
- Wood Chips	0203	Træ	111A		
- Wood Waste	0203	Træ	111A		
- Biogas, Other	0203	Biogas	309A		
<b>Horticulture</b>					
- LPG					
- Motor Gasoline					
- Gas-/Diesel Oil					
- Fuel Oil	0203	Fuelolie & Spildolie	203A		
- Petroleum Coke	0203	Petrokoks	110A		
- Natural Gas	0203	Naturgas	301A		
- Coal	0203	Kul	102A		
- Wood Waste	0203	Træ	111A		
<b>Fishing</b>					
- LPG					
- Motor Gasoline					
- Other Kerosene					
- Gas-/Diesel Oil					
- Fuel Oil					
<b>Manufacturing Industry</b>					
- Refinery Gas	0320	Raffinaderigas	308A		
- LPG					
- Naphtha (LVN)					
- Motor Gasoline					
- Other Kerosene	0320	Petroleum	206A		
- Gas-/Diesel Oil					
- Fuel Oil	0320	Fuelolie & Spildolie	203A		
- Waste Oil	0320	Fuelolie & Spildolie	203A		
- Petroleum Coke	0320	Petrokoks	110A		
- Natural Gas	0320	Naturgas	301A		
- Coal	0320	Kul	102A		
- Coke	0320	Koks	107A		
- Brown Coal Briquettes	0320	Brunkul	106A		
- Wood Pellets	0320	Træ	111A		
- Wood Waste	0320	Træ	111A		
- Biogas, Landfill	0320	Biogas	309A		
- Biogas, Other	0320	Biogas	309A		
- Wastes, Non-renewable	0320	Affald	114A		
- Wastes, Renewable	0320	Affald	114A		
- Town Gas	0320	Naturgas	301A		
<b>Construction</b>					
- LPG	0320	LPG	303A		
- Motor Gasoline					
- Other Kerosene	0320	Petroleum	206A		
- Gas-/Diesel Oil					
- Fuel Oil	0320	Fuelolie & Spildolie	203A		
- Natural Gas	0320	Naturgas	301A		
<b>Wholesale</b>					
- LPG	0201	LPG	303A		
- Motor Gasoline	0201	Petroleum	206A		
- Other Kerosene	0201	Gas & Dieselolie	204A		
- Gas-/Diesel Oil	0201	Fuelolie & Spildolie	203A		
- Petroleum Coke	0201	Petrokoks	110A		
- Natural Gas	0201	Naturgas	301A		
- Wood Waste	0201	Træ	111A		
<b>Retail Trade</b>					
- LPG	0201	LPG	303A		
- Other Kerosene	0201	Petroleum	206A		
- Gas-/Diesel Oil	0201	Gas & Dieselolie	204A		
- Fuel Oil	0201	Fuelolie & Spildolie	203A		
- Petroleum Coke	0201	Petrokoks	110A		
- Natural Gas	0201	Naturgas	301A		
<b>Private Service</b>					
- LPG	0201	LPG	303A		
- Other Kerosene	0201	Petroleum	206A		

Unit: TJ	End-use		Transformation 1980-1993	
	SNAP	Fuel ( <i>in Danish</i> )	Fuel-code	SNAP Fuel-code
- Gas-/Diesel Oil	0201	Gas & Dieselolie	204A	
- Fuel Oil	0201	Fuelolie & Spildolie	203A	
- Waste Oil	0201	Fuelolie & Spildolie	203A	
- Petroleum Coke	0201	Petrokoks	110A	
- Natural Gas	0201	Naturgas	301A	
- Wood Chips	0201	Træ	111A	
- Wood Waste	0201	Træ	111A	
- Biogas, Landfill	0201	Biogas	309A	
- Biogas, Sludge	0201	Biogas	309A	
- Biogas, Other	0201	Biogas	309A	
- Wastes, Non-renewable	0201	Affald	114A	
- Wastes, Renewable	0201	Affald	114A	
- Town Gas	0201	Naturgas	301A	
<b>Public Service</b>				
- LPG	0201	LPG	303A	
- Other Kerosene	0201	Petroleum	206A	
- Gas-/Diesel Oil	0201	Gas & Dieselolie	204A	
- Fuel Oil	0201	Fuelolie & Spildolie	203A	
- Petroleum Coke	0201	Petrokoks	110A	
- Natural Gas	0201	Naturgas	301A	
- Coal	0201	Kul	102A	
- Brown Coal Briquettes	0201	Brunkul	106A	
- Wood Chips	0201	Træ	111A	
- Wood Pellets	0201	Træ	111A	
- Town Gas	0201	Naturgas	301A	
<b>Single Family Houses</b>				
- LPG	0202	LPG	303A	
- Motor Gasoline				
- Other Kerosene	0202	Petroleum	206A	
- Gas-/Diesel Oil	0202	Gas & Dieselolie	204A	
- Fuel Oil	0202	Fuelolie & Spildolie	203A	
- Petroleum Coke	0202	Petrokoks	110A	
- Natural Gas	0202	Naturgas	301A	
- Coal	0202	Kul	102A	
- Coke	0202	koks	107A	
- Brown Coal Briquettes	0202	Brunkul	106A	
- Straw	0202	Halm	117A	
- Firewood	0202	Træ	111A	
- Wood Chips	0202	Træ	111A	
- Wood Pellets	0202	Træ	111A	
- Town Gas	0202	Naturgas	301A	
<b>Multi-family Houses</b>				
- LPG	0202	LPG	303A	
- Other Kerosene	0202	Petroleum	206A	
- Gas-/Diesel Oil	0202	Gas & Dieselolie	204A	
- Fuel Oil	0202	Fuelolie & Spildolie	203A	
- Petroleum Coke	0202	Petrokoks	110A	
- Natural Gas	0202	Naturgas	301A	
- Coal	0202	Kul	102A	
- Coke	0202	Koks	107A	
- Brown Coal Briquettes	0202	Brunkul	106A	
- Town Gas	0202	Naturgas	301A	



### Annex 3A-10 Time-series 1980/1985-2013



Continued




Figure 3A-10.1 Time-series for fuel consumption and emissions, 1980/1985 - 2013.

### **Annex 3A-11 QA/QC for stationary combustion**

An updated quality manual for the Danish emission inventories has been published in 2013 (Nielsen et al. 2013). The quality manual describes the concepts of quality work and definitions of sufficient quality, critical control points and a list of Point for Measuring (PM

Documentation concerning verification of the Danish emission inventories has been published by Fauser et al. (2013). The reference approach for the energy sector is shown in the annual National Inventory Report, Chapter 3.4.

The sector report for stationary combustion (Nielsen et al. 2014) has been reviewed by external experts in 2004, 2006, 2009 and 2014 (Nielsen et al. 2004; Nielsen et al. 2006; Nielsen et al. 2009; Nielsen et al., 2014). This forms a vital part of the QA activities for stationary combustion.

Source specific QA/QC and PM's are shown below.

### Data storage, level 1

Table 3A-11.1 lists the sectoral PM's for data storage level 1.

Table 3A-11.1 List of PM, data storage level 1.

Level	CCP	Id	Description	Sectoral/general	Stationary combustion
Data Storage level 1	1. Accuracy	DS.1.1.1	General level of uncertainty for every dataset including the reasoning for the specific values.	Sectoral	Uncertainties are estimated and references given in IIR chapter 3.2.
	2. Comparability	DS1.2.1	Comparability of the emission factors/calculation parameters with data from international guidelines, and evaluation of major discrepancies.	Sectoral	In general, if national referenced emission factors differ considerably from IPCC Guideline/EEA Guidebook values this is discussed in NIR chapter 3.2.4. This documentation is improved annually based on reviews. At CRF level, a project has been carried out comparing the Danish inventories with those of other countries (Fauser et al. 2013).
	3. Completeness	DS.1.3.1	Ensuring that the best possible national data for all sources are included, by setting down the reasoning behind the selection of datasets.	Sectoral	A list of external data are shown and discussed below.
	4. Consistency	DS.1.4.1	The original external data has to be archived with proper reference.	Sectoral	It is ensured that all external data are archived at DCE. Subsequent data processing takes place in other spreadsheets or databases. The datasets are archived annually in order to ensure that the basic data for a given report are always available in their original form.  In addition all references are archived.
	6. Robustness	DS.1.6.1	Explicit agreements between the external institution holding the data and DCE about the conditions of delivery	Sectoral	For stationary combustion, a data delivery agreement is made with the DEA. DCE and DEA have renewed the data delivery agreement in 2014. Most of the other external data sources are available due to legislative requirements. See Table 3.2.39.
	7. Transparency	DS.1.7.1	Listing of all archived datasets and external contacts.	Sectoral	A list of external datasets and external contacts is shown in Table 3A-11.2 below.

Table 3A-11.2 List of external data sources.

Dataset	Description	AD or Emf.	Reference	Contact(s)	Data agreement/ Comment
Energiproducenttællingen.xls	Data set for all electricity and heat producing plants.	Activity data	The Danish Energy Agency (DEA)	Kaj Stærkind	Data agreement.
Gas consumption for gas engines and gas turbines 1990-1994	Historical data set for gas engines and gas turbines.	Activity data	The Danish Energy Agency (DEA)	Jane Rusbjerg	No data agreement. Historical data
Basic data (Grunddata.xls)	The Danish energy statistics. Data set applied for both the reference approach and the national approach.	Activity data	The Danish Energy Agency (DEA)	Jane Rusbjerg	Data agreement. However, the data set is also published as part of national energy statistics
Energy statistics for industrial sub-sectors	Disaggregation of the industrial fuel consumption. The data set have been applied for the first time in the inventory reported in 2012.	Activity data	The Danish Energy Agency (DEA)	Jane Rusbjerg	Data agreement.
SO <sub>2</sub> & NO <sub>x</sub> data, plants>25 MW <sub>e</sub>	Annual emission data for all power plants > 25 MW <sub>e</sub> . Includes information on methodology: measurements or emission factor.	Emissions	Energinet.dk	Christian F.B. Nielsen	No data agreement.
Emission factors	Emission factors refer to a large number of sources.	Emission factors	See chapter regarding emission factors		Some of the annually updated CO <sub>2</sub> emission factors are based on EU ETS data, see below. For the other emission factors no formal data delivery agreement.
HM and PM from public power plants	Emissions from the two large power plant operators in DK DONG Energy and Vattenfall	Emissions	Dong Energy Vattenfall	Marina Snowman Møller, Heidi Demant	No formal data agreement.
Annual environmental reports / environmental data	Emissions from plants defined as large point sources	Emissions	Various plants		No data agreement necessary. Plants are obligated by law and data published on the Danish EPA homepage.
EU ETS data	Plant specific CO <sub>2</sub> emission factors	Emission factors and fuel consumption	The Danish Energy Agency (DEA)	Dorte Maimann Helen Falster	Plants are obligated by law. The availability of detailed information is part of the data agreement with DEA.

### **Energiproducenttaellingen - statistic on fuel consumption from district heating and power plants (DEA)**

The data set includes all plants producing power or district heating. The spreadsheet from DEA is listing fuel consumption of all plants included as large point sources in the emission inventory. The statistic on fuel consumption from district heating and power plants is regarded as complete and with no significant uncertainty since the plants are bound by law to report their fuel consumption and other information.

### **Gas consumption for gas engines and gas turbines 1990-1994 (DEA)**

For the years 1990-1994, DEA has estimated consumption of natural gas and biogas in gas engines and gas turbines. DCE assesses that the estimation by the DEA are the best available data.

### **Basic data (DEA)**

The Danish energy statistics. The spreadsheet from DEA is used for the CO<sub>2</sub> emission calculation in accordance with the IPCC reference approach and is also the first data set applied in the national approach. The data set is included in the data delivery agreement with DEA, but it is also published annually on DEA's homepage.

### **Energy statistics for industrial subsectors (DEA)**

The data includes disaggregation of the fuel consumption for industrial plants. The data set is estimated for the reporting to Eurostat. The dataset is included in the data agreement with DEA.

### **SO<sub>2</sub> and NO<sub>x</sub> emission data from electricity producing plants > 25MW<sub>e</sub> (Energinet.dk)**

Plants larger than 25 MW<sub>e</sub> are obligated to report emission data for SO<sub>2</sub> and NO<sub>x</sub> to the DEA annually. Data are on production unit level and classified. The data on plant level are part of the plants annually environmental reports. DCE's QC of the data consists of a comparison with data from previous years and with data from the plants' annual environmental reports.

### **Emission factors**

For specific references, see the chapter regarding emission factors. Some of the annually updated CO<sub>2</sub> emission factors are based on EU ETS data, see below.

### **Data for emission of heavy metals and particles from central power plants, DONG Energy and Vattenfall**

The two major Danish power plant operators assess heavy metal emissions from their plants using model calculations based on fuel data and type of flue gas cleaning. DCE's QC of the data consists of a comparison with data from previous years and with data from the plants' annual environmental reports.

### **Annual environmental reports (DEPA)**

A large number of plants are obligated by law to report annual environmental data including emission data. DCE compares the data with those from previous years and large discrepancies are checked.

### **EU ETS data (DEA)**

EU ETS data are information on fuel consumption, heating values, carbon content of fuel, oxidation factor and CO<sub>2</sub> emissions. DCE receives the veri-

fied reports for all plants which utilises a detailed estimation methodology. DCE's QC of the received data consists of comparing to calculation using standard emission factors as well as comparing reported values with those for previous years.

### Data processing, level 1

Table 3A-11.3 lists the sectoral PM's for data processing level 1.

Table 3A-11.3 List of PM, data processing level 1.

Level	CCP	Id	Description	Sectoral / general	Stationary combustion	
Data Processing level 1	1. Accuracy	DP.1.1.1	Uncertainty assessment for every data source not part of DS.1.1.1 as input to Data Storage level 2 in relation to type and scale of variability.	Sectoral	Uncertainties are estimated and references given in NIR chapter 3.2.5.	
	2.Comparability	DP.1.2.1	The methodologies have to follow the international guidelines suggested by UNFCCC and IPCC.	Sectoral	The methodological approach is consistent with international guidelines. An overview of tiers is given in NIR Chapter 3.2.5	
	3.Completeness	DP.1.3.1	Identification of data gaps with regard to data sources that could improve quantitative knowledge.	Sectoral	The energy statistics is considered complete.	
	4.Consistency	DP.1.4.1	Documentation and reasoning of methodological changes during the time series and the qualitative assessment of the impact on time series consistency.	Sectoral	The two main methodological changes in the time series; implementation of Energiproducentaellingen (plant specific fuel consumption data) from 1994 onwards and implementation of EU ETS data from 2006 onwards is discussed in NIR chapter 3.2.	
	5.Correctness		DP.1.5.2	Verification of calculation results using time series	Sectoral	Time series for activity data on SNAP and CRF source category level are used to identify possible errors. Time series for emission factors and the emission from CRF subcategories are also examined.
			DP.1.5.3	Verification of calculation results using other measures	Sectoral	The IPCC reference approach validates the fuel consumption rates and CO <sub>2</sub> emission. Both differ less than 2.0 % (1990-2013). The reference approach is further discussed in NIR Chapter 3.4.
	7.Transparency		DP.1.7.1	The calculation principle, the equations used and the assumptions made must be described.	Sectoral	This is included in NIR chapter 3.2.5.
			DP.1.7.2	Clear reference to dataset at Data Storage level 1	Sectoral	This is included in NIR chapter 3.2.5.
			DP.1.7.3	A manual log to collect information about recalculations.	Sectoral	-

## Data storage, level 2

Table 3A-11.4 lists the sectoral PM's for data storage level 2.

Table 3A-11.4 List of PM, data storage level 2.

Level	CCP	Id	Description	Sectoral / general	Stationary combustion
Data Storage level 2	5. Correctness	DS.2.5.1	Check if a correct data import to level 2 has been made	Sectoral	To ensure a correct connection between data on level 2 and level 1 different controls are in place, e.g. control of sums and random tests.

## Data storage level 4

Table 3A-11.5 lists the sectoral PM's for data storage level 4.

Table 3A-11.5 List of PM, data storage level 4.

Level	CCP	Id	Description	Sectoral / general	Stationary combustion
Data Storage level 4	4. Consistency	DS.4.4.3	The IEFs from the CRF are checked both regarding level and trend. The level is compared to relevant emission factors to ensure correctness. Large dips/jumps in the time series are explained.	Sectoral	Large dips/jumps in time series are discussed and explained in NIR chapter 3.2.

## Other QC procedures

The emission from each large point source is compared with the emission reported the previous year.

Some automated checks have been prepared for the emission databases:

- Check of units for fuel rate, emission factors and plant-specific emissions.
- Check of emission factors for large point sources. Emission factors for pollutants that are not plant-specific should be the same as those defined for area sources.
- Additional checks on database consistency.
- Emission factor references are included in this report.
- Annual environmental reports are kept for subsequent control of plant-specific emission data.
- QC checks of the country-specific emission factors have not been performed, but most factors are based on input from companies that have implemented some QA/QC work. The major power plant owner/operators in Denmark, DONG Energy and Vattenfall have obtained the ISO 14001 certification for an environmental management system. The Danish Gas Technology Centre and Force Technology both run accredited laboratories for emission measurements.

## National external review

The sector report for stationary combustion has been reviewed by external experts in 2004, 2006 and 2009 (Nielsen et al., 2004; Nielsen et al., 2006; Nielsen et al., 2009; Nielsen et al., 2014). This forms a vital part of the QA activities for stationary combustion.



## **Annex 3B - Transport and other mobile sources**

### **List of content**

Annex 3B-1: Fleet data 1985-2013 for road transport (No. vehicles)

Annex 3B-2: Mileage data 1985-2013 for road transport (km)

Annex 3B-3: EU directive emission limits for road transportation vehicles

Annex 3B-4: Basis emission factors for road transportation vehicles (g/km)

Annex 3B-5: Reduction factors for road transport emission factors

Annex 3B-6: Deterioration factors for road transport emission factors

Annex 3B-7: Final fuel consumption factors (MJ/km) and emission factors (g/km) in 2013

Annex 3B-8: Fuel consumption (GJ) and emissions (tons) per vehicle category and as totals

Annex 3B-9: COPERT IV:DEA statistics fuel use ratios and mileage adjustment factors

Annex 3B-10-1: Correspondence table between actual aircraft type codes and representative aircraft types

Annex 3B-10-2: LTO no. per representative aircraft type for domestic and int. flights (Copenhagen and other airports)

Annex 3B-10-3: No. of flights between Danish airports and airports in Greenland and Faroe Islands

Annex 3B-10-4: LTO fuel consumption and emission factors per representative aircraft type for Copenhagen Airport and other airports

Annex 3B-11-1: Stock data for diesel tractors 1985-2013

Annex 3B-11-2: Stock data for gasoline tractors 1985-2013

Annex 3B-11-3: Stock data for harvesters 1985-2013

Annex 3B-11-4: Stock data for fork lifts 1985-2013

Annex 3B-11-5: Stock data for construction machinery 1985-2013

Annex 3B-11-6: Stock data for machine pools 1985-2013

Annex 3B-11-7: Stock data for household and gardening machinery 1985-2013

Annex 3B-11-8: Stock data and engine size data for recreational craft 1985-2013

Annex 3B-11-9: Basis fuel consumption and emission factors, deterioration factors and transient factors for non road working machinery and equipment and recreational craft. Stock and activity data for certain specific types of machinery

Annex 3B-12-1: Annual traffic data (no. of round trips) for Danish domestic ferries 1990-2013

Annex 3B-12-2: Ferry service, ferry name, engine year, main engine MCR (kW), engine type, specific fuel consumption (sfc), aux. engine (kW)

Annex 3B-12-3: Sailing time (single trip) for Danish domestic ferries

Annex 3B-12-4: Engine load factor (% MCR) for Danish domestic ferries

Annex 3B-12-5: Round trip shares for Danish domestic ferries

Annex 3B-13-1: Specific fuel consumption, NO<sub>x</sub>, CO, VOC, NMVOC and CH<sub>4</sub> emission factors (g pr kWh) per engine year for diesel ship engines

Annex 3B-13-2: S-%, SO<sub>2</sub>, PM and BC emission factors (g/kg fuel and g/GJ) per fuel type for diesel ship engines

Annex 3B-14: Fuel sales figures from DEA, and further processed fuel consumption data suited for the Danish inventory

Annex 3B-15-1: Emission factors for 1990 in CollectER format

Annex 3B-15-2: Emission factors for 2013 in CollectER format

Annex 3B-15-3: Emissions for 1990 in CollectER format

Annex 3B-15-4: Emissions for 2013 in CollectER format

Annex 3B-15-5: Non-exhaust emission factors, activity data and total non-exhaust emissions of TSP, PM<sub>1</sub>, PM<sub>2.5</sub>, BC and heavy metals in 2013

Annex 3B-16-1: Fuel consumption 1985-2013 in NFR format

Annex 3B-16-2: Emissions 1985-2013 in NFR format

Annex 3B-17: Uncertainty estimates

### Annex 3B-1: Fleet data 1985-2013 for road transport (No. vehicles)

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-2: Mileage data 1985-2013 for road transport (km)

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-3: EU directive emission limits for road transportation vehicles

Private cars and light duty vehicles I (<1305 kg).

G prkm		EURO 1	EURO 2	EURO 3 <sup>1)</sup>	EURO 4	EURO 5	EURO 6
<u>Normal temp.</u>							
CO	Gasoline	2.72	2.2	2.3	1.0	1.0	1.0
	Diesel	2.72	1.0	0.64	0.5	0.5	0.5
HC	Gasoline	-	-	0.20	0.10	0.1	0.1
NMHC	Gasoline	-	-	-	-	0.068	0.068
NO <sub>x</sub>	Gasoline	-	-	0.15	0.08	0.06	0.06
	Diesel	-	-	0.5	0.25	0.18	0.08
HC+NO <sub>x</sub>	Gasoline	0.97	0.5	-	-	-	-
	Diesel	0.97	0.7/0.9 <sup>2)</sup>	0.56	0.30	0.23	0.17
Particulates	Diesel	0.14	0.08/0.10 <sup>2)</sup>	0.05	0.025	0.005	0.005
Particulates (#)		-	-	-	-	-	6x10 <sup>11 4)</sup>
<u>Low temp.</u>							
CO	Gasoline	-	-	-	15	15	15
HC	Gasoline	-	-	-	1.8	1.8	1.8
<u>Evaporation</u>							
HC <sup>3)</sup>	Gasoline	2.0	2.0	2.0	2.0	2.0	2.0

<sup>1)</sup> Changed test procedure at normal temperatures (40 s warm-up phase omitted) and for evaporation measurements. <sup>2)</sup> Less stringent emission limits for direct injection diesel engines. <sup>3)</sup> Unit: g/test. <sup>4)</sup> Applicable for diesel and gasoline direct injection (GDI). 6x10<sup>12</sup> within first three years of Euro 6 effective dates

Light duty vehicles II (1305-1760 kg)

G pr km		EURO 1	EURO 2	EURO 3 <sup>1)</sup>	EURO 4	EURO 5	EURO 6
<u>Normal temp.</u>							
CO	Gasoline	5.17	4.0	4.17	1.81	1.81	1.81
	Diesel	5.17	1.25	0.80	0.63	0.63	0.63
HC	Gasoline	-	-	0.25	0.13	0.13	0.13
NMHC	Gasoline	-	-	-	-	0.9	0.9
NO <sub>x</sub>	Gasoline	-	-	0.18	0.10	0.75	0.75
	Diesel	-	-	0.65	0.33	0.235	0.105
HC+NO <sub>x</sub>	Gasoline	1.4	0.6	-	-	-	-
	Diesel	1.4	1.0/1.3 <sup>2)</sup>	0.72	0.39	0.295	0.195
Particulates	Gasoline					0.005	0.005
	Diesel	0.19	0.12/0.14 <sup>2)</sup>	0.07	0.04	0.005	0.005
Particulates (#)		-	-	-	-	-	6x10 <sup>11 4)</sup>
<u>Low temp.</u>							
CO	Gasoline	-	-	-	24	24	24
HC	Gasoline	-	-	-	2.7	2.7	2.7
<u>Evaporation</u>							
HC <sup>3)</sup>	Gasoline	2.0	2.0	2.0	2.0	2.0	2.0

<sup>1)</sup> Changed test procedure at normal temperatures (40 s warm-up phase omitted) and for evaporation measurements. <sup>2)</sup> Less stringent emission limits for direct injection diesel engines. <sup>3)</sup> Unit: g/test. <sup>4)</sup> Applicable for diesel and gasoline direct injection (GDI). 6x10<sup>12</sup> within first three years of Euro 6 effective dates

Light duty vehicles III (>1760 kg).

G pr km		EURO 1	EURO 2	EURO 3 <sup>1)</sup>	EURO 4	EURO 5	EURO 6
<u>Normal temp.</u>							
CO	Gasoline	6.9	5.0	5.22	2.27	2.27	2.27
	Diesel	6.9	1.5	0.95	0.74	0.74	0.74
HC	Gasoline	-	-	0.29	0.16	0.16	0.16
NMHC	Gasoline					0.108	0.108
NO <sub>x</sub>	Gasoline	-	-	0.21	0.11	0.082	0.082
	Diesel	-	-	0.78	0.39	0.28	0.125
HC+NO <sub>x</sub>	Gasoline	1.7	0.7	-	-	-	-
	Diesel	1.7	1.2/1.6 <sup>2)</sup>	0.86	0.46	0.35	0.215
Particulates	Gasoline					0.005	0.005
	Diesel	0.25	0.17/0.20 <sup>2)</sup>	0.10	0.06	0.005	0.005
Particulates (#)		-	-	-	-	-	6x10 <sup>11 4)</sup>
<u>Low temp.</u>							
CO	Gasoline	-	-	-	30	30	30
HC	Gasoline	-	-	-	3.2	3.2	3.2
<u>Evaporation</u>							
HC <sup>3)</sup>	Gasoline	2.0	2.0	2.0	2.0	2.0	2.0

<sup>1)</sup> Changed test procedure at normal temperatures (40 s warm-up phase omitted) and for evaporation measurements. <sup>2)</sup> Less stringent emission limits for direct injection diesel engines. <sup>3)</sup> Unit: g/test. <sup>4)</sup> Applicable for diesel and gasoline direct injection (GDI). 6x10<sup>12</sup> within first three years of Euro 6 effective dates.

Heavy duty diesel vehicles.

(g pr kWh)	Test <sup>1)</sup>	EURO I	EURO II	EURO III	EURO			EEV <sup>2)</sup>
		1993	1996	2001	IV 2006	EURO V 2009	EURO VI 2014	
CO	ECE/ESC	4.5	4.0	2.1	1.5	1.5	1.5	1.5
	ETC	-	-	(5.45)	4.0	4.0	4.0	3.0
HC	ECE/ESC	1.1	1.1	0.66	0.46	0.46	0.13	0.25
	ETC	-	-	(0.78)	0.55	0.55	0.16	0.40
NO <sub>x</sub>	ECE/ESC	8.0	7.0	5.0	3.5	2.0	0.4	2.0
	ETC	-	-	(5.0)	3.5	2.0	0.4	2.0
Particulates <sup>3)</sup>	ECE/ESC	0.36/0.61	0.15/0.25	0.10/0.13	0.02	0.02	0.01	0.02
	ETC	-	-	(0.16/0.21)	0.03	0.03	0.01	0.02
	ELR	-	-	0.8	0.5	0.5		0.15
NH <sub>3</sub>	ECE/ESC						10 (ppm)	
	ETC						10 (ppm)	

<sup>1)</sup> Test procedure: Euro 1 og Euro 2: ECE (stationary)

Euro 3: ESC (stationary) + ELR (load response)

Euro 4, Euro 5 og EEV: ESC (stationary) + ETC (transient) + ELR (load response)

<sup>2)</sup> EEV: Emission limits for extra environmental friendly vehicles, used as a basis for economical incitements (gas fueled vehicles).

<sup>3)</sup> For Euro 1, Euro 2 og Euro 3 less stringent emission limits apply for small engines:

Euro 1: <85 kW

Euro 2: <0,7 l

Euro 3: <0,75 l

### Annex 3B-4: Basis emission factors for road transportation vehicles (g/km)

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-5: Reduction factors for road transport emission factors

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-6: Deterioration factors for road transport emission factors

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-7: Final fuel consumption factors (MJ/km) and emission factors (g/km) in 2013

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-8: Fuel consumption (GJ) and emissions (tonnes) per vehicle category and as totals

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

### Annex 3B-9: COPERT IV:DEA statistics fuel use ratios and mileage adjustment factors

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-10-1: Correspondence table between actual aircraft type codes and representative aircraft types**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-10-2: LTO no. per representative aircraft type for domestic and int. flights (Copenhagen and other airports)**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-10-3: No. of flights between Danish airports and airports in Greenland and Faroe Islands**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-10-4: LTO fuel consumption and emission factors per representative aircraft type for Copenhagen Airport and other airports**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-1: Stock data for diesel tractors 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-2: Stock data for gasoline tractors 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-3: Stock data for harvesters 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-4: Stock data for fork lifts 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-5: Stock data for construction machinery 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-6: Stock data for machine pools 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-7: Stock data for household and gardening machinery 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-8: Stock data and engine size data for recreational craft 1985-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-11-9: Basis fuel consumption and emission factors, deterioration factors and transient factors for non road working machinery and equipment and recreational craft. Stock and activity data for certain specific types of machinery**

Basis factors for diesel fuelled non road machinery.

Engine size [P=kW]	Emission Level	NO <sub>x</sub>	VOC	CO	N <sub>2</sub> O [g pr kWh]	NH <sub>3</sub>	TSP	Fuel
P<19	<1981	12,00	5,00	7,00	0,035	0,002	2,80	300
P<19	1981-1990	11,50	3,80	6,00	0,035	0,002	2,30	285
P<19	1991-Stage I	11,20	2,50	5,00	0,035	0,002	1,60	270
P<19	Stage I	11,20	2,50	5,00	0,035	0,002	1,60	270
P<19	Stage II	11,20	2,50	5,00	0,035	0,002	1,60	270
P<19	Stage IIIA	11,20	2,50	5,00	0,035	0,002	1,60	270
P<19	Stage IIIB	11,20	2,50	5,00	0,035	0,002	1,60	270
P<19	Stage IV	11,20	2,50	5,00	0,035	0,002	1,60	270
19<=P<37	<1981	18,00	2,50	6,50	0,035	0,002	2,00	300
19<=P<37	1981-1990	18,00	2,20	5,50	0,035	0,002	1,40	281
19<=P<37	1991-Stage I	9,80	1,80	4,50	0,035	0,002	1,40	262
19<=P<37	Stage I	9,80	1,80	4,50	0,035	0,002	1,40	262
19<=P<37	Stage II	6,50	0,60	2,20	0,035	0,002	0,40	262
19<=P<37	Stage IIIA	6,08	0,60	2,20	0,035	0,002	0,40	262
19<=P<37	Stage IIIB	6,08	0,60	2,20	0,035	0,002	0,40	262
19<=P<37	Stage IV	6,08	0,60	2,20	0,035	0,002	0,40	262
37<=P<56	<1981	7,70	2,40	6,00	0,035	0,002	1,80	290
37<=P<56	1981-1990	8,60	2,00	5,30	0,035	0,002	1,20	275
37<=P<56	1991-Stage I	11,50	1,50	4,50	0,035	0,002	0,80	260
37<=P<56	Stage I	7,70	0,60	2,20	0,035	0,002	0,40	260
37<=P<56	Stage II	5,50	0,40	2,20	0,035	0,002	0,20	260
37<=P<56	Stage IIIA	3,81	0,40	2,20	0,035	0,002	0,20	260
37<=P<56	Stage IIIB	3,81	0,28	2,20	0,035	0,002	0,03	260
37<=P<56	Stage IV	3,81	0,28	2,20	0,035	0,002	0,03	260
56<=P<75	<1981	7,70	2,40	6,00	0,035	0,002	1,80	290
56<=P<75	1981-1990	8,60	2,00	5,30	0,035	0,002	1,20	275
56<=P<75	1991-Stage I	11,50	1,50	4,50	0,035	0,002	0,80	260
56<=P<75	Stage I	7,70	0,60	2,20	0,035	0,002	0,40	260
56<=P<75	Stage II	5,50	0,40	2,20	0,035	0,002	0,20	260
56<=P<75	Stage IIIA	3,81	0,40	2,20	0,035	0,002	0,20	260
56<=P<75	Stage IIIB	2,97	0,28	2,20	0,035	0,002	0,03	260
56<=P<75	Stage IV	0,40	0,28	2,20	0,035	0,002	0,03	260
75<=P<130	<1981	10,50	2,00	5,00	0,035	0,002	1,40	280
75<=P<130	1981-1990	11,80	1,60	4,30	0,035	0,002	1,00	268
75<=P<130	1991-Stage I	13,30	1,20	3,50	0,035	0,002	0,40	255
75<=P<130	Stage I	8,10	0,40	1,50	0,035	0,002	0,20	255
75<=P<130	Stage II	5,20	0,30	1,50	0,035	0,002	0,20	255
75<=P<130	Stage IIIA	3,24	0,30	1,50	0,035	0,002	0,20	255
75<=P<130	Stage IIIB	2,97	0,13	1,50	0,035	0,002	0,03	255
75<=P<130	Stage IV	0,40	0,13	1,50	0,035	0,002	0,03	255
130<=P<560	<1981	17,80	1,50	2,50	0,035	0,002	0,90	270
130<=P<560	1981-1990	12,40	1,00	2,50	0,035	0,002	0,80	260
130<=P<560	1991-Stage I	11,20	0,50	2,50	0,035	0,002	0,40	250
130<=P<560	Stage I	7,60	0,30	1,50	0,035	0,002	0,20	250
130<=P<560	Stage II	5,20	0,30	1,50	0,035	0,002	0,10	250
130<=P<560	Stage IIIA	3,24	0,30	1,50	0,035	0,002	0,10	250
130<=P<560	Stage IIIB	1,80	0,13	1,50	0,035	0,002	0,03	250
130<=P<560	Stage IV	0,40	0,13	1,50	0,035	0,002	0,03	250



## Basis factors for 4-stroke gasoline non road machinery.

Engine	Size code	Size classe [S=ccm]	Emission Level	NO <sub>x</sub>	VOC	CO	N <sub>2</sub> O [g pr kWh]	NH <sub>3</sub>	TSP	Fuel
4-stroke	SH2	20<=S<50	<1981	2.4	33	198	0.002	0.03	0.09	496
4-stroke	SH2	20<=S<50	1981-1990	3.5	27.5	165	0.002	0.03	0.08	474
4-stroke	SH2	20<=S<50	1991-Stage I	4.7	22	132	0.002	0.03	0.06	451
4-stroke	SH2	20<=S<50	Stage I	4.7	22	132	0.002	0.03	0.06	406
4-stroke	SH2	20<=S<50	Stage II	4.7	22	132	0.002	0.03	0.06	406
4-stroke	SH3	S>=50	<1981	2.4	33	198	0.002	0.03	0.09	496
4-stroke	SH3	S>=50	1981-1990	3.5	27.5	165	0.002	0.03	0.08	474
4-stroke	SH3	S>=50	1991-Stage I	4.7	22	132	0.002	0.03	0.06	451
4-stroke	SH3	S>=50	Stage I	4.7	22	132	0.002	0.03	0.06	406
4-stroke	SH3	S>=50	Stage II	4.7	22	132	0.002	0.03	0.06	406
4-stroke	SN1	S<66	<1981	1.2	26.9	822	0.002	0.03	0.09	603
4-stroke	SN1	S<66	1981-1990	1.8	22.5	685	0.002	0.03	0.08	603
4-stroke	SN1	S<66	1991-Stage I	2.4	18	548	0.002	0.03	0.06	603
4-stroke	SN1	S<66	Stage I	4.3	16.1	411	0.002	0.03	0.06	475
4-stroke	SN1	S<66	Stage II	4.3	16.1	411	0.002	0.03	0.06	475
4-stroke	SN2	66<=S<100	<1981	2.3	10.5	822	0.002	0.03	0.09	627
4-stroke	SN2	66<=S<100	1981-1990	3.5	8.7	685	0.002	0.03	0.08	599
4-stroke	SN2	66<=S<100	1991-Stage I	4.7	7	548	0.002	0.03	0.06	570
4-stroke	SN2	66<=S<100	Stage I	4.7	7	467	0.002	0.03	0.06	450
4-stroke	SN2	66<=S<100	Stage II	4.7	7	467	0.002	0.03	0.06	450
4-stroke	SN3	100<=S<225	<1981	2.6	19.1	525	0.002	0.03	0.09	601
4-stroke	SN3	100<=S<225	1981-1990	3.8	15.9	438	0.002	0.03	0.08	573
4-stroke	SN3	100<=S<225	1991-Stage I	5.1	12.7	350	0.002	0.03	0.06	546
4-stroke	SN3	100<=S<225	Stage I	5.1	11.6	350	0.002	0.03	0.06	546
4-stroke	SN3	100<=S<225	Stage II	5.1	9.4	350	0.002	0.03	0.06	546
4-stroke	SN4	S>=225	<1981	1.3	11.1	657	0.002	0.03	0.09	539
4-stroke	SN4	S>=225	1981-1990	2	9.3	548	0.002	0.03	0.08	514
4-stroke	SN4	S>=225	1991-Stage I	2.6	7.4	438	0.002	0.03	0.06	490
4-stroke	SN4	S>=225	Stage I	2.6	7.4	438	0.002	0.03	0.06	490
4-stroke	SN4	S>=225	Stage II	2.6	7.4	438	0.002	0.03	0.06	490

Basis factors for 2-stroke gasoline non road machinery.

Engine	Size code	Size classe [ccm]	Emission Level	NO <sub>x</sub>	VOC	CO	N <sub>2</sub> O [g pr kWh]	NH <sub>3</sub>	TSP	Fuel
2-stroke	SH2	20<=S<50	<1981	1	305	695	0.002	0.01	7	882
2-stroke	SH2	20<=S<50	1981-1990	1	300	579	0.002	0.01	5.3	809
2-stroke	SH2	20<=S<50	1991-Stage I	1.1	203	463	0.002	0.01	3.5	735
2-stroke	SH2	20<=S<50	Stage I	1.5	188	379	0.002	0.01	3.5	720
2-stroke	SH2	20<=S<50	Stage II	1.5	44	379	0.002	0.01	3.5	500
2-stroke	SH3	S>=50	<1981	1.1	189	510	0.002	0.01	3.6	665
2-stroke	SH3	S>=50	1981-1990	1.1	158	425	0.002	0.01	2.7	609
2-stroke	SH3	S>=50	1991-Stage I	1.2	126	340	0.002	0.01	1.8	554
2-stroke	SH3	S>=50	Stage I	2	126	340	0.002	0.01	1.8	529
2-stroke	SH3	S>=50	Stage II	1.2	64	340	0.002	0.01	1.8	500
2-stroke	SN1	S<66	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN1	S<66	Stage II	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN2	66<=S<100	Stage II	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN3	100<=S<225	Stage II	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	<1981	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	1981-1990	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	1991-Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	Stage I	0.5	155	418	0.002	0.01	2.6	652
2-stroke	SN4	S>=225	Stage II	0.5	155	418	0.002	0.01	2.6	652

Fuel consumption and emission factors for LPG fork lifts.

NO <sub>x</sub>	VOC	CO	NH <sub>3</sub>	N <sub>2</sub> O	TSP	FC
[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]	[g pr kWh]
19	2.2	1.5	0.003	0.05	0.07	311

Fuel consumption and emission factors for All Terrain Vehicles (ATV's).

ATV type	NO <sub>x</sub>	VOC	CO	NH <sub>3</sub>	N <sub>2</sub> O	TSP	Fuel
	[g pr GJ]	[g pr GJ]	[g pr GJ]	[g pr GJ]	[g pr GJ]	[g pr GJ]	[kg pr hour]
Professional	108	1077	16306	2	2	32	1.125
Private	128	1527	22043	2	2	39	0.75

Fuel consumption and emission factors for recreational craft.

Fuel type	Vessel type	Engine	Engine type	Direktiv	Engine size	CO	VOC	N <sub>2</sub> O	NH <sub>3</sub>	NO <sub>x</sub>	TSP	Fuel
					[kW]	[g pr kWh]						
Gasoline	Other boats (< 20 ft)	Out board	2-stroke	2003/44	8	202.5	45.9	0.01	0.002	2.0	10.00	791
Gasoline	Other boats (< 20 ft)	Out board	2-stroke	Konv.	8	427.0	257.0	0.01	0.002	2.0	10.00	791
Gasoline	Other boats (< 20 ft)	Out board	4-stroke	2003/44	8	202.5	24.0	0.03	0.002	7.0	0.08	426
Gasoline	Other boats (< 20 ft)	Out board	4-stroke	Konv.	8	520.0	24.0	0.03	0.002	7.0	0.08	426
Gasoline	Yawls and cabin boats	Out board	2-stroke	2003/44	20	162.0	36.5	0.01	0.002	3.0	10.00	791
Gasoline	Yawls and cabin boats	Out board	2-stroke	Konv.	20	374.0	172.0	0.01	0.002	3.0	10.00	791
Gasoline	Yawls and cabin boats	Out board	4-stroke	2003/44	20	162.0	14.0	0.03	0.002	10.0	0.08	426
Gasoline	Yawls and cabin boats	Out board	4-stroke	Konv.	20	390.0	14.0	0.03	0.002	10.0	0.08	426
Gasoline	Sailing boats (< 26 ft)	Out board	2-stroke	2003/44	10	189.0	43.0	0.01	0.002	2.0	10.00	791
Gasoline	Sailing boats (< 26 ft)	Out board	2-stroke	Konv.	10	427.0	257.0	0.01	0.002	2.0	10.00	791
Gasoline	Sailing boats (< 26 ft)	Out board	4-stroke	2003/44	10	189.0	24.0	0.03	0.002	7.0	0.08	426
Gasoline	Sailing boats (< 26 ft)	Out board	4-stroke	Konv.	10	520.0	24.0	0.03	0.002	7.0	0.08	426
Gasoline	Speed boats	In board	4-stroke	2003/44	90	141.0	10.0	0.03	0.002	12.0	0.08	426
Gasoline	Speed boats	In board	4-stroke	Konv.	90	346.0	10.0	0.03	0.002	12.0	0.08	426
Gasoline	Speed boats	Out board	2-stroke	2003/44	50	145.8	31.8	0.01	0.002	3.0	10.00	791
Gasoline	Speed boats	Out board	2-stroke	Konv.	50	374.0	172.0	0.01	0.002	3.0	10.00	791
Gasoline	Speed boats	Out board	4-stroke	2003/44	50	145.8	14.0	0.03	0.002	10.0	0.08	426
Gasoline	Speed boats	Out board	4-stroke	Konv.	50	390.0	14.0	0.03	0.002	10.0	0.08	426
Gasoline	Water scooters	Built in	2-stroke	2003/44	45	147.0	32.2	0.01	0.002	3.0	10.00	791
Gasoline	Water scooters	Built in	2-stroke	Konv.	45	374.0	172.0	0.01	0.002	3.0	10.00	791
Gasoline	Water scooters	Built in	4-stroke	2003/44	45	147.0	14.0	0.03	0.002	10.0	0.08	426
Gasoline	Water scooters	Built in	4-stroke	Konv.	45	390.0	14.0	0.03	0.002	10.0	0.08	426
Diesel	Motor boats (27-34 ft)	In board		2003/44	150	5.0	1.7	0.035	0.002	8.6	1.00	275
Diesel	Motor boats (27-34 ft)	In board		Konv.	150	5.3	2.0	0.035	0.002	8.6	1.20	275
Diesel	Motor boats (> 34 ft)	In board		2003/44	250	5.0	1.6	0.035	0.002	8.6	1.00	275
Diesel	Motor boats (> 34 ft)	In board		Konv.	250	5.3	2.0	0.035	0.002	8.6	1.20	275
Diesel	Motor boats (< 27 ft)	In board		2003/44	40	5.0	1.8	0.035	0.002	9.8	1.00	281
Diesel	Motor boats (< 27 ft)	In board		Konv.	40	5.5	2.2	0.035	0.002	18.0	1.40	281
Diesel	Motor sailers	In board		2003/44	30	5.0	1.9	0.035	0.002	9.8	1.00	281
Diesel	Motor sailers	In board		Konv.	30	5.5	2.2	0.035	0.002	18.0	1.40	281
Diesel	Sailing boats (> 26 ft)	In board		2003/44	30	5.0	1.9	0.035	0.002	9.8	1.00	281
Diesel	Sailing boats (> 26 ft)	In board		Konv.	30	5.5	2.2	0.035	0.002	18.0	1.40	281

CH<sub>4</sub> shares of VOC for diesel, gasoline and LPG.

Fuel type	CH <sub>4</sub> share of VOC
Diesel	0.024
Gasoline 4-stroke	0.034
Gasoline 2-stroke	0.07
LPG	0.05

Deterioration factors for diesel machinery.

Emission Level	NO <sub>x</sub>	VOC	CO	TSP
<1981	0.024	0.047	0.185	0.473
1981-1990	0.024	0.047	0.185	0.473
1991-Stage I	0.024	0.047	0.185	0.473
Stage I	0.024	0.036	0.101	0.473
Stage II	0.009	0.034	0.101	0.473
Stage IIIA	0.008	0.027	0.151	0.473
Stage IIIB	0.008	0.027	0.151	0.473
Stage IV	0.008	0.027	0.151	0.473

Deterioration factors for gasoline 2-stroke machinery.

Engine	Size code	Size classe	Emission Level	NO <sub>x</sub>	VOC	CO	TSP
2-stroke	SH2	20<=S<50	<1981	0	0.2	0.2	0
2-stroke	SH2	20<=S<50	1981-1990	0	0.2	0.2	0
2-stroke	SH2	20<=S<50	1991-Stage I	0	0.2	0.2	0
2-stroke	SH2	20<=S<50	Stage I	0	0.29	0.24	0
2-stroke	SH2	20<=S<50	Stage II	0	0.29	0.24	0
2-stroke	SH3	S>=50	<1981	-0.031	0.2	0.2	0
2-stroke	SH3	S>=50	1981-1990	-0.031	0.2	0.2	0
2-stroke	SH3	S>=50	1991-Stage I	-0.031	0.2	0.2	0
2-stroke	SH3	S>=50	Stage I	0	0.266	0.231	0
2-stroke	SH3	S>=50	Stage II	0	0.266	0.231	0
2-stroke	SN1	S<66	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN1	S<66	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN1	S<66	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN1	S<66	Stage I	-0.33	0.266	1.109	5.103
2-stroke	SN1	S<66	Stage II	-0.33	0	1.109	5.103
2-stroke	SN2	66<=S<100	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN2	66<=S<100	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN2	66<=S<100	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN2	66<=S<100	Stage I	-0.33	0.266	1.109	5.103
2-stroke	SN2	66<=S<100	Stage II	-0.33	0	1.109	5.103
2-stroke	SN3	100<=S<225	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN3	100<=S<225	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN3	100<=S<225	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN3	100<=S<225	Stage I	-0.33	0.266	1.109	5.103
2-stroke	SN3	100<=S<225	Stage II	-0.33	0	1.109	5.103
2-stroke	SN4	S>=225	<1981	-0.6	0.201	0.9	1.1
2-stroke	SN4	S>=225	1981-1990	-0.6	0.201	0.9	1.1
2-stroke	SN4	S>=225	1991-Stage I	-0.6	0.201	0.9	1.1
2-stroke	SN4	S>=225	Stage I	-0.274	0	0.887	1.935
2-stroke	SN4	S>=225	Stage II	-0.274	0	0.887	1.935

Deterioration factors for gasoline 4-stroke machinery.

Engine	Size code	Size classe	Emission Level	NO <sub>x</sub>	VOC	CO	TSP
4-stroke	SN1	S<66	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN1	S<66	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN1	S<66	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN1	S<66	Stage I	-0.3	1.753	1.051	1.753
4-stroke	SN1	S<66	Stage II	-0.3	1.753	1.051	1.753
4-stroke	SN2	66<=S<100	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN2	66<=S<100	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN2	66<=S<100	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN2	66<=S<100	Stage I	-0.3	1.753	1.051	1.753
4-stroke	SN2	66<=S<100	Stage II	-0.3	1.753	1.051	1.753
4-stroke	SN3	100<=S<225	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN3	100<=S<225	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN3	100<=S<225	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN3	100<=S<225	Stage I	-0.3	1.753	1.051	1.753
4-stroke	SN3	100<=S<225	Stage II	-0.3	1.753	1.051	1.753
4-stroke	SN4	S>=225	<1981	-0.6	1.1	0.9	1.1
4-stroke	SN4	S>=225	1981-1990	-0.6	1.1	0.9	1.1
4-stroke	SN4	S>=225	1991-Stage I	-0.6	1.1	0.9	1.1
4-stroke	SN4	S>=225	Stage I	-0.599	1.095	1.307	1.095
4-stroke	SN4	S>=225	Stage II	-0.599	1.095	1.307	1.095
4-stroke	SH2	20<=S<50	<1981	0	0	0	0
4-stroke	SH2	20<=S<50	1981-1990	0	0	0	0
4-stroke	SH2	20<=S<50	1991-Stage I	0	0	0	0
4-stroke	SH2	20<=S<50	Stage I	0	0	0	0
4-stroke	SH2	20<=S<50	Stage II	0	0	0	0
4-stroke	SH3	S>=50	<1981	0	0	0	0
4-stroke	SH3	S>=50	1981-1990	0	0	0	0
4-stroke	SH3	S>=50	1991-Stage I	0	0	0	0
4-stroke	SH3	S>=50	Stage I	0	0	0	0
4-stroke	SH3	S>=50	Stage II	0	0	0	0

Transient factors for diesel machinery.

Emission Level	Load	Load factor	NO <sub>x</sub>	VOC	CO	TSP	Fuel
<1981	High	> 0.45	0.95	1.05	1.53	1.23	1.01
1981-1990	High	> 0.45	0.95	1.05	1.53	1.23	1.01
1991-Stage I	High	> 0.45	0.95	1.05	1.53	1.23	1.01
Stage I	High	> 0.45	0.95	1.05	1.53	1.23	1.01
Stage II	High	> 0.45	0.95	1.05	1.53	1.23	1.01
Stage IIIA	High	> 0.45	1.04	1.05	1.53	1.47	1.01
Stage IIIB	High	> 0.45	1	1	1	1	1
Stage IV	High	> 0.45	1	1	1	1	1
<1981	Middle	0.25>= LF <= 0.45	1.025	1.67	2.05	1.6	1.095
1981-1990	Middle	0.25>= LF <= 0.45	1.025	1.67	2.05	1.6	1.095
1991-Stage I	Middle	0.25>= LF <= 0.45	1.025	1.67	2.05	1.6	1.095
Stage I	Middle	0.25>= LF <= 0.45	1.025	1.67	2.05	1.6	1.095
Stage II	Middle	0.25>= LF <= 0.45	1.025	1.67	2.05	1.6	1.095
Stage IIIA	Middle	0.25>= LF <= 0.45	1.125	1.67	2.05	1.92	1.095
Stage IIIB	Middle	0.25>= LF <= 0.45	1	1	1	1	1
Stage IV	Middle	0.25>= LF <= 0.45	1	1	1	1	1
<1981	Low	<0.25	1.1	2.29	2.57	1.97	1.18
1981-1990	Low	<0.25	1.1	2.29	2.57	1.97	1.18
1991-Stage I	Low	<0.25	1.1	2.29	2.57	1.97	1.18
Stage I	Low	<0.25	1.1	2.29	2.57	1.97	1.18
Stage II	Low	<0.25	1.1	2.29	2.57	1.97	1.18
Stage IIIA	Low	<0.25	1.21	2.29	2.57	2.37	1.18
Stage IIIB	Low	<0.25	1	1	1	1	1
Stage IV	Low	<0.25	1	1	1	1	1

Annual working hours, load factors and lifetimes for agricultural tractors.

Tractor type	Annual working hours	Load factor	Lifetime (yrs)
Diesel	500 (0-7 years)	0.5	30
	500-100 (7-16 years)		
	100 (>16 years)		
Gasoline (certified)	100	0.4	37
Gasoline (non certified)	50	0.4	37

Annual working hours, load factors and lifetimes for harvesters.

Annual working hours	Load factor	Lifetime (yrs)
250-100 (linear decrease 0-24 years)	0.8	25

Annual working hours, load factors and lifetime for machine pool machinery.

Tractor type	Hours pr yr	Load factor	Lifetime (yrs)
Tractors	750	0.5	7
Harvesters	100	0.8	11
Self-propelled vehicles	500	0.75	6

Operational data for other machinery types in agriculture.

Machinery type	Fuel type	Load factor	Lifetime (yrs)	Hours	Size (kW)
ATV private	Gasoline	-	6	250	-
ATV professional	Gasoline	-	8	400	-
Bedding machines	Gasoline	0.3	10	50	3
Fodder trucks	Gasoline	0.4	10	200	8
Other (gasoline)	Gasoline	0.4	10	50	5
Scrapers	Gasoline	0.3	10	50	3
Self-propelled vehicles	Diesel	0.75	15	150	60
Sweepers	Gasoline	0.3	10	50	3

Annual working hours, load factors and lifetimes for forestry machinery.

Machinery type	Hours	Load factors	Lifetime
Chippers	1200	0.5	6
Tractors (other)	100 (1990) 400 (2004)	0.5	15
Tractors (silvicultural)	800	0.5	6
Harvesters	1200	0.5	8
Forwarders	1200	0.5	8
Chain saws (forestry)	800	0.4	3

Annual working hours, load factors and lifetime for fork lifts.

Hours pr yr	Load factor	Lifetime (yrs)
1200 (>=50 kW and <=10 years old)	0.27	20
650 (>=50 kW and >10 years old)		
650 (<50 kW)		

Operational data for construction machinery.

Machinery type	Load factor	Lifetime	Hours	Size
Track type dozers	0.5	10	1100	140
Track type loaders	0.5	10	1100	100 (1990) 150 (2004)
Wheel loaders (0-5 tonnes)	0.5	10	1200	20
Wheel loaders (> 5,1 tonnes)	0.5	10	1200	120
Wheel type excavators	0.6	10	1200	100
Track type excavators (0-5 tonnes)	0.6	10	1100	20
Track type excavators (>5,1 tonnes)	0.6	10	1100	120
Excavators/Loaders	0.45	10	700	50
Dump trucks	0.4	10	900 (1990) 1200 (2004)	60 (1990) 180 (2004)
Mini loaders	0.5	14	700	30
Telescopic loaders	0.5	14	1000	35

Stock and operational data for other machinery types in industry.

Sector	Fuel type	Machinery type	Size (kW)	No Load Factor	Hours	
Construction machinery	Diesel	Tampers/Land rollers	30	2800	0.45	600
Construction machinery	Diesel	Generators (diesel)	45	5000	0.5	200
Construction machinery	Diesel	Kompressors (diesel)	45	5000	0.5	500
Construction machinery	Diesel	Pumps (diesel)	75	1000	0.5	5
Construction machinery	Diesel	Asphalt pavers	80	300	0.35	700
Construction machinery	Diesel	Motor graders	100	100	0.4	700
Construction machinery	Diesel	Refuse compressors	160	100	0.25	1300
Construction machinery	Gasoline	Generators (gasoline)	2.5	11000	0.4	80
Construction machinery	Gasoline	Pumps (gasoline)	4	10000	0.4	300
Construction machinery	Gasoline	Kompressors (gasoline)	4	500	0.35	15
Industry	Diesel	Refrigerating units (distribution)	8	3000	0.5	1250
Industry	Diesel	Refrigerating units (long distance)	15	3500	0.5	200
Industry	Diesel	Tractors (transport, industry)	50	3000	0.4	500
Airport GSE and other	Diesel	Airport GSE and other (light duty)	100	500	0.5	400
Airport GSE and other	Diesel	Airport GSE and other (medium duty)	125	350	0.5	300
Airport GSE and other	Diesel	Airport GSE and other (Heavy duty)	175	650	0.5	200
Building and construction	Diesel	Vibratory plates	6	3500	0.6	300
Building and construction	Diesel	Aereal lifts (diesel)	30	150	0.4	400
Building and construction	Diesel	Sweepers (diesel)	30	200	0.4	300
Building and construction	Diesel	High pressure cleaners (diesel)	30	50	0.8	500
Building and construction	Gasoline	Rammers	2.5	3000	0.4	80
Building and construction	Gasoline	Drills	3	100	0.4	10
Building and construction	Gasoline	Vibratory plates (gasoline)	4	2500	0.5	200
Building and construction	Gasoline	Cutters	4	800	0.5	50
Building and construction	Gasoline	Other (gasoline)	5	1000	0.5	40
Building and construction	Gasoline	High pressure cleaners (gasoline)	5	500	0.6	200
Building and construction	Gasoline	Sweepers (gasoline)	10	500	0.4	150
Building and construction	Gasoline	Slicers	10	100	0.7	150
Building and construction	Gasoline	Aereal lifts (gasoline)	20	50	0.4	400

Operational data for the most important types of household and gardening machinery.

Machinery type	Engine	Size (kW)	Hours	Load factor	Lifetime (yrs)
Chain saws (private)	2-stroke	2	5	0.3	10
Chain saws (professional)	2-stroke	3	270	0.4	3
Cultivators (private-large)	4-stroke	3.7	5	0.6	5
Cultivators (private-small)	4-stroke	1	5	0.6	15
Cultivators (professional)	4-stroke	7	360	0.6	8
Hedge cutters (private)	2-stroke	0.9	10	0.5	10
Hedge cutters (professional)	2-stroke	2	300	0.5	4
		2.5	25		
		(2000)			
		3.5			
Lawn movers (private)	4-stroke	(2004)		0.4	8
		2.5	250		
		(2000)			
		3.5			
Lawn movers (professional)	4-stroke	(2004)		0.4	4
Riders (private)	4-stroke	11	50	0.5	12
Riders (professional)	4-stroke	13	330	0.5	5
Shrub clearers (private)	2-stroke	1	15	0.6	10
Shrub clearers (professional)	2-stroke	2	300	0.6	4
Trimmers (private)	2-stroke	0.9	20	0.5	10
Trimmers (professional)	2-stroke	0.9	200	0.5	4



Stock and operational data for other machines in household and gardening.

Machinery type	Engine	No.	Size		Load factor	Lifetime (yrs)
			(kW)	Hours		
Chippers	2-stroke	200	10	100	0.7	10
Garden shredders	2-stroke	500	3	20	0.7	10
Other (gasoline)	2-stroke	200	2	20	0.5	10
Suction machines	2-stroke	300	4	80	0.5	10
Wood cutters	4-stroke	100	4	15	0.5	10

Operational data for recreational craft.

Fuel type	Vessel type	Engine type	Stroke	Hours	Lifetime	Load factor
Gasoline	Other boats (<20 ft)	Out board engine	2-stroke	30	10	0.5
Gasoline	Other boats (<20 ft)	Out board engine	4-stroke	30	10	0.5
Gasoline	Yawls and cabin boats	Out board engine	2-stroke	50	10	0.5
Gasoline	Yawls and cabin boats	Out board engine	4-stroke	50	10	0.5
Gasoline	Sailing boats (<26ft)	Out board engine	2-stroke	5	10	0.5
Gasoline	Sailing boats (<26ft)	Out board engine	4-stroke	5	10	0.5
Gasoline	Speed boats	In board engine	4-stroke	75	10	0.5
Gasoline	Speed boats	Out board engine	2-stroke	50	10	0.5
Gasoline	Speed boats	Out board engine	4-stroke	50	10	0.5
Gasoline	Water scooters	Built in	2-stroke	10	10	0.5
Gasoline	Water scooters	Built in	4-stroke	10	10	0.5
Diesel	Motor boats (27-34 ft)	In board engine		150	15	0.5
Diesel	Motor boats (>34 ft)	In board engine		100	15	0.5
Diesel	Motor boats (<27 ft)	In board engine		75	15	0.5
Diesel	Motor sailors	In board engine		75	15	0.5
Diesel	Sailing boats (<26ft)	In board engine		25	15	0.5

**Annex 3B-12-1: Annual traffic data (no. of round trips) for Danish domestic ferries 1990-2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-12-2: Ferry service, ferry name, engine year, main engine MCR (kW), engine type, specific fuel consumption (sfc), aux. engine (kW)**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-12-3: Sailing time (single trip) for Danish domestic ferries**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-12-4: Engine load factor (% MCR) for Danish domestic ferries**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-12-5: Round trip shares for Danish domestic ferries**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-13-1: Specific fuel consumption, NO<sub>x</sub>, CO, VOC, NMVOC and CH<sub>4</sub> emission factors (g pr kWh) per engine year for diesel ship engines**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-13-2: S-%, SO<sub>2</sub>, PM and BC emission factors (g/kg fuel and g/GJ) per fuel type for diesel ship engines**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-14: Fuel sales figures from DEA, and further processed fuel consumption data suited for the Danish inventory**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-15-1: Emission factors for 1990 in CollectER format**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-15-2: Emission factors for 2013 in CollectER format**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-15-3: Emissions for 1990 in CollectER format**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-15-4: Emissions for 2013 in CollectER format**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-15-5: Non-exhaust emission factors, activity data and total non-exhaust emissions of TSP, PM<sub>1</sub>, PM<sup>2.5</sup>, BC and heavy metals in 2013**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-16-1: Fuel consumption 1985-2013 in NFR format**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-16-2: Emissions 1985-2013 in NFR format**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Annex 3B-17: Uncertainty estimates**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

## Annex 3C - Industrial processes (NFR 2)

### Non-energy products from fuels and solvent use (NFR 2D)

Annex 3C-1:	Activity data Solvent use (Gg per year)
Annex 3C-2:	Emission factors Solvent use (Gg NMVOC per Gg)
Annex 3C-3:	Emissions Solvent use (Gg NMVOC per year)
Annex 3C-4:	Activity data for asphalt in road paving (Mg per year)
Annex 3C-5:	NMVOC and CO emissions from road paving with asphalt (Gg per year)
Annex 3C-6:	Activity data for asphalt roofing (Mg per year)
Annex 3C-7:	NMVOC and CO emissions from asphalt roofing (Gg per year)

### Other product use (NFR 2G)

Annex 3C-8:	Activity data Other product use (Paraffin wax, Fireworks, Charcoal, Tobacco) (Gg per year)
Annex 3C-9:	Emission factors Other product use (Paraffin wax)
Annex 3C-10:	Emission factors Other product use (Fireworks)
Annex 3C-11:	Emission factors Other product use (Charcoal for barbeques)
Annex 3C-12:	Emission factors Other product use (Tobacco)
Annex 3C-13:	Emissions Other product use (Paraffin wax)
Annex 3C-14:	Emissions Other product use (Fireworks)
Annex 3C-15:	Emissions Other product use (Charcoal for barbeques)
Annex 3C-16:	Emissions Other product use (Tobacco)

All annexes are available online at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

## Annex 3D - Agriculture

**Table 3D-1: Number of animals allocated on subcategories. See:**  
<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Table 3D-2a: Nitrogen excretion rates in average, kg N per head per year. See:**  
<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Table 3D-2b: Nitrogen excretion given as TAN (Total Ammonical Nitrogen), kg N per head per year. See:**  
<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Table 3D-3: Changes in housing type. See:**  
<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3D-4: Cover of slurry tanks 1985-2013, pct. with no or full cover

	1985-1999	2000-2001	2002	2003-2013
Cattle				
		Percent		
No cover	20	5	5	2
Full cover	80	95	95	98
Swine				
No cover	40	20	10	5%
Full cover	60	80	90	95
Fur animals				
No cover	20	5	5	2
Full cover	80	95	95	98

Ref: COWI (2000)

**Table 3D-5: PM emission from housings, Gg TSP, PM<sub>10</sub> and PM<sub>2.5</sub>. See:**  
<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

Table 3D-6: Assumptions for synthetic fertiliser

EMEP/EEA fertiliser types <sup>1</sup>	Danish fertiliser types
Ammonium nitrate (AN)	Ammonium nitrate
Anhydrous ammonia	Liquid ammonia
Ammonium phosphates (MAP, DAP)	Calcium and boron calcium nitrate Diammonphosphate Other NP fertiliser types Magnesium fertiliser
Ammonium sulphate (AS)	Ammonium sulphate
Calcium ammonium nitrate (CAN)	Calcium ammonium nitrate and other nitrate types
Calcium nitrate (CN)	-
Ammonia solutions (AN)	Other nitrogen fertiliser
Ammonia solutions (Urea AN)	-
Urea ammonium solutions (UAS)	-
Urea	Urea
Other NK and NPK	NPK-fertiliser NK fertiliser

<sup>1</sup> EMEP/EEA emission inventory guidebook 2013, Table 3-2 Emission factors for total NH<sub>3</sub> emissions from soils due to N fertiliser volatilization.

**Table 3D-7: Area of cultivated crops. See:**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Table 3D-8a-d: Number of treatments. See:**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Table 3D-9: Activity data for field burning of agricultural wastes. See:**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

**Table 3D-10: Emissions of pollutants from field burning of agricultural wastes. See:**

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

#### References

COWI, 2000: Overdækning af gyllebeholdere og kommunernes tilsyn hermed – undersøgelsesrapport. Danish Forest and Nature Agency. December 2000. (In Danish).

## Annex 3E - Waste

Annex 3E-1:	Human cremation activity data, 1980-2013
Annex 3E-2:	Animal cremation activity data, 1980-2013
Annex 3E-3:	Emissions from human cremation, 1980-2013
Annex 3E-4:	Emissions from animal cremation, 1980-2013
Annex 3E-5:	Compost production activity data, 1985-2013
Annex 3E-6:	Emissions from composting, 1985-2013
Annex 3E-7:	Combusted biogas at biogas production plants activity data, 1994-2004
Annex 3E-8:	Combusted biogas at biogas production plants emissions, 1994-2004
Annex 3E-9:	Occurrence of all fires, building and vehicle fires, 1980-2013
Annex 3E-10:	Accidental building fires full scale equivalent activity data, 1980-2013
Annex 3E-11:	Emission factors for building fires, 1980-2013
Annex 3E-12:	Average building floor space, 1980-2013
Annex 3E-13:	Emissions from building fires, 1980-2013
Annex 3E-14:	Full scale vehicle fires, 1980-2013
Annex 3E-15:	Average vehicle weight, 1980-2013
Annex 3E-16:	Accidental vehicle fires activity data, 1980-2013
Annex 3E-17:	Emissions from accidental vehicle fires, 1980-2013

All annexes are available online at:

<http://envs.au.dk/videnudveksling/luft/emissioner/supporting-documentation/air-pollution-iir/>

## Annex 4 – Completeness and use of notation keys

### Not estimated categories

The Danish air emission inventory is generally complete. However, some categories and/or pollutants are reported as NE (Not estimated).

### Mobile combustion

PAH emissions from tire and brake wear are not estimated, due to lack of emission factors.

### Industrial processes

- Emissions from quarrying and mining of minerals other than coal have not been estimated.
- Emissions from construction and demolition are not estimated.
- Emissions from storage, handling and transport of mineral products are not estimated
- Emissions from storage, handling and transport of chemical products have not been estimated.
- Emissions from storage, handling and transport of metal products have not been estimated.
- Emissions from pulp and paper production have not been estimated.
- Emissions from wood processing have not been estimated.
- Emissions from production of POPs have not been estimated due to lack of emission factors.
- Emissions from consumption of POPs and heavy metals have not been estimated.
- Emissions of PM, BC and PAH from road paving with asphalt and asphalt roofing have not been estimated.
- Emissions of PM from food and beverages industry have not been estimated.
- Emissions from some product uses have not been estimated, e.g. use of shoes.

### Agriculture

Emissions of PM from off-farm storage, handling and transport of bulk agricultural products have not been estimated.

Emissions of NMVOC from the use of pesticides have not been estimated.

### Waste

Emissions from solid waste disposal on land have not been estimated due to lack of resources.

Emissions from wastewater handling have not been estimated due to lack of resources.

Emissions from small scale waste burning have not been estimated due to lack of resources.

The emission of selenium and HCB from accidental fires has not been estimated due to lack of available emission factors.

### **Categories reported as IE (Included Elsewhere)**

The table below indicates the categories where the notation key IE has been used in the reporting for some or all pollutants.

Table A3.1 List of categories reported as included elsewhere.

Category reported as IE	Emissions where emissions are included
2A1 Cement production	1A2f Manufacturing industries and construction: Non-metallic minerals
2A2 Lime production	1A2f Manufacturing industries and construction: Non-metallic minerals
2A3 Glass production	1A2f Manufacturing industries and construction: Non-metallic minerals

Emissions from cement production (2A1), lime production (2A2) and glass production (2A3) are included in manufacturing industries and construction (1A2f). For some or all pollutants, it is not possible to separate the process emissions from the energy related emissions.

For some pollutants in other categories IE is also used. The specific reasons are explained in the sectoral chapters of the report.



*[Blank page]*

# ANNUAL DANISH INFORMATIVE INVENTORY REPORT TO UNECE

Emission inventories from the base year of the protocols to  
year 2013

This report is a documentation report on the emission inventories for Denmark as reported to the UNECE Secretariat under the Convention on Long Range Transboundary Air Pollution due by 15 February 2015. The report contains information on Denmark's emission inventories regarding emissions of (1) SO<sub>x</sub> for the years 1980-2013, (2) NO<sub>x</sub>, CO, NMVOC and NH<sub>3</sub> for the years 1985-2013, (3) Particulate matter: TSP, PM<sub>10</sub>, PM<sub>2.5</sub> for the years 2000-2013, (4) Heavy Metals: Pb, Cd, Hg, As, Cr, Cu, Ni, Se and Zn for the years 1990-2013, (5) Polycyclic aromatic hydrocarbons (PAH): Benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene and indeno(1,2,3-cd)pyrene, PCDD/F and HCB for the years 1990-2013. Further, the report contains information on background data for emissions inventory.